




Australian Government
**Department of Agriculture
and Water Resources**

Final report for the non-regulated analysis of existing policy for table grapes from India

July 2016



© Commonwealth of Australia 2016

Ownership of intellectual property rights

Unless otherwise noted, copyright (and any other intellectual property rights, if any) in this publication is owned by the Commonwealth of Australia (referred to as the Commonwealth).

Creative Commons licence

All material in this publication is licensed under a Creative Commons Attribution 3.0 Australia Licence, save for content supplied by third parties, photographic images, logos and the Commonwealth Coat of Arms.



Creative Commons Attribution 3.0 Australia Licence is a standard form licence agreement that allows you to copy, distribute, transmit and adapt this publication provided you attribute the work. A summary of the licence terms is available from creativecommons.org/licenses/by/3.0/au/deed.en. The full licence terms are available from creativecommons.org/licenses/by/3.0/au/legalcode.

Inquiries about the licence and any use of this document should be sent to copyright@agriculture.gov.au.

This publication (and any material sourced from it) should be attributed as: Australian Government Department of Agriculture and Water Resources 2016, *Final report for the non-regulated analysis of existing policy for table grapes from India*. CC BY 3.0

Cataloguing data

Australian Government Department of Agriculture and Water Resources 2016, *Final report for the non-regulated analysis of existing policy for table grapes from India*, the Australian Government Department of Agriculture and Water Resources, Canberra.

This publication is available at agriculture.gov.au.

Australian Government Department of Agriculture and Water Resources

Postal address: GPO Box 858 Canberra ACT 2601

Switchboard: +61 2 6272 3933 or 1800 900 090

Facsimile: +61 2 6272 3307

Email: plant@agriculture.gov.au

Liability

The Australian Government acting through the Australian Government Department of Agriculture and Water Resources has exercised due care and skill in preparing and compiling the information in this publication. Notwithstanding, the Australian Government Department of Agriculture and Water Resources, its employees and advisers disclaim all liability, including liability for negligence and for any loss, damage, injury, expense or cost incurred by any person as a result of accessing, using or relying upon any of the information or data in this publication to the maximum extent permitted by law.

Contents

| | |
|---|-------------|
| Acronyms and abbreviations..... | viii |
| Summary | 1 |
| 1 Introduction | 3 |
| 1.1 Australia’s biosecurity policy framework..... | 3 |
| 1.2 This import risk analysis..... | 3 |
| 2 Method for pest risk analysis | 6 |
| 2.1 Stage 1 Initiation..... | 6 |
| 2.2 Stage 2 Pest risk assessment | 7 |
| 2.3 Stage 3 Pest risk management..... | 14 |
| 3 India’s commercial production practices for table grapes | 16 |
| 3.1 Assumptions used in estimating unrestricted risk | 16 |
| 3.2 India’s table grape production areas | 16 |
| 3.3 Climate in production areas..... | 16 |
| 3.4 Pre-harvest | 21 |
| 3.5 Harvesting and handling procedures..... | 27 |
| 3.6 Post-harvest | 27 |
| 3.7 Export capability | 32 |
| 4 Pest risk assessments for quarantine pests..... | 34 |
| Assessments for quarantine pests for which a full pest risk assessment is conducted | 38 |
| 4.1 Grapevine bacterial canker disease..... | 38 |
| 4.2 Tomato black ring virus..... | 47 |
| Assessments for quarantine pests for which the URE outcome is adopted from previous assessments | 58 |
| 4.3 Grapevine phylloxera | 58 |
| 4.4 European fruit lecanium | 59 |
| 4.5 Mealybugs | 60 |
| 4.6 Large grape plume moth..... | 61 |
| 4.7 Thrips..... | 62 |
| 4.8 Kanzawa spider mite | 63 |
| 4.9 Black rot..... | 64 |
| 4.10 Brown rot | 65 |
| 4.11 Grapevine leaf rust | 66 |
| 4.12 Phomopsis cane and leaf spot..... | 67 |
| Assessments for quarantine pests for which some of the likelihood and/or consequence ratings are adopted from previous assessments..... | 68 |

| | | |
|-------------------|--|------------|
| 4.13 | Fruit flies..... | 68 |
| 4.14 | Leaf rolling moth..... | 72 |
| 4.15 | Citrus pyralid..... | 76 |
| 4.16 | Bitter rot..... | 80 |
| 4.17 | Fruit rot..... | 86 |
| 4.18 | White rot..... | 92 |
| 4.19 | Tobacco necrosis viruses..... | 98 |
| | Assessments for quarantine pests for which the likelihood and consequence ratings have been determined in a previous assessment..... | 103 |
| 4.20 | Spotted wing drosophila..... | 103 |
| 4.21 | Pest risk assessment conclusions..... | 105 |
| 5 | Pest risk management..... | 111 |
| 5.1 | Pest risk management measures and phytosanitary procedures..... | 111 |
| 5.2 | Operational system for the maintenance and verification of phytosanitary status..... | 119 |
| 5.3 | Responsibility of competent authority..... | 123 |
| 5.4 | Uncategorised pests..... | 124 |
| 5.5 | Review of processes..... | 124 |
| 5.6 | Meeting Australia's food laws..... | 124 |
| 6 | Conclusion..... | 126 |
| Appendix A | Initiation and categorisation for pests of table grapes from India..... | 127 |
| Appendix B | Issues raised in stakeholder comments..... | 206 |
| | Glossary..... | 209 |
| | References..... | 213 |

Figures

| | | |
|----------|--|-----|
| Figure 1 | Diagram of grapes..... | vii |
| Figure 2 | Monthly mean maximum and minimum temperatures and monthly mean rainfall in the table grape producing districts of Nashik, Pune, Gulberga, Bangalore, Anantapur, Hyderabad, Madurai, Hisar and Bhatinda in India..... | 19 |
| Figure 3 | A grape bunch covered with a paper bag..... | 22 |
| Figure 4 | Grapes being trimmed in the packing house..... | 28 |
| Figure 5 | Grapes being packed in the packing house..... | 28 |
| Figure 6 | Examples of labels showing trace back information on the carton..... | 29 |
| Figure 7 | Summary of vineyard and post-harvest processes practiced in India, for table grapes for export..... | 31 |

Tables

| | | |
|-----------|--|----|
| Table 2.1 | Nomenclature of qualitative likelihoods..... | 10 |
|-----------|--|----|

| | |
|--|-----|
| Table 2.2 Matrix of rules for combining qualitative likelihoods | 11 |
| Table 2.3 Decision rules for determining the consequence impact score based on the magnitude of consequences at four geographic scales..... | 13 |
| Table 2.4 Decision rules for determining the overall consequence rating for each pest..... | 13 |
| Table 2.5 Risk estimation matrix..... | 14 |
| Table 3.1 Pests and their management measures for grapevine in India | 24 |
| Table 3.2 Production and per cent share of total grape production in India in 2013–14 | 32 |
| Table 3.3 Export volumes of fresh grapes from India to the top five export markets from 2010 to 2014..... | 32 |
| Table 4.1 Quarantine pests for table grapes from India for which a full pest risk assessment is conducted..... | 36 |
| Table 4.2 Quarantine pests for table grapes from India for which the URE outcome is adopted from previous assessments..... | 36 |
| Table 4.3 Quarantine pests for table grapes from India for which some of the likelihood and/or consequence ratings are adopted from previous assessments | 37 |
| Table 4.4 Quarantine pests for table grapes from India for which the likelihood and consequence ratings have been determined in a previous assessment..... | 37 |
| Table 4.5 Summary of unrestricted risk estimates for quarantine pests associated with table grapes from India for which a full pest risk assessment is conducted | 106 |
| Table 4.6 Summary of unrestricted risk estimates for quarantine pests associated with table grapes from India for which the URE outcome is adopted from previous assessments..... | 106 |
| Table 4.7 Summary of unrestricted risk estimates for quarantine pests associated with table grapes from India for which some of the likelihood and/or consequence ratings are adopted from previous assessments..... | 108 |
| Table 4.8 Summary of unrestricted risk estimates for quarantine pests associated with table grapes from India for which the likelihood and consequence ratings have been determined in a previous assessment..... | 109 |
| Table 5.1 Phytosanitary measures recommended for quarantine pests for table grapes from India..... | 112 |

Maps

| | |
|--|----|
| Map 1 Map of Australia..... | vi |
| Map 2 A guide to Australia’s bio-climatic zones..... | vi |
| Map 3 Main table grape production areas in India | 18 |

Map 1 Map of Australia



Map 2 A guide to Australia's bio-climatic zones





Figure 1 Diagram of grapes

A shows the main parts of a grape cluster, B shows detail of the berry attachment



Source: Pratt (1988)

Acronyms and abbreviations

| Term or abbreviation | Definition |
|----------------------|---|
| ACT | Australian Capital Territory |
| ALOP | Appropriate level of protection |
| CABI | CAB International, Wallingford, UK |
| CSIRO | Commonwealth Scientific and Industrial Research Organisation |
| DAC | India's Department of Agriculture and Cooperation |
| DAFF | Acronym of the former Australian Government Department of Agriculture, Fisheries and Forestry, which is now the Australian Government Department of Agriculture and Water Resources |
| DAFWA | Government Department of Agriculture and Food, Western Australia |
| DPP | Directorate of Plant Protection, Ministry of Agriculture, India |
| EP | Existing policy |
| EPPO | European and Mediterranean Plant Protection Organization |
| FAO | Food and Agriculture Organization of the United Nations |
| ICON | The Australian Government Department of Agriculture and Water Resources Import Conditions database |
| IPC | International Phytosanitary Certificate |
| IPPC | International Plant Protection Convention |
| IRA | Import risk analysis |
| ISPM | International Standard for Phytosanitary Measures |
| NSW | New South Wales |
| NPPO | National Plant Protection Organisation |
| NT | Northern Territory |
| PIRSA | Department of Primary Industries and Regions of South Australia |
| PRA | Pest risk analysis |
| Qld | Queensland |
| SA | South Australia |
| SPS | Sanitary and Phytosanitary |
| Tas. | Tasmania |
| USA | The United States of America |
| Vic. | Victoria |
| WA | Western Australia |
| WTO | World Trade Organization |

Summary

The Australian Government Department of Agriculture and Water Resources has prepared this final report to assess the proposal by India for market access to Australia for table grapes.

Australia has existing policy for the import of table grapes for human consumption from Chile, the United States of America (California), New Zealand, the People's Republic of China, the Republic of Korea and Japan.

This final report recommends that the importation of table grapes from all commercial production areas of India be permitted, subject to a range of quarantine conditions.

Written submissions on the draft report were received from five stakeholders. This final report takes into account stakeholder comments on the draft report.

This final report identifies pests that require phytosanitary measures to manage risks to a very low level in order to achieve Australia's appropriate level of protection (ALOP). Fifteen pests were identified as requiring phytosanitary measures. Out of these 15 pests, 12 are arthropods and three are pathogens.

The 12 arthropod pests requiring measures are: *Planococcus ficus* (grapevine mealybug), *Planococcus lilanicus* (coffee mealybug), *Planococcus minor* (Pacific mealybug), *Rastrococcus iceryoides* (Downey snowline mealybug), *Tetranychus kanzawai* (Kanzawa spider mite), *Archips machlopi* (leaf rolling moth), *Retithrips syriacus* (black vine thrips), *Rhipiphorothrips cruentatus* (grapevine thrips), *Bactrocera correcta* (guava fruit fly), *Bactrocera dorsalis* (oriental fruit fly), *Drosophila suzukii* (spotted wing drosophila) and *Daktulosphaira vitifoliae* (grapevine phylloxera).

The three pathogen pests requiring measures are: *Guignardia bidwellii* (black rot), *Monilinia fructigena* (brown rot) and *Phakopsora euvitis* (grapevine leaf rust).

The recommended phytosanitary measures take account of regional differences within Australia. One arthropod pest requiring measures, Kanzawa spider mite, has been identified as a quarantine pest for Western Australia.

This final report recommends a range of risk management measures, combined with a system of operational procedures to ensure quarantine standards are met. These measures will reduce the risk posed by the 15 quarantine pests, and achieve Australia's ALOP. These measures include:

- visual inspection and, if detected, remedial action for the mealybugs, spider mite, leaf rolling moth and thrips
- area freedom or fruit treatment (cold disinfestation or irradiation) for fruit flies
- area freedom, fruit treatment (irradiation, methyl bromide fumigation or combined sulphur dioxide/carbon dioxide fumigation followed by cold disinfestation) or a systems approach approved by the Australian Government Department of Agriculture and Water Resources for spotted wing drosophila

- area freedom or fruit treatment (sulphur pad or combined sulphur dioxide/carbon dioxide fumigation) for grapevine phylloxera
- area freedom or a systems approach approved by the Australian Government Department of Agriculture and Water Resources for black rot, brown rot and grapevine leaf rust
- a supporting operational system to maintain and verify the phytosanitary status of export consignments.

The Australian Government Department of Agriculture and Water Resources has made some changes to the risk analysis following consideration of stakeholder comments on the draft report and subsequent review of literature. These changes include:

- the addition of two pests, *Cryptoblabes gnidiella* (Citrus pyralid) and *Strawberry latent ringspot virus*, in the pest categorisation (Appendix A)
- the addition of a pest risk assessment for one of these two pests, Citrus pyralid (the unrestricted risk estimate for this pest achieves Australia's ALOP and risk management measures are not required)
- additional text in the pest categorisation regarding the potential establishment of seed-borne pathogens through infected grapevine seedlings
- the addition of Appendix B Issues raised in stakeholder comments, which summarises key stakeholder comments and how they were considered in the final report
- minor corrections, rewording and editorial changes for consistency, clarity and web accessibility.

1 Introduction

1.1 Australia's biosecurity policy framework

Australia's biosecurity policies aim to protect Australia against the risks that may arise from exotic pests entering, establishing and spreading in Australia, thereby threatening Australia's unique flora and fauna, as well as those agricultural industries that are relatively free from serious pests.

The risk analysis process is an important part of Australia's biosecurity policies. It enables the Australian Government to formally consider the risks that could be associated with proposals to import new products into Australia. If the risks are found to exceed Australia's appropriate level of protection (ALOP), risk management measures are proposed to reduce the risks to an acceptable level. But, if it is not possible to reduce the risks to an acceptable level, then no trade will be allowed.

Successive Australian Governments have maintained a stringent, but not a zero risk, approach to the management of biosecurity risks. This approach is expressed in terms of Australia's ALOP, which reflects community expectations through government policy and is currently described as providing a high level of protection aimed at reducing risk to a very low level, but not to zero.

Australia's risk analyses are undertaken by the Australian Government Department of Agriculture and Water Resources (the department) using technical and scientific experts in relevant fields, and involve consultation with stakeholders at various stages during the process.

The department's assessment may take the form of a biosecurity import risk analysis (BIRA) or a non-regulated risk analysis (such as scientific review of existing policy and import conditions).

Further information about Australia's biosecurity framework is provided in the *Biosecurity Import Risk Analysis Guidelines 2016* located on the [Australian Government Department of Agriculture and Water Resources](#) website.

1.2 This import risk analysis

1.2.1 Background

The Department of Agriculture and Cooperation of India formally requested market access for table grapes to Australia in a submission received in 2007 (DPP 2007). This submission included information on the pests associated with table grape crops in India, including the plant part affected, and the standard commercial production practices for table grapes in India (DPP 2007).

In February 2008, India advised that market access for table grapes was its top priority. Additional production and pest information were received from India in 2009 (DPP 2009) and 2012 (DPP 2012).

On 26 November 2010, the department formally announced the commencement of this risk analysis, advising that it would be progressed as a non-regulated review of existing policy.

Officers from the department visited table grape production areas in Maharashtra, one of India's major table grape producing states, in April 2010 and in November 2015 to observe production systems and packinghouse operations.

1.2.2 Scope

The scope of this risk analysis is to consider the biosecurity risk that may be associated with the importation of commercially produced table grapes (*Vitis vinifera* and hybrids) (henceforth these will be referred to as table grapes) from India, for human consumption in Australia.

In this risk analysis, table grapes are defined as table grape bunches or clusters, which include peduncles, rachises, laterals, pedicels and berries (Pratt 1988), but not other plant parts (Figure 1). This risk analysis covers all commercially produced table grapes from all table grape producing states of India.

1.2.3 Existing policy

International policy

Import policy exists for table grapes from the United States of America (California) (AQIS 1999, 2000; Biosecurity Australia 2006a; DAFF 2013), Chile (Biosecurity Australia 2005b), New Zealand (Department of Agriculture 2013a), the People's Republic of China (Biosecurity Australia 2011a), the Republic of Korea (Biosecurity Australia 2011b) and Japan (Department of Agriculture 2014).

The [import requirements](#) for these commodity pathways can be found at the department's website. The department has considered all the pests previously identified in the existing policies and where relevant, the information in these assessments has been taken into account in this risk analysis.

Domestic arrangements

The Australian Government is responsible for regulating the movement of plants and plant products into and out of Australia. However, the state and territory governments are responsible for plant health controls within their individual jurisdiction. Legislation relating to resource management or plant health may be used by state and territory government agencies to control interstate movement of plants and their products. Once plant and plant products have been cleared by Australian biosecurity officers, they may be subject to interstate movement conditions. It is the importer's responsibility to identify, and ensure compliance with all requirements.

Under Western Australia legislation, grape fruit, seeds and plants and machinery used in the growing or processing of grapes are prescribed potential carriers of various declared pests and are restricted entry into Western Australia. Import permits may be issued for the entry of grape plants and propagative material subject to post entry quarantine requirements.

On 15 September 2011, the Government Department of Agriculture and Food, Western Australia (DAFWA) announced the formal commencement of a pest risk analysis considering the importation of table grapes into Western Australia from other Australian states and territories. In June 2015, DAWFA released a draft report for this pest risk analysis for stakeholder

consultation until 1 August 2015 (DAFWA 2015b, a). On 16 October 2015, DAFWA released the final report for this pest risk analysis (DAFWA 2015d, c).

1.2.4 Contaminating pests

In addition to the pests associated with table grapes from India that are assessed in this risk analysis, there are other organisms that may arrive with the imported commodity. These organisms could include pests of other crops or predators and parasitoids of other arthropods. The department considers these organisms to be contaminating pests that could pose sanitary and phytosanitary risks. These risks are addressed by existing operational procedures that require a 600 unit inspection of all consignments, or equivalent, and investigation of any pest that may be of quarantine concern to Australia.

1.2.5 Consultation

On 26 November 2010, the department notified stakeholders in Biosecurity Advice 2010/37 of the formal commencement of a non-regulated analysis of existing policy to consider a proposal from India for market access to Australia for table grapes.

The department has regularly consulted with India's DAC and Australian state and territory government departments during the preparation of this final report.

The department provided a draft pest categorisation to Australian state and territory government departments on 29 October 2012 for their advance consideration of regional pests, prior to the formal release of the draft report.

The draft report was released on 22 July 2015 (Biosecurity Advice 2015/09) for comment and consultation with stakeholders, for a period of 30 days that concluded on 21 August 2015. The department received five submissions on the draft report. All submissions were carefully considered and, where relevant, changes were made to the final report. A summary of major stakeholder comments and how they were considered is contained in Appendix B.

2 Method for pest risk analysis

This chapter sets out the method used for the pest risk analysis (PRA) in this report. The department has conducted this PRA in accordance with the International Standards for Phytosanitary Measures (ISPMs), including ISPM 2: *Framework for pest risk analysis* (FAO 2007b) and ISPM 11: *Pest risk analysis for quarantine pests* (FAO 2013) that have been developed under the SPS Agreement (WTO 1995).

A PRA is 'the process of evaluating biological or other scientific and economic evidence to determine whether an organism is a pest, whether it should be regulated, and the strength of any phytosanitary measures to be taken against it' (FAO 2015a). A pest is 'any species, strain or biotype of plant, animal, or pathogenic agent injurious to plants or plant products' (FAO 2015a).

Quarantine risk consists of two major components: the probability of a pest entering, establishing and spreading in Australia from imports; and the consequences should this happen. These two components are combined to give an overall estimate of the risk.

Unrestricted risk is estimated taking into account the existing commercial production practices of the exporting country and that, on arrival in Australia, the department will verify that the consignment received is as described on the commercial documents and its integrity has been maintained.

Restricted risk is estimated with phytosanitary measure(s) applied. A phytosanitary measure is 'any legislation, regulation or official procedure having the purpose to prevent the introduction and/or spread of quarantine pests, or to limit the economic impact of regulated non-quarantine pests' (FAO 2015a).

A glossary of the terms used is provided at the back of this report.

The PRAs are conducted in the following three consecutive stages: initiation, pest risk assessment and pest risk management.

2.1 Stage 1 Initiation

Initiation identifies the pest(s) and pathway(s) that are of quarantine concern and should be considered for risk analysis in relation to the identified PRA area.

Appendix A of this risk analysis report lists the pests with the potential to be associated with the exported commodity produced using commercial production and packing procedures. Appendix A does not present a comprehensive list of all the pests associated with the entire plant, but concentrates on the pests that could be on the assessed commodity. Contaminating pests that have no specific relation to the commodity or the export pathway have not been listed and would be addressed by Australia's current approach to contaminating pests.

The identity of the pests is given in Appendix A. The species name is used in most instances but a lower taxonomic level is used where appropriate. Synonyms are provided where the current scientific name differs from that provided by the exporting country's National Plant Protection Organisation (NPPO) or where the cited literature used a different scientific name.

For this risk analysis, the 'PRA area' is defined as Australia for pests that are absent, or of limited distribution and under official control. For areas with regional freedom from a pest, the 'PRA area' may be defined on the basis of a state or territory of Australia or may be defined as a region of Australia consisting of parts of a state or territory or several states or territories.

For pests that had been considered by the department in other risk assessments and for which import policies already exist, a judgement was made on the likelihood of entry of pests on the commodity and whether existing policy is adequate to manage the risks associated with its import. Where appropriate, the previous risk assessment was taken into consideration when developing the new policy.

2.2 Stage 2 Pest risk assessment

A pest risk assessment (for quarantine pests) is the 'evaluation of the probability of the introduction and spread of a pest and of the magnitude of the associated potential economic consequences' (FAO 2015a).

The following three, consecutive steps were used in pest risk assessment:

2.2.1 Pest categorisation

Pest categorisation identifies which of the pests with the potential to be on the commodity are quarantine pests for Australia and require pest risk assessment. A 'quarantine pest' is a pest of potential economic importance to the area endangered thereby and not yet present there, or present but not widely distributed and being officially controlled (FAO 2015a).

The pests identified in Stage 1 were categorised using the following primary elements to identify the quarantine pests for the commodity being assessed:

- identity of the pest
- presence or absence in the PRA area
- regulatory status
- potential for establishment and spread in the PRA area
- potential for economic consequences (including environmental consequences) in the PRA area.

The results of pest categorisation are set out in Appendix A. The quarantine pests identified during categorisation were carried forward for pest risk assessment and are listed in Tables 4.1 to 4.4.

2.2.2 Assessment of the probability of entry, establishment and spread

Details of how to assess the 'probability of entry', 'probability of establishment' and 'probability of spread' of a pest are given in ISPM 11 (FAO 2013). A summary of this process is given below, followed by a description of the qualitative methodology used in this risk analysis.

Probability of entry

The probability of entry describes the probability that a quarantine pest will enter Australia as a result of trade in a given commodity, be distributed in a viable state in the PRA area and subsequently be transferred to a host. It is based on pathway scenarios depicting necessary steps in the sourcing of the commodity for export, its processing, transport and storage, its use in Australia and the generation and disposal of waste. In particular, the ability of the pest to survive is considered for each of these various stages.

The probability of entry estimates for the quarantine pests for a commodity are based on the use of the existing commercial production, packaging and shipping practices of the exporting country. Details of the existing commercial production practices for the commodity are set out in Chapter 3. These practices are taken into consideration by the department when estimating the probability of entry.

For the purpose of considering the probability of entry, the department divides this step into two components:

- **Probability of importation**—the probability that a pest will arrive in Australia when a given commodity is imported.
- **Probability of distribution**—the probability that the pest will be distributed, as a result of the processing, sale or disposal of the commodity, in the PRA area and subsequently transfer to a susceptible part of a host.

Factors considered in the probability of importation include:

- distribution and incidence of the pest in the source area
- occurrence of the pest in a life-stage that would be associated with the commodity
- mode of trade (for example, bulk, packed)
- volume and frequency of movement of the commodity along each pathway
- seasonal timing of imports
- pest management, cultural and commercial procedures applied at the place of origin
- speed of transport and conditions of storage compared with the duration of the lifecycle of the pest
- vulnerability of the life-stages of the pest during transport or storage
- incidence of the pest likely to be associated with a consignment
- commercial procedures (for example, refrigeration) applied to consignments during transport and storage in the country of origin, and during transport to Australia.

Factors considered in the probability of distribution include:

- commercial procedures (for example, refrigeration) applied to consignments during distribution in Australia
- dispersal mechanisms of the pest, including vectors, to allow movement from the pathway to a host

- whether the imported commodity is to be sent to a few or many destination points in the PRA area
- proximity of entry, transit and destination points to hosts
- time of year at which import takes place
- intended use of the commodity (for example, for planting, processing or consumption)
- risks from by-products and waste.

Probability of establishment

Establishment is defined as the 'perpetuation for the foreseeable future, of a pest within an area after entry' (FAO 2015a). In order to estimate the probability of establishment of a pest, reliable biological information (for example, lifecycle, host range, epidemiology, survival) is obtained from the areas where the pest currently occurs. The situation in the PRA area can then be compared with that in the areas where it currently occurs and expert judgement used to assess the probability of establishment.

Factors considered in the probability of establishment in the PRA area include:

- availability of hosts, alternative hosts and vectors
- suitability of the environment
- reproductive strategy and potential for adaptation
- minimum population needed for establishment
- cultural practices and control measures.

Probability of spread

Spread is defined as 'the expansion of the geographical distribution of a pest within an area' (FAO 2015a). The probability of spread considers the factors relevant to the movement of the pest, after establishment on a host plant or plants, to other susceptible host plants of the same or different species in other areas. In order to estimate the probability of spread of the pest, reliable biological information is obtained from areas where the pest currently occurs. The situation in the PRA area is then carefully compared with that in the areas where the pest currently occurs and expert judgement used to assess the probability of spread.

Factors considered in the probability of spread include:

- suitability of the natural and/or managed environment for natural spread of the pest
- presence of natural barriers
- potential for movement with commodities, conveyances or by vectors
- intended use of the commodity
- potential vectors of the pest in the PRA area
- potential natural enemies of the pest in the PRA area.

Assigning qualitative likelihoods for entry, establishment and spread

In its qualitative PRAs, the department uses the term 'likelihood' for the descriptors it uses for its estimates of probability of entry, establishment and spread. Qualitative likelihoods are assigned to each step of entry, establishment and spread. Six descriptors are used: high; moderate; low; very low; extremely low; and negligible (Table 2.1). Descriptive definitions for these descriptors and their indicative probability ranges are given in Table 2.1. The indicative probability ranges are only provided to illustrate the boundaries of the descriptors and are not used beyond this purpose in qualitative PRAs. These indicative probability ranges provide guidance to the risk analyst and promote consistency between different pest risk assessments.

Table 2.1 Nomenclature of qualitative likelihoods

| Likelihood | Descriptive definition | Indicative probability (P) range |
|-------------------|--|---|
| High | The event would be very likely to occur | $0.7 < P \leq 1$ |
| Moderate | The event would occur with an even probability | $0.3 < P \leq 0.7$ |
| Low | The event would be unlikely to occur | $0.05 < P \leq 0.3$ |
| Very low | The event would be very unlikely to occur | $0.001 < P \leq 0.05$ |
| Extremely low | The event would be extremely unlikely to occur | $0.000001 < P \leq 0.001$ |
| Negligible | The event would almost certainly not occur | $0 < P \leq 0.000001$ |

Combining likelihoods

The likelihood of entry is determined by combining the likelihood that the pest will be imported into the PRA area and the likelihood that the pest will be distributed within the PRA area, using a matrix of rules (Table 2.2). This matrix is then used to combine the likelihood of entry and the likelihood of establishment, and the likelihood of entry and establishment is then combined with the likelihood of spread to determine the overall likelihood of entry, establishment and spread.

For example, if the likelihood of importation is assigned a descriptor of 'low' and the likelihood of distribution is assigned a descriptor of 'moderate', then they are combined to give a likelihood of 'low' for entry. The likelihood for entry is then combined with the likelihood assigned for establishment of 'high' to give a likelihood for entry and establishment of 'low'. The likelihood for entry and establishment is then combined with the likelihood assigned for spread of 'very low' to give the overall likelihood for entry, establishment and spread of 'very low'. This can be summarised as:

| | |
|--|----------------------------------|
| importation x distribution = entry [E] | low x moderate = low |
| entry x establishment = [EE] | low x high = low |
| [EE] x spread = [EES] | low x very low = very low |

Table 2.2 Matrix of rules for combining qualitative likelihoods

| | High | Moderate | Low | Very low | Extremely low | Negligible |
|---------------|------|----------|----------|---------------|---------------|------------|
| High | High | Moderate | Low | Very low | Extremely low | Negligible |
| Moderate | | Low | Low | Very low | Extremely low | Negligible |
| Low | | | Very low | Very low | Extremely low | Negligible |
| Very low | | | | Extremely low | Extremely low | Negligible |
| Extremely low | | | | | Negligible | Negligible |
| Negligible | | | | | | Negligible |

Time and volume of trade

One factor affecting the likelihood of entry is the volume and duration of trade. If all other conditions remain the same, the overall likelihood of entry will increase as time passes and the overall volume of trade increases.

The department normally considers the likelihood of entry on the basis of the estimated volume of one year's trade. This is a convenient value for the analysis that is relatively easy to estimate and allows for expert consideration of seasonal variations in pest presence, incidence and behaviour to be taken into account. The consideration of the likelihood of entry, establishment and spread and subsequent consequences takes into account events that might happen over a number of years even though only one year's volume of trade is being considered. This difference reflects biological and ecological facts, for example where a pest or disease may establish in the year of import but spread may take many years.

The use of a one year volume of trade has been taken into account when setting up the matrix that is used to estimate the risk and therefore any policy based on this analysis does not simply apply to one year of trade. Policy decisions that are based on the department's method that uses the estimated volume of one year's trade are consistent with Australia's policy on appropriate level of protection and meet the Australian Government's requirement for ongoing quarantine protection. If there are substantial changes in the volume and nature of the trade in specific commodities then the department will review the risk analysis and, if necessary, provide updated policy advice.

In assessing the volume of trade in this risk analysis, the department assumed that a substantial volume of trade will occur.

2.2.3 Assessment of potential consequences

The objective of the consequence assessment is to provide a structured and transparent analysis of the likely consequences if the pests or disease agents were to enter, establish and spread in Australia. The assessment considers direct and indirect pest effects and their economic and environmental consequences. The requirements for assessing potential consequences are given in Article 5.3 of the SPS Agreement (WTO 1995), ISPM 5 (FAO 2015a) and ISPM 11 (FAO 2013).

Direct pest effects are considered in the context of the effects on:

- plant life or health

- other aspects of the environment.

Indirect pest effects are considered in the context of the effects on:

- eradication, control
- domestic trade
- international trade
- environment.

For each of these six criteria, the consequences were estimated over four geographic levels, defined as:

Local: an aggregate of households or enterprises (a rural community, a town or a local government area).

District: a geographically or geopolitically associated collection of aggregates (generally a recognised section of a state or territory, such as 'Far North Queensland').

Regional: a geographically or geopolitically associated collection of districts in a geographic area (generally a state or territory, although there may be exceptions with larger states such as Western Australia).

National: Australia wide (Australian mainland states and territories and Tasmania).

For each criterion, the magnitude of the potential consequence at each of these levels was described using four categories, defined as:

Indiscernible: pest impact unlikely to be noticeable.

Minor significance: expected to lead to a minor increase in mortality/morbidity of hosts or a minor decrease in production but not expected to threaten the economic viability of production. Expected to decrease the value of non-commercial criteria but not threaten the criterion's intrinsic value. Effects would generally be reversible.

Significant: expected to threaten the economic viability of production through a moderate increase in mortality/morbidity of hosts, or a moderate decrease in production. Expected to significantly diminish or threaten the intrinsic value of non-commercial criteria. Effects may not be reversible.

Major significance: expected to threaten the economic viability through a large increase in mortality/morbidity of hosts, or a large decrease in production. Expected to severely or irreversibly damage the intrinsic 'value' of non-commercial criteria.

The estimates of the magnitude of the potential consequences over the four geographic levels were translated into a qualitative impact score (A–G) using Table 2.3. For example, a consequence with a magnitude of 'significant' at the 'district' level will have a consequence impact score of D.

Table 2.3 Decision rules for determining the consequence impact score based on the magnitude of consequences at four geographic scales

| Magnitude | Geographic scale | | | |
|--------------------|------------------|----------|--------|--------|
| | Local | District | Region | Nation |
| Indiscernible | A | A | A | A |
| Minor significance | B | C | D | E |
| Significant | C | D | E | F |
| Major significance | D | E | F | G |

Note: In earlier qualitative PRAs, the scale for the impact scores went from A to F and did not explicitly allow for the rating 'indiscernible' at all four levels. This combination might be applicable for some criteria. In this report, the impact scale of A to F has been changed to become B–G and a new lowest category A ('indiscernible' at all four levels) was added. The rules for combining impacts in Table 2.4 were adjusted accordingly.

The overall consequence for each pest is achieved by combining the qualitative impact scores (A–G) for each direct and indirect consequence using a series of decision rules (Table 2.4). These rules are mutually exclusive, and are assessed in numerical order until one applies.

Table 2.4 Decision rules for determining the overall consequence rating for each pest

| Rule | The impact scores for consequences of direct and indirect criteria | Overall consequence rating |
|------|--|----------------------------|
| 1 | Any criterion has an impact of 'G'; or more than one criterion has an impact of 'F'; or a single criterion has an impact of 'F' and each remaining criterion an 'E'. | Extreme |
| 2 | A single criterion has an impact of 'F'; or all criteria have an impact of 'E'. | High |
| 3 | One or more criteria have an impact of 'E'; or all criteria have an impact of 'D'. | Moderate |
| 4 | One or more criteria have an impact of 'D'; or all criteria have an impact of 'C'. | Low |
| 5 | One or more criteria have an impact of 'C'; or all criteria have an impact of 'B'. | Very Low |
| 6 | One or more but not all criteria have an impact of 'B', and all remaining criteria have an impact of 'A'. | Negligible |

2.2.4 Estimation of the unrestricted risk

Once the assessment of the probability of entry, establishment and spread and for potential consequences is completed, the unrestricted risk can be determined for each pest or groups of pests. This is determined by using a risk estimation matrix (Table 2.5) to combine the estimates of the probability of entry, establishment and spread and the overall consequences of pest establishment and spread. Therefore, risk is the product of likelihood and consequence.

When interpreting the risk estimation matrix, note the descriptors for each axis are similar (for example, low, moderate, high) but the vertical axis refers to likelihood and the horizontal axis refers to consequences. Accordingly, a 'low' likelihood combined with 'high' consequences, is not the same as a 'high' likelihood combined with 'low' consequences—the matrix is not symmetrical. For example, the former combination would give an unrestricted risk rating of 'moderate', whereas, the latter would be rated as a 'low' unrestricted risk.

Table 2.5 Risk estimation matrix

| Likelihood of pest entry, establishment and spread | Consequences of pest entry, establishment and spread | | | | | |
|--|--|-----------------|-----------------|-----------------|-----------------|---------------|
| | Negligible | Very low | Low | Moderate | High | Extreme |
| High | Negligible risk | Very low risk | Low risk | Moderate risk | High risk | Extreme risk |
| Moderate | Negligible risk | Very low risk | Low risk | Moderate risk | High risk | Extreme risk |
| Low | Negligible risk | Negligible risk | Very low risk | Low risk | Moderate risk | High risk |
| Very low | Negligible risk | Negligible risk | Negligible risk | Very low risk | Low risk | Moderate risk |
| Extremely low | Negligible risk | Negligible risk | Negligible risk | Negligible risk | Very low risk | Low risk |
| Negligible | Negligible risk | Negligible risk | Negligible risk | Negligible risk | Negligible risk | Very low risk |

2.2.5 Australia's appropriate level of protection (ALOP)

The SPS Agreement defines the concept of an 'appropriate level of sanitary or phytosanitary protection (ALOP)' as the level of protection deemed appropriate by the WTO Member establishing a sanitary or phytosanitary measure to protect human, animal or plant life or health within its territory.

Like many other countries, Australia expresses its ALOP in qualitative terms. Australia's ALOP, which reflects community expectations through government policy, is currently expressed as providing a high level of sanitary or phytosanitary protection aimed at reducing risk to a very low level, but not to zero. The band of cells in Table 2.5 marked 'very low risk' represents Australia's ALOP.

2.3 Stage 3 Pest risk management

Pest risk management describes the process of identifying and implementing phytosanitary measures to manage risks to achieve Australia's ALOP, while ensuring that any negative effects on trade are minimised.

The conclusions from pest risk assessment are used to decide whether risk management is required and if so, the appropriate measures to be used. Where the unrestricted risk estimate exceeds Australia's ALOP, risk management measures are required to reduce this risk to a very low level. The guiding principle for risk management is to manage risk to achieve Australia's ALOP. The effectiveness of any proposed phytosanitary measures (or combination of measures) is evaluated, using the same approach as used to evaluate the unrestricted risk, to ensure it reduces the restricted risk for the relevant pest or pests to meet Australia's ALOP.

ISPM 11 (FAO 2013) provides details on the identification and selection of appropriate risk management options and notes that the choice of measures should be based on their effectiveness in reducing the probability of entry of the pest.

Examples given of measures commonly applied to traded commodities include:

- options for consignments—for example, inspection or testing for freedom from pests, prohibition of parts of the host, a pre-entry or post-entry quarantine system, specified conditions on preparation of the consignment, specified treatment of the consignment, restrictions on end-use, distribution and periods of entry of the commodity
- options preventing or reducing infestation in the crop—for example, treatment of the crop, restriction on the composition of a consignment so it is composed of plants belonging to resistant or less susceptible species, harvesting of plants at a certain age or specified time of the year, production in a certification scheme
- options ensuring that the area, place or site of production or crop is free from the pest—for example, pest-free area, pest-free place of production or pest-free production site
- options for other types of pathways—for example, consider natural spread, measures for human travellers and their baggage, cleaning or disinfestations of contaminated machinery
- options within the importing country—for example, surveillance and eradication programmes
- prohibition of commodities—if no satisfactory measure can be found.

Risk management measures are identified for each quarantine pest where the risk exceeds Australia's ALOP. These are presented in Chapter 5: Pest risk management, of this report.

3 India's commercial production practices for table grapes

This chapter provides information on the pre-harvest, harvest and post-harvest practices, considered to be standard practices in India for the production of table grapes for export. The export capability of India is also outlined.

3.1 Assumptions used in estimating unrestricted risk

India provided Australia with information on the standard commercial practices used in the production of table grapes in different regions and for all commercially produced table grape cultivars in India. This information was complemented with data from other sources and was taken into consideration when estimating the unrestricted risks of pests that may be associated with the import of this commodity.

Officers from the Australian Government Department of Agriculture and Water Resources (the department) visited table grape production areas in Maharashtra, one of India's major table grape producing states, in April 2010 and November 2015 to observe production systems and packinghouse operations. The department's observations and additional information provided during the visit confirmed the production and processing procedures described in this chapter as standard commercial production practices for table grapes for export.

In estimating the likelihood of pest introduction it was assumed that the pre-harvest, harvest and post-harvest production practices for table grapes as described in this chapter are implemented for all regions and for all table grape cultivars within the scope of this analysis. Where a specific practice described in this chapter is not taken into account to estimate the unrestricted risk, it is clearly identified and explained in Chapter 4.

3.2 India's table grape production areas

While grapes are grown across the sub-continent, commercially grown table grapes are mainly produced in the peninsular region of the country, as shown in Map 3 (DPP 2012; Shikhamany 2001). The major grape growing states are Maharashtra and Karnataka, accounting for about 95 per cent of India's total grape production. Other table grape growing states include Tamil Nadu, Telangana, Andhra Pradesh, Mizoram, Punjab, Madhya Pradesh, Jammu and Kashmir, Nagaland, Haryana and Rajasthan (APEDA 2015; DPP 2007, 2012). While most of the grapes are likely to be exported from the two major producing states, Maharashtra and Karnataka, India may also export grapes from other grape growing states.

3.3 Climate in production areas

Table grapes are grown under a variety of soil and climatic conditions in three distinct agro-climatic zones in India: sub-tropical, hot tropical and mild tropical climatic regions (DPP 2012). Climate data, including mean maximum and minimum temperatures, and mean rainfall for each of the production regions are presented in Figure 2.

3.3.1 Sub-tropical region

This region covers the north-western plains and lies between the latitudes of 28 and 32 degrees north, including Haryana (Hissar and Jind districts) and Punjab (Bathinda, Ferozpur, Gurdaspur and Ludhiana districts) (DPP 2012; Shikhamany 2001). Winter temperatures in this region rarely go below 0 degrees Celsius. Vines are dormant over winter and bud break starts in the first week of March. With rain arriving in the first week of June, only 90–95 days are available from the initiation of growth to harvest. Single pruning and a single harvest is the accepted practice here (DPP 2012; Shikhamany 2001).

3.3.2 Hot tropical region

This region lies between the latitudes of 15 and 20 degrees north, covering Maharashtra (Nashik, Sangli, Solapur, Pune, Satara, Latur and Osmanabad districts), Andhra Pradesh and Telangana (Hyderabad, Ranga Reddy, Mahabubnagar, Anantapur and Medak districts) and northern Karnataka (Bijapur, Bagalkot, Belgaum and Gulberga districts). This is the major viticulture region accounting for 70 per cent of the grape producing areas in India as well as 70 per cent of the total harvest. Vines do not undergo dormancy. Maximum and minimum temperature is 42 and 8 degrees Celsius respectively. Double pruning and a single harvest is the general practice in this region (DPP 2012; Shikhamany 2001).

3.3.3 Mild tropical region

This region lies between the latitudes of 10 and 15 degrees north, covering Karnataka (Bangalore and Kolar districts), Andhra Pradesh (Chittoor district) and Tamil Nadu (Coimbatore, Madurai and Theni districts). Maximum temperatures in a year seldom exceed 36 degrees Celsius, while the minimum is about 12 degrees Celsius. Generally two crops are harvested in a year in this region (DPP 2012; Shikhamany 2001).


Map 3 Main table grape production areas in India (APEDA 2015) marked by a star (★) Note: in 2014, the state of Andhra Pradesh was split into two states, Andhra Pradesh and Telangana, both major table grape production areas, nomenclature and territorial boundaries may not necessarily reflect government policy




Source: Adapted from CIA (2013)

Figure 2 Monthly mean maximum (—◆—) and minimum (—■—) temperatures (degrees Celsius) and monthly mean rainfall (millimetres) (—▲—) in the table grape producing districts of Nashik, Pune, Gulberga, Bangalore, Anantapur, Hyderabad, Madurai, Hisar and Bhatinda in India, based on average monthly weather data from 1901 to 2000






Sub-tropical



Hot tropical



Mild tropical



Source: India Meteorological Department (2000).

3.4 Pre-harvest

3.4.1 Cultivars

Grapes were introduced into India from Persia around the 14th century. Grapes are harvested almost all year round. However, in the main commercial production areas of Maharashtra and Karnataka, grapes are harvested between mid-February and the end of April (DAFF 2010; DPP 2009). More than 20 grape cultivars are grown in India (APEDA 2015). More than half of the grape cultivars produced in India are seedless, with Thompson Seedless being the dominant cultivar (DPP 2012). The cultivars produced commercially include Anab-e-Shahi (white, seeded), Bangalore Blue (black, seeded, also known as Isabella), Beauty Seedless (bluish black, seedless), Bhokri (white, seeded), Flame Seedless (red, seedless), Gulabi (purple, seeded, also known as Muscat Hamburg), Perlette (white, seedless), Pusa Seedless (white, seedless), Sharad Seedless (black, seedless) and Thompson Seedless and its mutants (white, seedless) (APEDA 2015; DPP 2012).

Anab-e-Shahi, Sharad Seedless, Flame Seedless and Thompson Seedless and its mutants are grown in the hot tropical region of India. Perlette and Thompson Seedless are grown in the sub-tropical region. Bangalore Blue, Anab-e-Shahi, Gulabi, Bhokri and Thompson Seedless are grown in the mild tropical region (DPP 2012).

The total commercial production area for grapes is approximately 55 000 hectares, distributed mainly in the states of Maharashtra and Karnataka (DPP 2012). Approximately 85 per cent of total grape production is consumed fresh, both for domestic and export industries. The rest are used for raisins, juice and in wine making (DPP 2012).

The table grape cultivars India intends to export to Australia include Thompson Seedless and a mutant, Muscat, Bangalore Blue, Flame Seedless, Sharad Seedless, Anab-e-shahi and Perlette (DPP 2009).

3.4.2 Cultivation practices

Planting

Grapevines in India are planted by bed and furrow with spacing of 3 meters between rows and 2 meters between plants within a row. Several rootstocks are used in India, for example Dogridge, 110R, 99R, Ramsey and St. George (NRC 2013). The National Research Centre (NRC) for grapes has a disease-free rootstock planting programme. The NRC produces disease-free rootstocks on which they graft a desired cultivar. It takes 18 months to produce a grafted rootstock ready for planting (DAFF 2010).

Trellis systems

Two types of trellis system, T framed and Y framed, are used. The T framed trellis provides full canopy between the rows with drooping bunches. The benefits of this are minimal weed growth, cooler air and soil temperatures, minimal berry rot and drop and lower evaporation rates (DAFF 2010). The Y framed trellis promotes growth across the furrow and produces an open, incomplete canopy derived from vines trained to two horizontal cordons (bilateral cordon training). However, the canopy gap between rows can allow considerable sunlight onto the

vineyard floor, which is conducive to weed growth, particularly under the drip line on beds and in furrows (DAFF 2010).

Pruning

In the tropical areas of India, pruning is done in March-April (summer) and September-October (winter) (Chand, Patil & Kishun 1991; DAFF 2010). The summer pruning, called back pruning, leaves basal single buds on the shoots near the cordon. The winter pruning cuts the mature canes back to the knot on subcanes or to the 6-7 bud position if the cane is straight. This is called fruit pruning because it prepares the vine for bearing fruits (NRC 2013).

Bagging

In India, paper bags are used to cover the grape bunches (Figure 3) in commercial vineyards, mainly to protect bunches from sunburn but may also help with pest management. Immediately prior to harvest, the paper bags are removed from the bunches and deposited in the irrigation channels. The bags are cleared from the channels after harvest, prior to flood irrigation (DAFF 2010).

Figure 3 A grape bunch covered with a paper bag



Irrigation

Drip irrigation is used during the dry season. The drip line is usually run 0.3–1.2 metres above the ground. One vineyard visited by officers from the department used a single line system hung at about 30–50 centimetres above the trunk base. This vineyard also used flood irrigation between harvest and pruning, which is generally done in October (DAFF 2010). Irrigation is scheduled as per advice received from the NRC which is based on evaluation of weather data (DAFF 2010).

3.4.3 Pest management

General pest surveillance and management programmes

India has a comprehensive pest monitoring and trace back system for table grapes for export to the European Union and other countries following the European Union food safety

requirements. The detailed processes are available in Procedures for export of fresh table grapes to the European Union (APEDA 2013). The Agricultural and Processed Food Products Export Development Authority (APEDA) is responsible for this system. GrapeNet is the internet based electronic system used by APEDA for quality assurance and trace back of all table grapes intended for export.

All farms intending to produce table grapes for export must be registered; this is done through the state agriculture departments. Each vineyard is given a unique registration number. The APEDA and the NRC for Grapes provide recommended spray and cultivation programmes to growers that comply with the European Union's maximum residue limits (MRL) for chemicals. The NRC for grapes also provides weather forecast based advice at three day intervals which predict the level of pest risk and recommend management strategies.

Pest and disease management programmes

Pest and disease management programmes in table grape vineyards use a broad range of practices including nutrient and water management, specific pruning and canopy management techniques and bagging. Biological and chemical control measures are also used. Table 3.1 shows the common pests and integrated pest management procedures for production of quality table grapes (NRC 2013).

Export vineyards need to maintain a record of practices that includes information on cultural practices and application of fertilizer and chemicals. Before harvest and chemical testing, each vineyard is inspected by the respective state's agriculture department (APEDA 2013). The state agriculture department will inspect the crop for quality, pest and disease incidence as well as verify the records (APEDA 2013). The growers not passing this inspection will not be qualified for export.

Before harvest, table grapes from vineyards registered for export are randomly sampled and tested for chemical residue by APEDA authorised laboratory workers (APEDA 2013). If the sample conforms to MRL requirements of the importing country, the grapes can be harvested for export.

Table 3.1 Pests and their management measures for grapevine in India (NRC 2013)

| Pest/pathogen | Common name | India's management measures |
|--|--------------------|---|
| <i>Elsinoe ampelina</i> | Anthracnose | <ul style="list-style-type: none"> • Remove all prunings • Sprays of carbendazim 50WP, ziram 27SL, chloronhalonil, COC 50WP, copper hydroxide 77WP, Bordeaux mixture, mancozeb, captan, difenconazole 25EC, fosetyl A1 and propineb 70WP at certain developmental times. |
| <i>Spodoptera litura</i> and others | Caterpillar | <ul style="list-style-type: none"> • Field sanitation • Summer ploughing • Regular scouting and monitoring • Light traps to catch moths • Pheromone traps for <i>Spodoptera litura</i> • Sprays of methomyl 40SP, emamectin benzoate 5SG and lambda cyhalothrin 5CS at certain developmental times. |
| <i>Plasmopara viticola</i> | downy mildew | <ul style="list-style-type: none"> • Prune after 15 October for fruit pruning • Sprays of Bordeaux mixture, copper fungicides, ziram, mancozeb, captan, chloronhalonil, sulphur 80WDG with Bordeaux mixture, fosetyl A1, propineb 70WP, metataxyl 8WP, cymoxanil, dimethomorph 50WP, fenamidone, azoxystrobin 23SC, iprovalicarb, famoxidone, metiram 60WG, pyraclostrobin and kresoxim methyl 44.3SC at certain developmental times. |
| <i>Rastrococcu iceryoides</i> , <i>Maconellicoccus hirsutus</i> and others | Mealybugs | <ul style="list-style-type: none"> • Field sanitation • Summer ploughing • Regular scouting and monitoring • Sprays of methomyl 40SP, bupprofezin 25SC and imidocloprid 17.8SL before veraison. |
| <i>Uncinula necator</i> | powdery mildew | <ul style="list-style-type: none"> • Remove all prunings • Sprays of sulphur 80WDG, potassium bicarbonate, Dinocap 48EC, hexaconazole, peconazole, flusilazole, myclobutanil, tetraconazole, azoxystrobin 23SC, kresoxim methyl 44.3SC, triademefon, mineral oils, fenarimol and difenoconazole at certain developmental times. |

| Pest/pathogen | Common name | India's management measures |
|---|---------------------|---|
| <i>Amrasca biguttula biguttula</i> | Leafhopper | <ul style="list-style-type: none"> • Field sanitation • Summer ploughing • Regular scouting and monitoring • Light traps • Sprays of fipronil 80WG, lambda cyhalothrin 5CS, imidacloprid 17.8SL and emamectin benzoate 5SG at certain developmental times. |
| <i>Phakopsora euvtis</i> | grapevine leaf rust | <ul style="list-style-type: none"> • Sprays of Bordeaux mixture, captafol, COC, copper hydroxide, difolatan, flusilazole 40EC, polycarbamate and prochloraz at certain developmental times • Use of resistant species and cultivars. |
| <i>Scelodonta strigicola</i> | flea beetle | <ul style="list-style-type: none"> • Field sanitation • Summer ploughing • Regular scouting and monitoring • Sprays of imidacloprid 17.8SL and lambda cyhalothrin 5SC at certain developmental times. |
| <i>Stromatium barbatum, Celosterna scrabrator</i> | stem borer | <ul style="list-style-type: none"> • Field sanitation • Summer ploughing • Regular scouting and monitoring • Make bore holes wider and try to hook the larvae out and kill them • Fine light traps for adults. |
| <i>Tetranychus kanzawai</i> | Kanzawa spider mite | <ul style="list-style-type: none"> • Field sanitation • Summer ploughing • Regular scouting and monitoring • If severe, use a jet spray of water before applying the chemical spray • Keep adequately irrigated • Sprays of sulphur 80WDG at certain developmental times. |

| Pest/pathogen | Common name | India's management measures |
|---|------------------------------------|--|
| <i>Tetranychus urticae</i> | red spider mite | <ul style="list-style-type: none"> • Field sanitation • Summer ploughing • Regular scouting and monitoring. • If severe, use a jet spray of water before applying the chemical spray • Keep adequately irrigated • Sprays of sulphur 80WDG at certain developmental times. |
| <i>Retithrips syriacus</i> , <i>Rhipiphorothrips cruentatus</i> , <i>Scirtothrips dorsalis</i> | Thrips | <ul style="list-style-type: none"> • Field sanitation • Summer ploughing • Regular scouting and monitoring • Sprays of fipronil 80WG, lambda cyhalothrin 5CS, imidacloprid 17.8SL and emamectin benzoate 5SG at certain developmental times. |
| <i>Xanthomonas campestris</i> pv. <i>viticola</i> | grapevine bacterial canker disease | <ul style="list-style-type: none"> • Copper fungicides can help to control this bacterial species. |

3.5 Harvesting and handling procedures

The grape harvest period in India may vary greatly. Nevertheless, in the main commercial production areas of Maharashtra and Karnataka, grapes are harvested between mid-February and the end of April (DAFF 2010; DPP 2009). Market preference for colour and sweetness determines harvest timing of each cultivar (DPP 2012).

Grape bunches for export typically carry 130 to 150 berries of 16 millimetres in diameter or more, with a brix level of 17–18 per cent. Each bunch can weigh up to 1 kilogram. The diameter is checked using a collapsible grape-sizing gauge, comprising a series of flat plastic fingers with punched holes, ranging from 16 to 29 millimetres. Grape berries for domestic consumption are smaller (less than 16 millimetres in diameter), sweeter and are left on the vine for longer to deepen in colour (DAFF 2010).

Bags (paper wrapping) are removed in the morning just before the grapes are harvested. The grapes are harvested manually using scissors (DAFF 2010; DPP 2009). The picker examines the bunches for berry size, colour, uniformity and any blemishes or damage. The harvested grape bunches are placed into field crates. Field crates used in India are stackable, low-walled, vented plastic crates, which are lined with a polyethylene foam insert. The dimensions of the field crates are 600 millimetres in length, 400 millimetres in width and 250 millimetres in height. The filled crates are either loaded on a small field trolley or on a large trailer and then loaded onto trucks in front of the property. The trucks are covered and sealed when full and the load is transported to the packing house for processing (DAFF 2010).

3.6 Post-harvest

3.6.1 Packing house

All packing houses wishing to process table grapes for export must be registered with APEDA. The packing house is inspected by the Packhouse Recognition Committee and if passed a packing house recognition certificate is issued.

Upon arrival at the packing house the harvested grape bunches are put into a receiving room, which is isolated from the grading room. The fruit is held in the receiving room for an hour to cool (DAFF 2010).

Sorting and trimming

After the grape bunches have cooled they are moved to the grading room. Post-harvest trimming of bunches is done to improve the appearance of the bunch and prevent postharvest decay. Trimming is performed with sharp, long-nosed scissors to remove defective berries (Figure 4). Berries that are too small, overripe, split, shrivelled, diseased, immature, or off-coloured are removed (by their pedicels) from the selected bunches. Bunches that are too compact, too straggly, sun scorched, damaged or containing excessive amounts of water are promptly removed for disposal (DAFF 2010; DPP 2012).

The bunches are graded for size and colour. Grade designation and quality of table grapes is specified under the *Fruits and vegetables grading and marking rules, 2004* (DAC 2004), amended in 2007 (DAC 2007), in Schedule II: Grade designation and quality of table grapes. Parameters

such as size, colour and Brix value are assessed and inspection made for visual diseases and defects (DPP 2009).

The graded bunches are put into clean crates and onto conveyor lines to move to the packing room (DPP 2009, 2012). A line supervisor inspects the graded product before it is packed.

Figure 4 Grapes being trimmed in the packing house



Packing

In the packing room, the bunches are trimmed to the desired weight and placed into 500 gram plastic punnets or polythene pouches. Each punnet or pouch contains two bunches weighing between 350 and 650 grams and are packed into 5 kilogram cardboard carton lined with a plastic bag or a 4 kilogram vented polystyrene carton (DAFF 2010; DPP 2012) (Figure 5).

Figure 5 Grapes being packed in the packing house





Before placing the punnets or pouches in a carton, bubble sheet is spread with its rough surface facing towards the base of the carton. A polythene liner is spread over the top of the bubble sheet. After pre-cooling, dual purpose sulphur dioxide releasing pads are placed over the punnets and the polythene liner is folded in.

Labelling

All cartons are labelled for trace back purposes using a unique identification code, identifying the packing house, vineyard and plot and the date packed. Cartons are also marked with information on the name of the commodity, variety, grade or fruit size, net weight, and any additional requirements specified by the importing country (DPP 2012) (Figure 6).

Figure 6 Examples of labels showing trace back information on the carton



Pre-cooling and storage

As soon as possible after packing and the phytosanitary checks, grapes are pre-cooled to less than 4 degrees Celsius by forced air. This pre-cooling is best if it occurs within 4–6 hours of harvest. The grapes are pre-cooled for 6-8 hours to reach a temperature of less than 4 degrees Celsius (DPP 2009, 2012; NRC 2013).

After pre-cooling, cartons are strapped and put into cold storage and held for shipment. Doors at the rear of each cool room open into a dispatch bay fitted with an insect zapper. Refrigerated shipping containers are backed up to the dispatch door for loading. Conditions are maintained at

0–2 degrees Celsius and 90–95 per cent humidity throughout storage and transport (DAFF 2010; DPP 2009, 2012). For air freight, there is no pre-cooling (DPP 2012). Figure 7 summarises the vineyard and post-harvest processes for Indian table grapes produced for export.


3.6.2 Export procedures

Cartons are randomly selected for inspection prior to storage. Quality control checks may be performed by packing house staff and phytosanitary checks for export certification are conducted by a district agriculture inspection officer from the National Horticulture Mission, which is a part of the Department of Agriculture and Cooperation, India. Once these checks are passed the palletised product is moved into pre-cooling.

3.6.3 Transport

Table grapes for export from India are transported by air or sea freight. When transported by sea freight, the grapes are in refrigerated containers, with temperature and humidity recorders (DPP 2009). The palletised boxes of grapes are loaded into a refrigerated container for shipment. Temperature and humidity recorders are placed inside the container and temperature sensors are calibrated prior to loading the container. When recommended temperature and humidity levels have been reached in the container, the pallets are arranged in an interlocking position to prevent movement within the container. The doors of the container are closed immediately after loading, and seals are affixed to the door lock (DPP 2012). For air freight, packed grapes are immediately loaded into refrigerated container trucks or vans and transported to the airport (DPP 2012).

Figure 7 Summary of vineyard and post-harvest processes practiced in India, for table grapes for export



3.7 Export capability

3.7.1 Production statistics

Approximately two and a half million tonnes of grapes are harvested annually in India. Table 3.2 shows the 2013–2014 grape production figures in India (APEDA 2015). Around 95 per cent of total production is from the two grape producing states, Maharashtra and Karnataka.

Table 3.2 Production and per cent share of total grape production in India in 2013–14 (APEDA 2015)

| State | Production in metric tonnes | Per cent share |
|-------------------|-----------------------------|----------------|
| Maharashtra | 2 160 000 | 83.55 |
| Karnataka | 302 390 | 11.70 |
| Tamil Nadu | 47 720 | 1.85 |
| Telangana | 25 790 | 1.00 |
| Mizoram | 23 870 | 0.92 |
| Punjab | 12 020 | 0.46 |
| Andhra Pradesh | 8 930 | 0.35 |
| Madhya Pradesh | 2 000 | 0.08 |
| Nagaland | 1 140 | 0.04 |
| Jammu and Kashmir | 740 | 0.03 |
| Haryana | 550 | 0.02 |
| Himachal Pradesh | 130 | 0.01 |
| Rajasthan | 70 | 0.00 |
| Total | 2 585 350 | - |

3.7.2 Export statistics

During the past five years, India exported between 64 000 and 148 000 tonnes of table grapes per year to more than 25 countries (International Trade Centre 2015). Key export markets include the Netherlands, Bangladesh, the United Arab Emirates, Russia and the United Kingdom (International Trade Centre 2015). Table 3.3 shows volumes of grapes exported from India to the top five countries from 2010 to 2014 (International Trade Centre 2015).

Table 3.3 Export volumes of fresh grapes from India to the top five export markets from 2010 to 2014 (International Trade Centre 2015)

| Destination | Volume (metric tonnes) | | | | |
|--------------------------------|------------------------|--------|---------|---------|---------|
| | 2010 | 2011 | 2012 | 2013 | 2014 |
| Netherlands | 25 050 | 12 988 | 27 187 | 39 327 | 41 674 |
| Russia | 1 790 | 1 937 | 7 429 | 20 573 | 21 093 |
| Bangladesh | 6 951 | 34 538 | 31 058 | 29 181 | 20 141 |
| UK | 11 438 | 3 913 | 10 632 | 14 580 | 16 779 |
| UAE | 6 095 | 7 842 | 13 023 | 13 622 | 10 028 |
| Total for top 5 export markets | 51 324 | 61 218 | 89 329 | 117 283 | 109 715 |
| Total for all export markets | 64 334 | 75 387 | 114 306 | 148 521 | 136 740 |

3.7.3 Export season

The main export season for table grapes from India is from February to the end of April (DAFF 2010; DPP 2012; International Trade Centre 2015). However, small volumes of table grapes may be exported at other times of the year.

4 Pest risk assessments for quarantine pests

Quarantine pests associated with table grapes from India are identified in the pest categorisation process (Appendix A). This chapter assesses the likelihood of the entry (importation and distribution), establishment and spread of these pests and the economic, including environmental, consequences these pests may cause if they were to enter, establish and spread in Australia.

Pest categorisation identified 27 quarantine pests associated with table grapes from India. Of these, 19 pests are of national concern and eight are of regional concern. Tables 4.1 to 4.4 identify these quarantine pests, and full details of the pest categorisation are given in Appendix A.

Assessments of risks associated with these pests are presented in this chapter unless otherwise indicated.

A full pest risk assessment is undertaken for each of the pests that have not been assessed previously by the department (Table 4.1).

Some of the pests considered here have been assessed previously by the department. For these pests, the outcomes of previous assessments will be adopted as appropriate, unless new information is available that suggests otherwise. Further explanation about the adoption of the outcomes of previous assessments is outlined below.

The likelihood of establishment and of spread of a pest in the PRA area will be comparable regardless of the fresh fruit commodity/country pathway in which the pest is imported into Australia, as these likelihoods relate specifically to events that occur in the PRA area and are independent of the importation pathway. The consequences of a pest are also independent of the importation pathway. For pests that have been assessed previously, the department reviewed the latest literature. If no new information is available that would significantly change the likelihood ratings for establishment and for spread, and the consequences the pests may cause, the ratings given in the previous assessments for these components will be adopted.

The need to reassess the likelihood of distribution for pests that have been assessed previously is considered on a case-by-case basis by comparing factors relevant to the distribution of table grapes from India with those assessed previously. These factors include the commodity type, time of year at which import takes place and availability and susceptibility of hosts during the time of import. After comparing these factors and reviewing the latest literature, the ratings of likelihood of distribution from the previous assessments will be adopted if the department considers that the likelihood of distribution for table grapes from India would be similar to, or at least not higher than, that given in the previous assessments. For some pests, the likelihood of distribution was reassessed and the reason for reassessing is provided in the introduction to the relevant pest risk assessment.

The need to reassess the likelihood of importation for pests that have been assessed previously is also considered on a case-by-case basis by comparing factors relevant to the importation of table grapes from India with those assessed previously. These factors include the commodity type, prevalence of the pest and commercial production practices. After comparing these factors

and reviewing the latest literature, the department considers it appropriate not to reassess the likelihood of importation where a) there is no information to suggest that the likelihood of importation for table grapes from India is not comparable to those assessed previously; or b) changes in the likelihood rating for importation will not change the overall outcome, that is the unrestricted risk estimate (URE) of achieving or exceeding Australia's ALOP.

The URE of achieving or exceeding Australia's ALOP will be adopted for pests for which the reassessment of both the likelihood of importation and the likelihood of distribution is considered not necessary because the URE outcome would not change from the previous assessment (Table 4.2).

Table 4.3 lists the pests for which the department considered that the differences in the information relating to the likelihood of importation and/or the likelihood of distribution are sufficient to warrant the reassessment of one or both of these components. For *Tobacco necrosis viruses*, the department also considered that new information warrants the reassessment of consequences of these pests.

In addition, the quarantine risks posed by *Drosophila suzukii* from all countries and for all commodities, including table grapes, were previously assessed in the final pest risk analysis report for *Drosophila suzukii* (Department of Agriculture 2013b). Therefore, there is no need to reassess this pest here (Table 4.4). A summary of pest information and the likelihood estimates from the final PRA report for *D. suzukii* is presented in this chapter for convenience.

Some pests identified in this assessment have been recorded in some regions of Australia, and due to interstate quarantine regulations are considered pests of regional concern. The acronym for the state for which the regional pest status is considered, such as 'WA' (Western Australia), is used to identify these organisms.

The department is aware of the recent changes in fungal nomenclature which ended the separate naming of different states of fungi with a pleiomorphic life cycle. However, as the nomenclature for these fungi is in a phase of transition and many priorities of names are still to be resolved, this report still uses dual names for most fungi. As official lists of accepted and rejected fungal names become available, these accepted names will be adopted.

Table grapes harvested, packed, stored and transported for export to Australia may need to travel variable distances to ports. Depending on the port of departure and arrival it could take up to four weeks for general sea freight from India to Australia. Table grapes could also potentially be air-freighted from India to Australia. While the assessments of the unrestricted risk undertaken in this risk analysis do not impose any mandatory measures during storage and transport, common commercial practices may impact on the survival of some pests. If these conditions are applied to all consignments for a minimum period of time, then those conditions can be considered as part of the assessment of the unrestricted risk.

Table 4.1 Quarantine pests for table grapes from India for which a full pest risk assessment is conducted

| Pest | Common name |
|---|------------------------------------|
| Bacteria | |
| <i>Xanthomonas campestris</i> pv. <i>viticola</i> | Grapevine bacterial canker disease |
| Viruses | |
| <i>Tomato black ring virus</i> | - |

Table 4.2 Quarantine pests for table grapes from India for which the URE outcome is adopted from previous assessments

| Pest | Common name |
|--|------------------------------|
| Phylloxera (Hemiptera: Phylloxeridae) | |
| <i>Daktulosphaira vitifoliae</i> (EP) | Grapevine phylloxera |
| Soft scales (Hemiptera: Coccidae) | |
| <i>Parthenolecanium corni</i> (EP, WA) | European fruit lecanium |
| Mealybugs (Hemiptera: Pseudococcidae) | |
| <i>Planococcus ficus</i> (EP) | Grapevine mealybug |
| <i>Planococcus lilacinus</i> (EP) | Coffee mealybug |
| <i>Planococcus minor</i> (EP, WA) | Pacific mealybug |
| <i>Rastrococcus iceryoides</i> (EP) | Downey snowline mealybug |
| Plume moth (Lepidoptera: Pterophoridae) | |
| <i>Platyptilia ignifera</i> (EP) | Large grape plume moth |
| Thrips (Thysanoptera: Thripidae) | |
| <i>Retithrips syriacus</i> (EP) | Black vine thrips |
| <i>Rhipiphorothrips cruentatus</i> (EP) | Grapevine thrips |
| Spider mite (Trombidiformes: Tetranychidae) | |
| <i>Tetranychus kanzawai</i> (EP, WA) | Kanzawa spider mite |
| Fungi | |
| <i>Guignardia bidwellii</i> (EP) | Black rot |
| <i>Monilinia fructigena</i> (EP) | Brown rot |
| <i>Phakopsora euvitis</i> (EP) | Grapevine leaf rust |
| <i>Phomopsis viticola</i> (EP, WA) | Phomopsis cane and leaf spot |

EP: Species has been assessed previously and import policy already exists.

WA: Regional pest for the state of Western Australia.

Table 4.3 Quarantine pests for table grapes from India for which some of the likelihood and/or consequence ratings are adopted from previous assessments

| Pest | Common name |
|--|--------------------|
| Fruit fly (Diptera: Tephritidae) | |
| <i>Bactrocera correcta</i> (EP) | Guava fruit fly |
| <i>Bactrocera dorsalis</i> (EP) | Oriental fruit fly |
| Tortricid moth (Lepidoptera: Tortricidae) | |
| <i>Archips machlopiis</i> | Leaf rolling moth |
| Pyralid moth (Lepidoptera: Pyralidae) | |
| <i>Cryptoblabes gnidiella</i> (EP) | Citrus pyralid |
| Fungi | |
| <i>Greeneria uvicola</i> (EP, WA) | Bitter rot |
| <i>Pestalotiopsis menezesiana</i> (EP, WA) | Fruit rot |
| <i>Pestalotiopsis uvicola</i> (EP, WA) | Fruit rot |
| <i>Pilidiella castaneicola</i> (EP, WA) | White rot |
| <i>Pilidiella diplodiella</i> (EP, WA) | White rot |
| Viruses | |
| <i>Tobacco necrosis viruses</i> (EP) | – |

EP: Species has been assessed previously and import policy already exists.

WA: Regional pest for the state of Western Australia.

Table 4.4 Quarantine pests for table grapes from India for which the likelihood and consequence ratings have been determined in a previous assessment

| Pest | Common name |
|--|-------------------------|
| Drosophila (Diptera: Drosophilidae) | |
| <i>Drosophila suzukii</i> (EP) | Spotted wing drosophila |

EP: Species has been assessed previously and import policy already exists.

Assessments for quarantine pests for which a full pest risk assessment is conducted

4.1 Grapevine bacterial canker disease

Xanthomonas campestris pv. *viticola*

Grapevine bacterial canker disease (GVBCD) was first detected in India in 1969 on 'Anab-e-Shahi' grapevines in Andhra Pradesh (Nayudu 1972). The causal organism was named as *Pseudomonas viticola* (Nayudu 1972), but was later changed to *Xanthomonas campestris* pv. *viticola* (Dye 1978). A large number of pathovars of *X. campestris* which were isolated in the 1950s and 60s, including *viticola*, have not been fully characterised and they are placed under *X. campestris* only provisionally (Parkinson et al. 2009).

Xanthomonas campestris pv. *viticola* is a gram negative, non-pigmented, plant pathogenic bacterium (Trindade et al. 2007). Natural hosts of this bacterium appear to be limited to grapevine. Susceptibility to GVBCD varies among different *Vitis* species (Chand et al. 1999). Chand (1999) reported that cultivars of *V. vinifera* are susceptible or highly susceptible to GVBCD whereas those of *V. labrusca* are resistant or moderately resistant to GVBCD. Many other *Vitis* species such as *V. rotundifolia* and *V. rupestris* are resistant and some *Vitis* species such as *V. riparia* and *V. parviflora* are highly resistant to GVBCD (Chand et al. 1999).

Several parts of grapevines including canes, leaves and grape bunches can be affected by GVBCD. Evidence of a systemic mechanism of spread of the bacterium in the conductive elements of the plant was recently reported (Tostes et al. 2014). The bacterium was detected in symptomatic and asymptomatic seeds of the grape cultivar Red Globe, both on the surface and internal tissue (Tostes et al. 2014).

The ultimate symptoms of GVBCD include stunting, cracking, irregular growth, a reduction in the health and vigour of the infected vines and considerable loss in yield and quality (Chand et al. 1999; Jambenal 2008). On grape bunches, symptoms develop on pedicels, rachises and berries as dark coloured lesions, cankers and vascular discoloration (Chand et al. 1999; Lima et al. 1999; Nascimento & Mariano 2004; Trindade et al. 2007). Infected berries are irregular in size and colour and severely infected berries are small, shrivelled, wilted and dried (Chand et al. 1999; Chand & Kishun 1990; Lima et al. 1999; Nascimento & Mariano 2004; Nascimento et al. 2006; Trindade et al. 2007).

In addition to host specificity, many *Xanthomonas* species and pathovars show tissue specificity as well, invading either intercellular spaces of mesophyll tissue (mesophyllic pathogens) and/or xylem elements of vascular tissue (vascular pathogens) (Ryan et al. 2011). Tostes et al. (2014) reported that *X. campestris* pv. *viticola* is both a mesophyllic and vascular pathogen.

Xanthomonas campestris pv. *viticola* survives mainly in the buds and canker lesions (Chand et al. 1999). The bacterium can survive in fallen leaves for about 45 days, and less under moist soil conditions (Chand et al. 1999).

The spread of *X. campestris* pv. *viticola* occurs through infected propagative material, agricultural equipment such as pruning and harvesting equipment, through dew, irrigation, rain

splash and wind-blown droplets (Chand et al. 1999). Although *X. campestris* pv. *viticola* has been detected in grapevine seeds (Tostes et al. 2014), seed transmission has not been demonstrated.

Epidemics of this disease in India are associated with rainfall in combination with wind, temperatures of between 20 and 30 degrees Celsius and humidity levels of around 80 per cent (Jambenal 2008). When bud burst coincides with frequent rains during early winter pruning, there is often a severe outbreak of the disease (Chand et al. 1999).

Grapevine bacterial canker disease has also been reported in Brazil (Halfeld-Vieira & De Lima Nechet 2006; Lima et al. 1999; Malavolta et al. 1999; Nascimento & Mariano 2004; Rodrigues Neto et al. 2011). The strains of the bacteria isolated from vineyards in Brazil were almost identical to the Indian strain of *X. campestris* pv. *viticola* (Trindade, Lima & Ferreira 2005).

The risk scenario of concern for *X. campestris* pv. *viticola* is that infected grape bunches with mild or no symptoms may escape detection during harvesting and packing procedures and hence may be exported to Australia.

4.1.1 Likelihood of entry

The likelihood of entry is considered in two parts, the likelihood of importation and the likelihood of distribution, which consider pre-border and post-border issues, respectively.

Likelihood of importation

The likelihood that *X. campestris* pv. *viticola* will arrive in Australia with the importation of table grapes from India is: **High**.

The following information provides supporting evidence for this assessment.

- Cultivars of *Vitis vinifera* are susceptible or highly susceptible to grapevine bacterial canker disease caused by *X. campestris* pv. *viticola* (Chand et al. 1999).
- In India, the disease is wide spread in the major grape growing regions of Maharashtra, Karnataka, Andhra Pradesh and Tamil Nadu (Chand et al. 1999).
- On grape bunches, the pathogen causes lesions and cankers on the pedicels and rachises (Chand et al. 1999; Trindade et al. 2007). Necrosis on pedicels and rachises occurring at the beginning of fruit set was followed by wilting and drying of berries (Lima et al. 1999).
- Infected berries are irregular in size and colour with brown to black necrotic lesions, and are small and shrivelled (Chand et al. 1999; Nascimento & Mariano 2004). These symptoms are observed at the beginning of fruit development at the pea-sized stage (Lima et al. 1999).
- The bacterium has also been detected on and in seeds of asymptomatic 'Red Globe' berries collected from vineyards effected with GVBCD (Tostes et al. 2014).
- Grape bunches showing obvious symptoms are likely to be removed during harvesting, grading and packing processes and would not be packed for export. However, grape bunches with no or mild symptoms could still be packed for export.
- The bacterium has been detected on grapevine leaf samples in Punjab (Chand et al. 1999) where the winter temperatures can go down to 0 degrees Celsius, therefore it is likely that the bacterium will survive low temperatures during transport and storage.

The prevalence of *X. campestris* pv. *viticola* in the major grape growing regions of India, the high susceptibility of cultivars of *V. vinifera* to GVBCD and the potential presence of the bacterium on and in grape seeds of asymptomatic berries support a likelihood estimate for importation of 'high'.

Likelihood of distribution

The likelihood that *X. campestris* pv. *viticola* will be distributed within Australia in a viable state as a result of the processing, sale or disposal of table grapes from India and subsequently transfer to a susceptible part of a host is: **Very low**.

The following information provides supporting evidence for this assessment.

- Imported grapes are intended for human consumption. It is expected that grape bunches will be distributed to many localities within all states and territories by wholesale and retail trade and by individual consumers.
- As grapes are easily damaged during handling (Mencarelli, Bellincontro & DiRenzo 2005), packed grapes may not be processed or handled again until they arrive at the retailers. Therefore, the bacterium, if present in packed grapes, is unlikely to be detected during transportation and distribution to retailers.
- Grape bunches with obvious symptoms of infection would not be marketable and would not be sold. Grape bunches without symptoms, or with only minor symptoms could be marketable and sold.
- Most fruit waste will be discarded into managed waste systems and will be disposed of in municipal tips and would therefore pose little risk of exposure to a suitable host.
- Consumers will discard small quantities of fruit waste in urban, rural and natural localities. Small amounts of fruit waste will be discarded in domestic compost. There is some potential for consumer waste being discarded near host plants, including commercially grown, household or wild host plants.
- In a study in Brazil the bacterium survived in infected grapevine tissues (fragmented shoots and leaves) on soil for at least 80 days but only 10 days when composted (Silva et al. 2012). In India the bacterium survived on fallen leaves normally for 45 days and up to 25 days under moist soil conditions (Chand et al. 1999).
- Generally survival of a pathogen in fruit waste is expected to be short due to dehydration and competition with other organisms. Based on the above studies in Brazil and India regarding survival of this bacterium on soil and fallen leaves, it is estimated that the bacterium could survive in fruit waste for some days.
- If present and still viable on fruit waste, the bacterium would then need to be transferred to a susceptible part of a host.
- To date, grapevine is the only confirmed natural host of *X. campestris* pv. *viticola*. In Australia, grapevines are grown in all states and territories, both commercially (ABS 2012a) and in household gardens.
- Peixoto (2007) reported that bacteria similar to *X. campestris* pv. *viticola* were isolated from the weeds *Alternanthera tenella*, *Amaranthus* sp., *Glycine* sp., and *Senna obtusifolia*. Artificial

inoculation studies suggested a number of other plant species as possible alternative hosts for *X. campestris* pv. *viticola*, namely, *Azadirachta indica*, *Phyllanthus maderaspatensis* (Nayudu 1972), *Mangifera indica* (Chand et al. 1999; Chand & Kishun 1990) and weed species *Chamaesyce hirta*, *Dactyloctenium aegyptium*, *Eragrostis pilosa* and *Pileas* sp. (Peixoto et al. 2007). Some of these plants are distributed throughout Australia. However, there have been no reports of infection of these hosts by *X. campestris* pv. *viticola* under natural conditions in the field.

- The bacterium can be transmitted by rain splash or wind driven rain. Outbreaks of the disease are correlated with frequent rains and cyclonic rains and hail storms (Chand et al. 1999). While rains with strong wind sometimes occur in parts of Australia, the transmission by rain splash and wind-blown droplets is still limited to a short distance for fruit waste on the ground.
- *Xanthomonas campestris* pv. *viticola* can be present in seeds of grape bunches from infected vines (Tostes et al. 2014). It is estimated that the majority of table grapes exported from India would be seedless, but some would be seeded.
- The proportion of grapevine seed that germinates depends on the cultivar, seed maturity, storage, stratification and planting conditions (Doijode 2001). Most grapevine seed is dormant and will not germinate unless it has been stratified. Night-time temperatures below 6 degrees Celsius during winter may be sufficient for stratification (Doijode 2001; Ellis, Hong & Roberts 1985). Seed of some cultivars will not germinate without stratification, other cultivars have very low germination rates when not stratified, but germination rates of up to 33 per cent from seed from fresh untreated berries of some cultivars has been reported (Forlani & Coppola 1977; Scott & Ink 1950; Singh 1961).
- Cold storage of imported table grapes during transport may stratify the seed and improve germination rates. Night-time temperatures in most temperate regions of Australia (Bureau of Meteorology 2010) may be low enough for stratification of grape seeds to occur naturally.
- A small proportion of grapevine seed from fruit waste may germinate. Successful germination will depend on local conditions. Many localities will not be suitable for grape seed germination.
- Grapevines are normally cultivated vegetatively, being propagated from cuttings by grafting onto rootstock or, less commonly, on their own roots (Zohary 1996). Seed is not used to establish vineyards because vines propagated from seed are likely to produce inferior berries; they are unlikely to be true to type after genetic segregation (Zohary 1996). This aspect of grapevine propagation is likely to deter members of the public from growing grapevines from seed from imported fruit, as will the relatively long time taken to grow a productive vine from seed (Olmo 1976) and the ready availability of grafted vines.
- Although *X. campestris* pv. *viticola* can be present in grapevine seeds (Tostes et al. 2014), seed to seedling transmission has not yet been demonstrated. A probability of seed transmission of 0.014 was reported for a related bacterium, *X. campestris* pv. *campestris* (Roberts et al. 1999).
- To date, there have been no vectors identified for this bacterium.

The potential presence of the bacterium in seeds, moderated by the limited potential for dispersal of the bacterium via rain splash or wind-blown droplets from infected fruit waste on

the ground to a susceptible part of a host which appears to be limited to grapevines, the small chance that grapevine seed will germinate and the lack of confirmed seed to seedling transmission of this bacterium support a likelihood estimate for distribution of 'very low'.

Overall likelihood of entry

The overall likelihood of entry is determined by combining the likelihood of importation with the likelihood of distribution using the matrix of rules shown in Table 2.2.

The likelihood that *X. campestris* pv. *viticola* will enter Australia as a result of trade in table grapes from India and be distributed in a viable state to a susceptible host is: **Very low**.

4.1.2 Likelihood of establishment

The likelihood that *X. campestris* pv. *viticola* will establish within Australia based on a comparison of factors in the source and destination areas that affect pest survival and reproduction, is: **Moderate**.

The following information provides supporting evidence for this assessment.

- Grapevines (*Vitis* spp.), the known host of *X. campestris* pv. *viticola*, are widely grown commercially and domestically across all states and territories of Australia.
- The pathogen can infect various tissues of grapevine, including leaves, grape bunches (rachises, pedicels, berries), canes (Chand et al. 1999; Nascimento & Mariano 2004; Trindade et al. 2007) and seeds (Tostes et al. 2014).
- The bacterium has also been detected on and in seeds, with and without symptoms, of asymptomatic berries collected from vineyards effected with GVBCD (Tostes et al. 2014). It is unlikely that asymptomatic infected hosts will be detected and destroyed.
- Peixoto (2007) reported that bacteria similar to *X. campestris* pv. *viticola* were isolated from the weeds *Alternanthera tenella*, *Amaranthus* sp., *Glycine* sp., and *Senna obtusifolia*. Artificial inoculation studies suggested that other plant species have potential to be alternative hosts of *X. campestris* pv. *viticola*, including *Phyllanthus maderaspatensis*, *Azadirachta indica*, *Chamaesyce hirta*, *Dactyloctenium aegyptium*, *Eragrostis pilosa*, *Mangifera indica* and *Phyllanthus maderaspatensis* (CABI 2012; Chand et al. 1999; Nayudu 1972; Peixoto et al. 2007). Some of these plants are distributed throughout Australia. However, there have been no reports of infection of these hosts by *X. campestris* pv. *viticola* under natural conditions.
- Cultivars of *V. vinifera* are susceptible or highly susceptible to *X. campestris* pv. *viticola* (Chand et al. 1999). The spread of *X. campestris* pv. *viticola* occurs through infected propagative material, agricultural equipment such as pruning and harvesting equipment, through dew, irrigation, rain splash and wind-blown droplets (Chand et al. 1999).
- Minimum inoculum dose required for successful infection in a field situation has not been determined for this pathogen. In artificial inoculations an inoculum concentration of 10¹² CFU/ml was most effective to induce the disease with minimum incubation period of 15 days (Chand et al. 1999).
- Optimum conditions for disease development are temperatures between 20 and 30 degrees Celsius, humidity levels of around 80 per cent, rain and wind associated with rain (Jambenal 2008). These conditions are not common in the grape production regions of Australia.

- Currently *X. campestris* pv. *viticola* is known to be well established only in India and Brazil. Both India and Brazil have been exporting table grapes to a number of countries. There is no information found with regards to specific conditions those importing countries require for *X. campestris* pv. *viticola* for table grapes from India or Brazil.
- The use of chemicals, including copper and antibiotics, was found not to be effective against *X. campestris* pv. *viticola*, especially in rainy weather (Chand, Patil & Kishun 1992; Jambenal 2008). The use of antibiotics to control plant diseases is currently not permitted in Australia. Copper-based chemicals are used in Australia but are unlikely to prevent the establishment of the disease.
- Presence of antagonistic organisms such as *Pseudomonas fluorescens* and *Bacillus subtilis* also give poor control on bunch infection in vivo (Jambenal 2008).
- GVBCD might establish in Australia from infected imported fruit if the infected seedlings survive.
- Few, if any, grapevine seedlings are likely to survive on agricultural land and in unmanaged localities in Australia. Seedling survival will depend on local conditions including rainfall.
- To date, there have been no vectors identified for this bacterium.

The ability of *X. campestris* pv. *viticola* to be transmitted from an infected volunteer grapevine by mechanical transmission, the high susceptibility of *Vitis vinifera* to this bacterium, asymptomatic plants are unlikely to be detected and destroyed, and the limited control measures available, moderated by the information that host of this bacterium is limited to grapevine, the likely limited climate conditions suitable for this bacterium in the grape production areas of Australia, the small chance that a volunteer grapevine seedling will survive and the lack of identified vector for this bacterium, support a likelihood estimate for establishment of 'moderate'.

4.1.3 Likelihood of spread

The likelihood that *X. campestris* pv. *viticola* will spread within Australia based on a comparison of factors in the source and destination areas that affect the expansion of the geographic distribution of the pest is: **Moderate**.

The following information provides supporting evidence for this assessment.

- Grapevines are grown commercially and domestically in all states and territories of Australia. Peixoto (2007) reported that bacteria similar to *X. campestris* pv. *viticola* were isolated from the weeds *Alternanthera tenella*, *Amaranthus* sp., *Glycine* sp., and *Senna obtusifolia*. Artificial inoculation studies suggested that other plant species have potential to be alternative hosts of *X. campestris* pv. *viticola*, including *Azadirachta indica*, *Chamaesyce hirta*, *Dactyloctenium aegyptium*, *Eragrostis pilosa*, *Mangifera indica* and *Phyllanthus maderaspatensis* (CABI 2012; Chand et al. 1999; Nayudu 1972; Peixoto et al. 2007). Some of these plants are distributed throughout Australia. However, there have been no reports of infection of these hosts under natural conditions in the field.
- Epidemics of GVBCD in India are associated with rainfall coupled with wind, temperatures of between 20 and 30 degrees Celsius and humidity levels of around 80 per cent (Jambenal 2008). These conditions are not common in the grape production regions of Australia.

- Currently *X. campestris* pv. *viticola* is known to be well established only in India and Brazil. Both India and Brazil have been exporting table grapes to a number of countries. There is no information found with regards to specific conditions those importing countries require for *X. campestris* pv. *viticola* for table grapes from India or Brazil.
- *Xanthomonas campestris* pv. *viticola* can be dispersed through infected propagative material, contaminated agricultural equipment such as containers, pruning shears and gloves, as well as rain splash and wind-blown droplets (Chand et al. 1999; Nascimento & Mariano 2004; Tostes et al. 2014).
- The long distance dispersal of *X. campestris* pv. *viticola* is more likely to be through the movement of infected grapevine planting material. The interstate movement of grapevine planting material is regulated in Australia (Plant Health Australia 2009b). Grapevine planting material certified as being free of pests and pathogens is available from accredited nurseries in Australia, as per the Vine Industry Nursery Accreditation Scheme (VINA 2008).
- It is possible that grapevine plants can be contaminated with *X. campestris* pv. *viticola* without showing symptoms (Tostes et al. 2014), which increases the potential for unintended spread of the bacterium.
- To date, there have been no vectors identified for this bacterium.

The ability of the bacterium to be dispersed through grapevine propagative materials and agricultural equipment and natural means, moderated by the limited natural dispersal of the bacterium, the information that host of this bacterium is limited to grapevine, the systems in place for the movement and certification of grapevine planting material in Australia, the likely limited climate conditions suitable for the development of GVBCD in the grape production areas of Australia, the lack of identified vector for this bacterium and the fact that the spread of the bacterium has so far been limited to India and Brazil support a likelihood estimate for spread of 'moderate'.

4.1.4 Overall likelihood of entry, establishment and spread

The overall likelihood of entry, establishment and spread is determined by combining the likelihoods of entry, of establishment and of spread using the matrix of rules shown in Table 2.2.

The overall likelihood that *X. campestris* pv. *viticola* will enter Australia as a result of trade in table grapes from India, be distributed in a viable state to a susceptible host, establish in Australia and subsequently spread within Australia is: **Very low**.

4.1.5 Consequences

The potential consequences of the establishment of *X. campestris* pv. *viticola* in Australia have been estimated according to the methods described in Table 2.3.

Based on the decision rules described in Table 2.4, that is, where the potential consequences of a pest with respect to one or more criteria is 'D', the overall consequences are estimated to be **Low**.

| Criterion | Estimate and rationale |
|----------------------------------|--|
| Direct | |
| Plant life or health | <p>D—Significant at the district level.</p> <p>In India, <i>X. campestris</i> pv. <i>viticola</i> is a pest of major economic significance throughout the major grape growing provinces of Maharashtra, Karnataka, Andhra Pradesh, and Tamil Nadu (Chand et al. 1999; Jambenal 2008). There are reports of this disease causing yield losses of between 60 and 80 per cent in these regions in September pruned vineyards (Chand et al. 1999; Chand, Patil & Kishun 1991; Jambenal 2008).</p> <p>In Brazil, the disease is considered the most important bacterial disease in the major grape growing region of São Francisco, causing yield losses to eight varieties of grapes that are commonly grown (Nascimento & Mariano 2004; Nascimento et al. 2006; Trindade et al. 2007). The incidence of 100 per cent disease symptoms and nearly total yield loss in some cases was reported (Lima et al. 1999).</p> <p>The natural host range of <i>X. campestris</i> pv. <i>viticola</i> appears to be limited to grapevine. Grapevine are grown commercially in all Australian states and territories for wine industry, table grapes, and dried grapes (ABS 2012a). In 2010–11, 1 597 669 tonnes of grapes produced in Australia were used for wine making (ABS 2012a). In 2007–8 the value of the Australian wine produced was \$4.77 billion (ABS 2009b). In 2013, the annual production of table grapes in Australia was about 120 000 tonnes with a farm gate value of \$330 million and Australia exported more than 70 000 tonnes to 52 countries earning about \$200 million (Australian Table Grape Association 2013). In 2010–11, approximately 11831 tonnes of grapes were used for drying (ABS 2012a).</p> <p>The bacterium is currently recorded in only two countries, India and Brazil. In India, even though the bacterium was found on grape leaves in Punjab, no economic damage was reported in this state.</p> <p>The extent of damage this bacterium may cause, if established, in Australia is uncertain. Optimum conditions for disease development are temperatures between 20 and 30 °C, humidity levels of around 80 per cent, rain and wind associated with rain (Jambenal 2008). These conditions are not common in the grape production regions of Australia. However, it is expected that the bacterium, if established, may cause significant damage to grapevine grown in some localised areas which have climate conditions suitable for GVBCD development.</p> |
| Other aspects of the environment | <p>A—Indiscernible at the local level.</p> <p>There are currently no known direct consequences of this bacterium on other aspects of the natural environment.</p> |
| Indirect | |
| Eradication, control | <p>D—Significant at the district level.</p> <p>The use of bactericides alone are not an effective control option and widespread control is mainly achieved through an integrated management system (Chand, Patil & Kishun 1992; Nascimento et al. 2006). The integrated management system involves using healthy propagative material, field inspection, chemical sprays, drastic pruning on infected plants, management of the time of production pruning, disinfection of agricultural equipment and vehicles, windbreaks to reduce pathogen dissemination and curbing the excess use of water (Jambenal 2008; Nascimento & Mariano 2004; Nascimento et al. 2006; Trindade et al. 2007). The integrated management system would incur significant management costs to the grape industry in areas which are suitable for disease development.</p> <p>The recent finding that the pathogen spreads systemically within the plants (Tostes et al. 2014) will require research investment on systemic control of the disease.</p> <p>While antibiotics are used in India to control GVBCD, antibiotics are currently not registered for use to control plant diseases in Australia.</p> <p>Eradication attempt on an outbreak in one property in the State of São Paulo in Brazil in 2009 resulted in the destruction of 4700 plants (Rodrigues Neto et al. 2011).</p> |
| Domestic trade | <p>D—Significant at the district level.</p> <p>The presence of <i>X. campestris</i> pv. <i>viticola</i> in commercial production areas is likely to result in interstate trade restrictions on table grapes, potential loss of markets and significant industry adjustment at the district level.</p> |

| Criterion | Estimate and rationale |
|----------------------------------|--|
| International trade | <p>D—Significant at the district level.</p> <p>At present, <i>X. campestris</i> pv. <i>viticola</i> is only recorded in India and Brazil (CABI 2012; Trindade et al. 2007).</p> <p>The European Union and the United States do not require measures specific to this bacterium for table grapes imported from India or Brazil.</p> <p>The presence of this pathogen in commercial production areas of table grapes in Australia could potentially limit access to some overseas markets that are free of this pathogen.</p> |
| Environmental and non-commercial | <p>B—Minor significance at the local level.</p> <p>Any additional usage of pesticide sprays may affect the environment.</p> <p>Streptomycin or any other antibiotic sprays are not currently registered for the control of plant pests in Australia. It is possible that the use of antibiotics could be permitted for emergency use under strict controls in an eradication programme. Registration for more widespread use of antibiotics to control plant pests would require the evaluation of the environmental impact. Significant issues that would need to be considered include the potential that resistance to antibiotics may develop and the potential for residues in other products such as honey. Streptomycin resistance of <i>X. campestris</i> pv. <i>viticola</i> has been reported (Chand et al. 1994; Reddy 2011).</p> <p>Copper sprays are already in use in Australia to control a range of plant pests.</p> |

4.1.6 Unrestricted risk estimate

Unrestricted risk is the result of combining the likelihoods of entry, establishment and spread with the outcome of overall consequences. Likelihoods and consequences are combined using the risk estimation matrix shown in Table 2.5.

Unrestricted risk estimate for *Xanthomonas campestris* pv. *viticola*

| | |
|---|------------|
| Overall likelihood of entry, establishment and spread | Very low |
| Consequences | Low |
| Unrestricted risk | Negligible |

As indicated, the unrestricted risk estimate for *Xanthomonas campestris* pv. *viticola* has been assessed as 'negligible' which achieves Australia's ALOP. Therefore, no specific risk management measures are required for this pest.

4.2 Tomato black ring virus

Tomato black ring virus

Tomato black ring virus (TBRV) is a member of the genus *Nepovirus*, Comoviridae family, and infects over 50 plant species, including grapevines and a wide range of economically important crop species (carrots, tomato, strawberry), cultivated garden plants (ash, daffodils) and weeds (chickweed, privet) (Harper et al. 2011; Harris, Gibbs & Gibbs 2002; Murrant 1983; Taylor & Brown 1997).

The virus has several strains including beet ringspot, celery yellow vein, lettuce ringspot, potato bouquet and potato pseudo-aucuba (CABI & EPPO 1997d; Murrant 1970). Antigenic variations between the strains separate the virus into two groups, the 'English' serotype contains the type, lettuce ringspot, celery yellow vein and potato bouquet isolates, whereas the 'Scottish' serotype contains the beet ringspot and potato pseudo-aucuba isolates (Harrison 1958; Murrant 1970).

The virus was first reported in India in 1966 and has since been confirmed to be present in Andhra Pradesh, Karnataka and Tamil Nadu, (CABI 2012; CABI & EPPO 1997d; Madhusudan & Govindu 1985). Grapes are commercially grown in these states (DPP 2007). It is unknown which virus strains are present in India.

Most naturally infected weed and crop plants may show few or no symptoms especially in the first year of infection (Murrant 1983), or when the infection occurs through the seed (CABI & EPPO 1997d). Despite this, infection with TBRV ultimately reduces plant growth and vigour (DPP 2012; Harper et al. 2011).

When symptoms are present on infected weed and crop plants, they may include systemic chlorotic ringspot, leaf mottle and deformation, black coalescent rings on the leaves, vein yellowing and stunting as well as flecking and reddish streaking on the petiole and stem on symptomatic host species (Brunt et al. 1996; Chowfla, Sharma & Thakur 1999). Where infection occurs through nematode transmission, TBRV infection often appears as patches of poor growth which slowly extend in size each year (CABI & EPPO 1997d).

On grapevine, it has been reported to cause yellow rings and blotches, malformed leaves, asymmetrical leaves, premature senescence and leaf fall (DPP 2012; Harris, Gibbs & Gibbs 2002; Martelli 1999). Fruit may be small and poorly set or malformed (DPP 2012; Stobbs & van Schagen 1984). *Tomato black ring virus* has also been detected in asymptomatic vines (Laveau et al. 2013).

The virus is transmitted and disseminated by several mechanisms. The virus is transmitted through the soil by nematode species. The English strain is efficiently transmitted by *Longidorus attenuatus* (Brown, Murrant & Trudgill 1989), whereas strains which are more related to the Scottish strain are more efficiently transmitted by *Longidorus elongatus* (Brown, Murrant & Trudgill 1989). Transmission of TBRV through grapevine seed to the emerging seedlings has not been studied. Martelli (1978) assumes that TBRV is seed-borne in grapevines. It is believed that nearly all of the nematode borne viruses, such as TBRV, are transmitted and distributed to some extent through the seed of their principal hosts (Murrant 1983).

Tomato black ring virus was shown to be transmitted through seed in 19 out of 28 species in a

study by Lister and Murrant (1967). The virus can also be transmitted by mechanical inoculation and grafting (Harrison 1996).

The virus is able to be transmitted horizontally by pollen and may infect the plant through the fertilised flower (Card, Pearson & Clover 2007). *Tomato black ring virus* is able to be transmitted via pollen in plants such as raspberry (*Rubus idaeus* L.) (Harris, Gibbs & Gibbs 2002; Harrison 1996; Lister & Murrant 1967; Murrant 1983).

The risk scenario of concern is the importation of fruit infected with TBRV, distribution of fruit waste, germination of some grape seeds from the waste, seed transmission and replication of the virus, survival of infected seedlings and the transmission of TBRV to other host plants in Australia.

Tomato black ring virus was included in the existing import policy for truss tomatoes from the Netherlands (Biosecurity Australia 2003). However, due to additional information and analysis, a full risk assessment is undertaken here for table grapes from India.

4.2.1 Likelihood of entry

The likelihood of entry is considered in two parts, the likelihood of importation and the likelihood of distribution, which consider pre-border and post-border issues, respectively.

Likelihood of importation

The likelihood that *Tomato black ring virus* will arrive in Australia with the importation of table grapes from India is: **Moderate**.

The following information provides supporting evidence for this assessment.

- *Tomato black ring virus* can be present in grapevine seed (Martelli 1978). Although there is some risk of fresh grapes with TBRV-infected seed being imported, it is estimated that the majority of table grapes exported from India would be seedless.
- *Tomato black ring virus* has been reported in grapevine in India (DPP 2007, 2012). The virus has been found in Andhra Pradesh, Karnataka and Tamil Nadu (CABI 2012; CABI & EPPO 1997d; Madhusudan & Govindu 1985). These states produced 1.27 per cent, 12.92 per cent and 1.75 per cent, respectively, of the total grape production in India in 2012-13 (APEDA 2015).
- *Tomato black ring virus* occurs systemically and most naturally infected weed and crop plants may show few or no symptoms especially in the year of infection, or when the infection occurs through the seed (CABI & EPPO 1997d; Murrant 1970). *Tomato black ring virus* has been detected in asymptomatic grapevines (Laveau et al. 2013; Stellmach 1970).
- Fruit from infected vines may be small and poorly set or malformed (DPP 2012; Stobbs & van Schagen 1984).
- Infected bunches showing symptoms are likely to be culled during harvesting, grading and packing.
- Healthy looking grape bunches carrying TBRV, and in some cases containing infected seeds, might be imported into Australia.

- Grapes are usually pre-cooled to less than 4 degrees Celsius after packing and cold stored at 0–2 degrees Celsius at 90–95 per cent humidity after palletizing until shipment (DPP 2009). Conditions are maintained at 0–2 degrees Celsius and 90–95 per cent humidity throughout transport (DAFF 2010; DPP 2009, 2012). TBRV is likely to survive conditions during transport and storage.

Systemic infection and asymptomatic infection of grape bunches in some cases, moderated by the fact that infected fruit which show symptoms are likely to be culled during harvesting and packing processes, support a likelihood estimate for importation of 'moderate'.

Likelihood of distribution

The likelihood that *Tomato black ring virus* will be distributed within Australia in a viable state as a result of the processing, sale or disposal of table grapes from India and subsequently transfer to a susceptible part of a host is: **Very low**.

The following information provides supporting evidence for this assessment.

- If table grapes are imported, they will be distributed through the domestic supply chain and sold to the public for consumption.
- Most fruit waste will be discarded into managed waste systems and will be disposed of in municipal tips. Consumers will discard small quantities of fruit waste in urban, rural and natural localities.
- Some table grape waste may go to household compost.
- For the virus to reach a susceptible part of a host from fruit waste, the only pathway considered feasible is transmission through the germination of infected grapevine seed.
- The proportion of grapevine seed that germinates depends on the cultivar, seed maturity, storage, stratification and planting conditions (Doijode 2001). Most grapevine seed is dormant and will not germinate unless it has been stratified. Night-time temperatures below 6 degrees Celsius during winter may be sufficient for stratification (Doijode 2001; Ellis, Hong & Roberts 1985). Seed of some cultivars will not germinate without stratification, other cultivars have very low germination rates when not stratified, but germination rates of up to 33 per cent from seed from fresh untreated berries of some cultivars has been reported (Forlani & Coppola 1977; Scott & Ink 1950; Singh 1961).
- Cold storage of imported table grapes during transport may stratify the seed and improve germination rates. Night time temperatures in most temperate regions of Australia (Bureau of Meteorology 2010) may be low enough for stratification of grape seeds to occur naturally.
- A small proportion of grapevine seed from fruit waste may germinate. Successful germination will depend on local conditions. Many localities will not be suitable for grape seed germination.
- Grapevines are normally cultivated vegetatively, being propagated from cuttings by grafting onto rootstock or, less commonly, on their own roots (Zohary 1996). Seed is not used to establish vineyards because vines propagated from seed are likely to produce inferior berries; they are unlikely to be true to type after genetic segregation (Zohary 1996). This aspect of grapevine propagation is likely to deter members of the public from growing

grapevines from seed from imported fruit, as will the relatively long time taken to grow a productive vine from seed (Olmo 1976) and the ready availability of grafted vines.

- Transmission of TBRV through grapevine seed to the emerging seedlings has not been studied. However, rates of TBRV transmission through seed have been documented in at least 24 other species in 13 botanical families, ranging from 3 to 100 per cent transmission effectiveness (CABI 2012; Murant 1983). The capacity to be seed transmitted is known to vary among strains of other virus species, and to vary between cultivars of the same plant species (Albrechtsen 2006); this may also be true of TBRV and *Vitis* species. Some strains of TBRV are probably seed transmitted in some grapevine cultivars.
- If grape seedlings grow from TBRV-infected seed, they may be infected with the virus.

The small chance that grapevine seed will germinate and the small chance that the virus will be transmitted from seed to seedling, supports a likelihood estimate for distribution of 'very low'.

Overall likelihood of entry

The overall likelihood of entry is determined by combining the likelihood of importation with the likelihood of distribution using the matrix of rules shown in Table 2.2.

The likelihood that *Tomato black ring virus* will enter Australia as a result of trade in table grapes from India and be distributed in a viable state to a susceptible host is: **Very low**.

4.2.2 Likelihood of establishment

The likelihood that *Tomato black ring virus* will establish within Australia, based on a comparison of factors in the source and destination areas that affect pest survival and reproduction, is: **Very low**.

The following information provides supporting evidence for this assessment.

- *Tomato black ring virus* has been reported in India, Turkey and the European Union (CABI & EPPO 1997d) and has demonstrated its ability to establish in a range of environments.
- As some strains of TBRV are probably seed transmitted in some grapevine cultivars, TBRV might establish in Australia from infected imported fruit if the infected seedlings survive.
- In Europe, volunteer grapevines grow as weeds in small numbers. Most of these weedy vines are probably rootstocks that have escaped cultivation and grown vegetatively, but some may have grown from seed (Arrigo & Arnold 2007; Ocete et al. 2008; Zohary 1996), suggesting seedlings sometimes survive in unmanaged environments. Small numbers of seedlings may survive in some regions of Europe because of the favourable climate and soils.
- *Vitis vinifera* is very infrequently encountered as a weed in Australia (Office of the Gene Technology Regulator 2003). There are reports of *V. vinifera* growing as a weed on roadsides and in disturbed areas in NSW, Vic. and WA (Richardson, Richardson & Shepherd 2006), but the number of plants is very small. Vines have been found near established vineyards and water-courses (Conn 2010). *Vitis vinifera* has been recorded as naturalised in WA and on the North Coast and North Western Slopes of NSW (Conn 2010). Reports indicating the origins of the naturalised plants were not found. It is likely that most or all of the plants found outside of vineyards have grown vegetatively from cultivated vines. Those found on

roadsides and in watercourses may have grown from plants taken from gardens or vineyards that have been discarded with other vegetation. Some weedy grapevines may be very old and the rate of successful invasion may be extremely small. If a plant grew from seed, it is likely the seed was from a rootstock, as rootstocks are more hardy.

- Few, if any, grapevine seedlings are likely to survive on agricultural land and in unmanaged localities in Australia. Seedling survival will depend on local conditions including rainfall.
- If an infected grapevine seedling survives, TBRV may be transmitted to other host plants through soil by species of the free living soil inhabiting nematodes *Longidorus elongatus* and *Longidorus attenuatus* (Chowfla, Sharma & Thakur 1999; Martelli 1999). *Longidorus elongatus* has been recorded in South Australia and Tasmania. Other unidentified specimens of *Longidorus* spp. have been recorded in NSW, Qld and Vic. (McLeod, Reay & Smyth 1994; Plant Health Australia 2001). Transmission efficacies of each of these nematode species varies from between 5 and 78 per cent depending on the serotype and the vector used (Brown, Murrant & Trudgill 1989; CABI 2012; Martelli 1978).
- Both larvae and adult nematodes transmit the virus but the virus does not multiply in the vector, and it is not retained after moulting, nor is it passed to nematode progeny (CABI 2012). Experiments have shown that *L. elongatus* retained infectivity only up to nine weeks when maintained in fallow soil (Lister & Murrant 1967).
- The vector potential of each of these nematodes is dependent somewhat on which virus serotype is present in India.
- Investigations have shown that nematode transmission of nepoviruses alone are not effective dispersal agents of the virus in terms of distance, and may only spread the disease between one and two metres per year, which is the case for similar viruses (Martelli 1978).
- *Tomato black ring virus* has been found naturally infecting over 30 species of commercial and cultivated crops and weeds including *Capsella bursa-pastoris* (shepherd's purse), *Cerastium vulgatum* (mouse-eared chickweed), *Chenopodium album* (fat hen), *Chenopodium quinoa* (quinoa), *Fragaria x ananassa* (strawberry), *Fraxinus* spp. (ash), *Gladiolus* spp., *Glycine max* (soybean), *Ligustrum vulgare* (privet), *Lycopersicon esculentum* (tomato), *Narcissus pseudonarcissus* (daffodil), *Petunia violacea* (petunia), *Poa annua* (winter grass), *Polygonum aviculare* (wireweed), *Polygonum convolvulus* (black bindweed), *Prunus dulcis* (almond), *Quercus robur* (English oak), *Ribes nigrum* (black currant), *Ribes rubrum* (red currant), *Ribes sanguineum* (flowering currant), *Rubus* (blackberry/raspberry), *Senecio vulgaris* (common groundsel), *Solanum melongena* (aubergine), *Solanum tuberosum* (potato), *Stellaria media* (chickweed) *Syringa vulgaris* (lilac), *Tulipa* (tulip) and *Vitis* (grapevine) (Harper et al. 2011; Harris, Gibbs & Gibbs 2002; Murrant 1983; Taylor & Brown 1997). Most of these host species grow in Australia and some are widely distributed.
- The natural infection of several plant species suggests that the vector nematodes are able to transmit TBRV between different plant species in the field.

The possibility of TBRV being transmitted through nematodes from an infected grapevine seedling to other host plants nearby, moderated by the small chance that a volunteer grapevine seedling will survive, the uncertainty about the presence of nematode vectors in parts of

Australia and the virus serotype present in India and the limited vector potential of the nematodes supports a likelihood estimate for establishment of 'very low'.

4.2.3 Likelihood of spread

The likelihood that *Tomato black ring virus* will spread within Australia, based on a comparison of factors in source and destination areas that affect the expansion of the geographic distribution of the pest, is: **Moderate**.

The following information provides supporting evidence for this assessment.

- Potential hosts of TBRV are widely available in Australia, both commercially and in home gardens.
- *Tomato black ring virus* is transmitted through the soil between host plants by species of the free living soil inhabiting nematodes, *Longidorus elongatus* and *Longidorus attenuatus* (Chowfla, Sharma & Thakur 1999; Martelli 1999). *Longidorus elongatus* has been recorded in South Australia and Tasmania. Other unidentified specimens of *Longidorus* spp. have been recorded in NSW, Qld and Vic. (McLeod, Reay & Smyth 1994; Plant Health Australia 2001). Transmission efficacies of each of these nematode species varies from between 5 and 78 per cent depending on the serotype and the vector used (Brown, Murant & Trudgill 1989; CABI 2012; Martelli 1978).
- Both larvae and adult nematodes transmit the virus but the virus does not multiply in the vector, and it is not retained after moulting, nor is it passed to nematode progeny (CABI 2012). Experiments have shown that *L. elongatus* retained infectivity only up to nine weeks when maintained in fallow soil (Lister & Murant 1967).
- The vector potential of each of these nematodes is dependent somewhat on which virus serotype is present in India.
- Investigations have shown that nematode transmission of nepoviruses alone are not effective dispersal agents of the virus in terms of distance, and may only spread the disease between one and two metres per year, which is the case for similar viruses (Martelli 1978).
- There is the potential that nematodes could be moved long distances through contaminated soil and machinery from farming practices (Watson 2004). Each could potentially contain nematodes which carry the virus which would facilitate the dispersal of the virus to new areas. This was believed to be the case in France, where nematode vectors (*Xiphinema index*) of Grapevine fanleaf virus were thought to be inadvertently transported short and long distances with both the movement of soil and farm machinery (Villate et al. 2008).
- *Tomato black ring virus* has the potential to transfer both horizontally and vertically (Card, Pearson & Clover 2007). This means that the virus is able to be transmitted horizontally by pollen and may infect the plant through the fertilised flower, and it may be transmitted vertically, in which case it may infect the seed and the seedling that will grow from that seed (Card, Pearson & Clover 2007).
- *Tomato black ring virus* is able to be transmitted via pollen in plants such as raspberry (*Rubus idaeus* L.) (Harris, Gibbs & Gibbs 2002; Harrison 1996; Lister & Murant 1967; Murant 1983). The capacity for TBRV to be pollen transmitted would likely vary between species

and cultivars, and it is possible that some grape cultivars will be effective pollen transmitters.

- The evidence of the role of nepovirus pollen transmission from virus contaminated pollen to the mother plant only through fertilization in the field is inconclusive (Mink 1993).
- *Vitis vinifera* can be wind pollinated, and studies have shown that pollen from some cultivars of *Vitis vinifera* has the potential to travel from between 500 metres and 3 kilometres throughout the landscape (Di Vecchi-Staraz et al. 2009).
- Transmission of TBRV through grapevine seed to the emerging seedlings has not been studied. However, rates of TBRV transmission through seed have been documented in at least 24 other species in 13 botanical families, ranging from 3 to 100 per cent transmission effectiveness (CABI 2012; Murrant 1983). The capacity for TBRV to be seed transmitted would likely vary between species and cultivars, and it is possible that some grape cultivars will be effective seed transmitters.
- Transmission of TBRV through seed of infected weeds could contribute to the dispersal of the virus over a wide area. Infected weed seeds could also provide a reservoir of the virus in the soil.
- Infection through the seed often leads to few or no visible symptoms and therefore, infected weeds or crops in commercial or ornamental plantings may not be detected (CABI & EPPO 1997d; Murrant 1983).
- *Tomato black ring virus* is transmitted by grafting and is disseminated with infected propagation material (Harrison 1996).
- In Germany, preliminary surveys for viruses affecting vineyards identified TBRV from only one region. However, nearly all vineyards in this region were infected and there was an active spread of the virus from vine to vine (Rüdel 1985).

The possibilities of transmission by two species of nematodes (*Longidorus elongatus* and *Longidorus attenuatus*) or spread in seed or pollen and the large host range, moderated by the uncertainty about the presence of nematode vectors in Australia and the limited vector potential of the nematodes, support a likelihood estimate for spread of 'moderate'.

4.2.4 Overall likelihood of entry, establishment and spread

The overall likelihood of entry, establishment and spread is determined by combining the likelihoods of entry, of establishment and of spread using the matrix of rules shown in Table 2.2.

The overall likelihood that *Tomato black ring virus* will enter Australia as a result of trade in table grapes from India, be distributed in a viable state to a susceptible host, establish in Australia and subsequently spread within Australia is: **Extremely low**.

4.2.5 Consequences

The potential consequences of the establishment of *Tomato black ring virus* in Australia have been estimated according to the methods described in Table 2.3.

Based on the decision rules described in Table 2.4, that is, where the potential consequences of a pest with respect to one or more criteria are 'E', the overall consequences are estimated to be **Moderate**.

| Criterion | Estimate and rationale |
|----------------------------------|--|
| Direct | |
| Plant life or health | <p>E—Significant at the regional level</p> <p><i>Tomato black ring virus</i> has a very wide host range with host species being grown in both commercial and ornamental situations (CABI 2012; Harris, Gibbs & Gibbs 2002). The main commercially grown crops which could be affected in Australia are asparagus, blackberry, currants, grapes, potatoes, raspberry, strawberry and tomato.</p> <p><i>Tomato black ring virus</i> causes a range of symptoms in grapevine which is similar to many of the nepoviruses (CABI 2012; Martelli 1978). These include distortion of the vine, malformation and chlorotic ringspots of the leaves, premature leaf fall and senescence, as well as the ultimate decline and death of the vine (Harris, Gibbs & Gibbs 2002; Martelli 1978, 1999). This virus also causes low quantity and quality of yield, reduced rooting ability, shortening of reproductive life and low graft take (Martelli 1978). In Germany, despite only being recorded in one region, observations indicated that this virus is capable of inducing very high crop losses in grape (Rüdel 1985).</p> <p><i>Tomato black ring virus</i> has been recorded to cause significant crop losses in strawberry crops, especially when present with other viral infections or adverse environmental conditions (Martin & Tzanetakis 2006). Symptoms in indicator plants vary from being asymptomatic to causing yellow blotching, ring spots, crinkling of the leaves, stunting and plant death (Martin & Tzanetakis 2006).</p> <p>In raspberry, TBRV causes ringspot on the leaves and may decrease the yield of 'tolerant' cultivars. In the cultivar, 'Seedling V', it causes many short, spindly and brittle young shoots with ill-defined chlorotic markings on the leaves. In 'Malling Exploit' leaves develop faint chlorotic mottling or ringspots initially and later the canes are stunted, yield is decreased and some drupelets are aborted. In 'Norfolk Giant' the leaves develop leaf-curl (Murant 1987). In red currant, TBRV was found in plants that showed pronounced yellow line-pattern symptoms (Jones & McGavin 1996).</p> <p>Yield losses due to TBRV have also been reported for various other crops, such as potatoes in Germany, a 20 per cent yield reduction of asparagus also infected with other viruses (Harris, Gibbs & Gibbs 2002) and the report that young tomato plants infected with the virus are frequently killed (Chowfla, Sharma & Thakur 1999). <i>Tomato black ring virus</i> also affects a range of ornamental plant species, which may affect a variety of environments such as street and city plantings, home gardens or the nursery sector.</p> <p>The value of total grape production for all uses (wine, dried and table) was \$1 040.6 million for the 2011–12 financial year (ABS 2012b).</p> <p>The production value in 2011–12 for the Australian strawberry industry was approximately \$200 million (Plant Health Australia 2014). This production occurs in Queensland, Victoria, South Australia, Western Australia, Tasmania and New South Wales (Plant Health Australia 2010). <i>Tomato black ring virus</i> is listed as a pathogen of quarantine concern to this industry (Plant Health Australia 2010).</p> <p>The annual Australian production in 2011–12 for <i>Rubus</i> spp. was over 1000 tonnes, worth approximately \$25 million. <i>Tomato black ring virus</i> is listed as a pathogen of quarantine concern to this industry (Plant Health Australia 2013b).</p> <p>Potato production in Australia in 2011–12 was 1 288 186 tonnes, which was worth \$625.6 million (Ausveg 2013a). <i>Tomato black ring virus</i> is listed as a pathogen of quarantine concern to this industry (Plant Health Australia 2013a).</p> <p>In 2010–11, Australia produced 10 276 tonnes of asparagus, which was worth \$68.7 million (Ausveg 2013b). The tomato production for 2011–12 was 371 514 tonnes and was worth \$351.8 million (Ausveg 2013a). <i>Tomato black ring virus</i> is listed as a pathogen of quarantine concern to the vegetable industry, although the overall risk is estimated as very low (Plant Health Australia 2011).</p> |
| Other aspects of the environment | <p>A—Indiscernible at local level</p> <p><i>Tomato black ring virus</i> naturally infects a range of common weeds including <i>Capsella bursa-pastoris</i> (shepherd's purse), <i>Ligustrum vulgare</i> (privet), <i>Senecio</i> (common groundsel), <i>Sonchus oleraceus</i> (sowthistle) and <i>Stellaria media</i> (chickweed) (CABI 2012). These weeds are distributed throughout Australia and infection may reduce the weed burden within some ecosystems.</p> |

| Criterion | Estimate and rationale |
|----------------------|---|
| Indirect | |
| Eradication, control | <p>E—Major significance at the district level</p> <p>In the absence of nematode vectors, eradication of TBRV could be achieved through removal and destruction of infected plants in combination with monitoring and weed control (Stobbs & van Schagen 1984).</p> <p>Eradication and control in the field can be difficult and may not be possible if weeds are infected and vector nematodes are present. It could also be difficult because the virus does not express clear symptoms in many of its possible hosts (Harper et al. 2011). The virus may be maintained in certain weeds and nematodes may spread it to new plantings.</p> <p>If a TBRV outbreak was detected in a vineyard, it is likely that local eradication would be undertaken. Detection in a vineyard may be delayed because symptoms take time to develop and are not diagnostically distinctive. Laboratory testing is required to confirm a diagnosis (Laveau et al. 2013). Vines infected with the virus would be destroyed. Properties in contact with the infected property would be traced and surveyed, and adjoining and nearby properties would be surveyed. Surveillance may continue for several seasons. Surveillance and testing is costly. Vines that are destroyed would probably be replaced with pathogen-free planting material. In Australia, disease tested grapevine planting material is available through the Vine Industry Nursery Accreditation Scheme (VINA 2008).</p> <p>If an outbreak was detected in a greenhouse or field tomato crop it would probably be eradicated through action coordinated at the state or national level. An outbreak of TBRV in tomato, potato or other crop hosts may not be detected until it has spread to several crops, properties and species. Quarantine would probably be enforced on infected properties. Laboratory testing would be required to confirm a diagnosis.</p> <p>If a potato crop was infected and detected then it would be destroyed. An infected tomato or grapevine crop might not be destroyed but infected plants identified through testing would be destroyed. Infected plant material would be buried or incinerated. Entire crops may be surveyed or random surveillance may be done because infected plants may be symptomless, especially in the first year of infection or when the infection occurs through the seed (CABI & EPPO 1997d). Tracing and surveillance would be done on other properties that are thought to be at risk of infection. Typically plants, propagating material, machinery and implements may not be moved from properties where a virus outbreak has been detected. Machinery and equipment would be disinfected. Potatoes produced on an infected property would be quarantined. Continued sales of tomato fruit or grapes produced on an infected property might be permitted.</p> <p>Methods used to control nematode-borne viruses include:</p> <ul style="list-style-type: none"> • Use of certified planting material • Soil fumigation with nematicides before planting • Weed control • Avoiding the movement of nematodes with contaminated equipment from an infected to an uninfected field • Removal of infected plants and neighbouring plants followed by spot treatment with nematicides • Crop rotation with crops that are not a host of the virus <p>Plants grown in potato seed certification schemes in Australia are currently inspected for symptoms of virus infection (DAFWA 2009a, b; ViCSPA 2009). Disease tested grapevine, strawberry and raspberry planting material is available in Australia through voluntary certification schemes (Menzies & Brien 2002; Plant Health Australia 2009a; VINA 2008). The virus has been eliminated from infected potato tubers by hot air treatment (Kaiser 1980).</p> |

| Criterion | Estimate and rationale |
|----------------------------------|---|
| Domestic trade | <p>D—Significant at the district level</p> <p>If TBRV became established in an Australian state, restrictions might be introduced on the interstate trade of affected produce and germplasm and, if this occurred it would lead to the loss of markets. As the virus can be transmitted through tomato seed, trade in tomato fruit might be affected. If potato crops were infected, trade in potato tubers from the district might cease. Trade restrictions might be limited to the affected properties, but could be placed on produce from a district, while the pest status of the district was determined and might be placed on trade from a state.</p> <p>Eradication campaigns have been launched in response to every recent outbreak of PSTVd in Australia. No movement of plants or machinery from the affected properties is permitted during the campaigns.</p> |
| International trade | <p>D—Significant at the district level.</p> <p><i>Tomato black ring virus</i> is a regulated pathogen in the North American plant protection organisation territory, as well as New Zealand (Harper et al. 2011). It is regulated for nursery stock, vegetative material, seed and pollen importation in New Zealand (MAF Biosecurity New Zealand 2011). If TBRV became established in Australia, additional restrictions might be introduced on the international trade of nursery stock and propagative material, and possibly some fruit with seed that is possibly infected. This could potentially lead to the loss of international markets and could lead to industry adjustment.</p> <p>Part of the Australian fresh tomato fruit crop is exported, as is a part of the Australian ware potato crop and seed potato crop. These exports might be affected if TBRV becomes established in Australia. Australia has markets for fresh tomatoes to New Zealand, Singapore, Hong Kong, Brunei, Malaysia, New Caledonia, Indonesia, French Polynesia, Fiji, and USA, and markets for potatoes to Republic of Korea, Malaysia, Mauritius, Singapore, Hong Kong, Indonesia, Philippines, United Arab Emirates, Thailand, Taiwan and Brunei, and markets for tomato seed to Thailand and New Zealand (DAFF 2008; HAL 2012).</p> |
| Environmental and non-commercial | <p>Impact score: B—Significant at the local level.</p> <p>The application of nematicides to the soil may affect the environment</p> |

4.2.6 Unrestricted risk estimate

Unrestricted risk is the result of combining the likelihoods of entry, establishment and spread with the outcome of overall consequences. Likelihoods and consequences are combined using the risk estimation matrix shown in Table 2.5.

Unrestricted risk estimate for *Tomato black ring virus*

| | |
|---|---------------|
| Overall likelihood of entry, establishment and spread | Extremely low |
| Consequences | Moderate |
| Unrestricted risk | Negligible |

As indicated, the unrestricted risk estimate for *Tomato black ring virus* has been assessed as 'negligible' which achieves Australia's ALOP. Therefore, no specific risk management measures are required for this pest.

Assessments for quarantine pests for which the URE outcome is adopted from previous assessments

4.3 Grapevine phylloxera

Daktulosphaira vitifoliae (EP)

Daktulosphaira vitifoliae was included in the final import policy for table grapes from China (Biosecurity Australia 2011a), from Korea (Biosecurity Australia 2011b) and from Japan (Department of Agriculture 2014). In these existing policies, the unrestricted risk estimate for *D. vitifoliae* was assessed as exceeding Australia's ALOP and therefore specific risk management measures are required for this pest.

The likelihood of establishment and spread of *D. vitifoliae* in Australia for table grapes from India will be comparable to previous assessments. These likelihoods relate specifically to events that occur in Australia and are principally independent of the importation pathway. The consequences of *D. vitifoliae* are also independent of the importation pathway.

Even though the main import windows differ between table grapes from the previous export areas and India, tissues susceptible to infection by *D. vitifoliae* will be available during the expected import window for table grapes from India as well as during the import windows for table grapes from the previously assessed export areas. Therefore, the likelihood of distribution for this pest for table grapes from India would be comparable to that for table grapes from the export areas assessed previously.

The department considered factors affecting the likelihood of importation for *D. vitifoliae* for table grapes from India and those previously assessed. The likelihood of importation for *D. vitifoliae* for table grapes from India would be comparable to that in the previous assessments.

In addition, the department has also reviewed the latest literature and no new information is available that would significantly change the risk ratings for importation, distribution, establishment, spread and consequences as set out for *D. vitifoliae* in the existing policies.

4.3.1 Unrestricted risk estimate

The unrestricted risk estimate for *D. vitifoliae* for table grapes from India is comparable to the estimates in previous assessments, exceeding Australia's ALOP. Therefore, specific risk management measures are required for this pest.

4.4 European fruit lecanium

***Parthenolecanium corni* (EP, WA)**

Parthenolecanium corni is not present in Western Australia and is a pest of regional quarantine concern for that state.

Parthenolecanium corni was included in the final import policy for table grapes from China (Biosecurity Australia 2011a), from California (DAFF 2013) and from Japan (Department of Agriculture 2014). In these existing policies, the unrestricted risk estimate for *P. corni* was assessed as achieving Australia's ALOP and therefore specific risk management measures are not required for this pest.

The likelihood of establishment and spread of *P. corni* in Western Australia will be comparable to previous assessments. These likelihoods relate specifically to events that occur in Australia and are principally independent of the importation pathway. The consequences of *P. corni* are also independent of the importation pathway.

Parthenolecanium corni has a wide host range and the likelihood of distribution for this pest for table grapes from India would be comparable to that for table grapes from the previously assessed export areas.

The department considered factors affecting the likelihood of importation for *P. corni* for table grapes from India and those previously assessed. The likelihood of importation for *P. corni* for table grapes from India would be comparable to that in the previous assessments. Also, if the likelihood of importation is assessed as 'high' (the possible highest estimate) for *P. corni* for table grapes from India, the unrestricted risk estimate will still achieve Australia's ALOP.

In addition, the department has also reviewed the latest literature and no new information is available that would significantly change the risk ratings for importation, distribution, establishment, spread and consequences as set out for *P. corni* in the existing policies.

4.4.1 Unrestricted risk estimate

The unrestricted risk estimate for *P. corni* for table grapes from India is comparable to the estimates in previous assessments, achieving Australia's ALOP. Therefore, specific risk management measures are not required for this pest.

4.5 Mealybugs

***Planococcus ficus* (EP), *Planococcus lilacinus* (EP), *Planococcus minor* (EP, WA) and *Rastrococcus ickeyroides* (EP)**

Planococcus ficus (grapevine mealybug), *Planococcus lilacinus* (coffee mealybug), *Planococcus minor* (Pacific mealybug) and *Rastrococcus iceryoides* (downey snowline mealybug) belong to the Pseudococcidae or mealybug family. The mealybug species assessed here have been grouped together because of their related biology and taxonomy, and they are predicted to pose a similar risk and to require similar mitigation measures.

Planococcus minor is not present in Western Australia and is a pest of regional quarantine concern for that state.

Several mealybug species were assessed previously in a number of existing import policies, for example, in the final import policy for mangoes from India (Biosecurity Australia 2008), unshu mandarins from Japan (Biosecurity Australia 2009) and pineapples from Malaysia (DAFF 2012). In these existing policies, the unrestricted risk estimate for mealybugs was assessed as exceeding Australia's ALOP and therefore specific risk management measures are required for the pests.

The likelihood of establishment and spread of mealybugs in Australia for table grapes from India will be comparable to previous assessments. These likelihoods relate specifically to events that occur in Australia and are principally independent of the importation pathway. The consequences of mealybugs are also independent of the importation pathway.

Mealybugs have a wide host range and the likelihood of distribution for these pests for table grapes from India would be comparable to that for commodities assessed previously.

The department considered factors affecting the likelihood of importation for mealybugs for table grapes from India and those previously assessed. The likelihood of importation for mealybugs for table grapes from India would be comparable to that in the previous assessments.

In addition, the department has also reviewed the latest literature and no new information is available that would significantly change the risk ratings for importation, distribution, establishment, spread and consequences as set out for mealybugs in the existing policies.

4.5.1 Unrestricted risk estimate

The unrestricted risk estimate for mealybugs for table grapes from India is comparable to the estimates in previous assessments, exceeding Australia's ALOP. Therefore, specific risk management measures are required for these pests.

4.6 Large grape plume moth

Platyptilia ignifera (EP)

Platyptilia ignifera was included in the final import policy for table grapes from Japan (Department of Agriculture 2014). In this existing policy, the unrestricted risk estimate for *P. ignifera* was assessed as achieving Australia's ALOP and therefore specific risk management measures are not required for this pest.

The likelihood of establishment and spread of *P. ignifera* in Australia for table grapes from India will be comparable to previous assessments. These likelihoods relate specifically to events that occur in Australia and are principally independent of the importation pathway. The consequences of *P. ignifera* are also independent of the importation pathway.

Even though the main import windows differ between table grapes from the previous export area and India, tissues susceptible to infection by *P. ignifera* will be available during the expected import window for table grapes from India as well as during the import window for table grapes from the previous export area. Therefore, the likelihood of distribution for this pest for table grapes from India would be comparable to that for table grapes from the previously assessed export area.

The department considered factors affecting the likelihood of importation for *P. ignifera* for table grapes from India and that previously assessed. The likelihood of importation for *P. ignifera* for table grapes from India would be comparable to that in the previous assessment.

In addition, the department has also reviewed the latest literature and no new information is available that would significantly change the risk ratings for importation, distribution, establishment, spread and consequences as set out for *P. ignifera* in the existing policies.

4.6.1 Unrestricted risk estimate

The unrestricted risk estimate for *P. ignifera* for table grapes from India is comparable to the estimates in previous assessments, achieving Australia's ALOP. Therefore, specific risk management measures are not required for this pest.

4.7 Thrips

***Retithrips syriacus* (EP) and *Rhipiphorothrips cruentatus* (EP)**

Retithrips syriacus (black vine thrips) and *Rhipiphorothrips cruentatus* (grapevine thrips) have been grouped together because of their related biology and taxonomy, and they are predicted to pose a similar risk and to require similar mitigation measures. Unless explicitly stated, the term 'thrips' is used to refer to both species and the information presented is considered as applicable to both species.

Several thrips species were assessed previously in a number of existing import policy, for example, in the import policy for persimmon from Israel (DAFF 2004b), mangoes from Taiwan (Biosecurity Australia 2006b) and table grapes from China (Biosecurity Australia 2011a). In these existing policies, the unrestricted risk estimate for thrips was assessed as exceeding Australia's ALOP and therefore specific risk management measures are required for the pests.

The likelihood of establishment and spread of thrips in Australia for table grapes from India will be comparable to previous assessments. These likelihoods relate specifically to events that occur in Australia and are principally independent of the importation pathway. The consequences of thrips are also independent of the importation pathway.

Thrips have a wide host range and the likelihood of distribution for these pests for table grapes from India would be comparable to that for commodities assessed previously.

The department considered factors affecting the likelihood of importation for thrips for table grapes from India and those previously assessed. The likelihood of importation for thrips for table grapes from India would be comparable to that in the previous assessments.

In addition, the department has also reviewed the latest literature and no new information is available that would significantly change the risk ratings for importation, distribution, establishment, spread and consequences as set out for thrips in the existing policies.

4.7.1 Unrestricted risk estimate

The unrestricted risk estimate for thrips for table grapes from India is comparable to the estimates in previous assessments, exceeding Australia's ALOP. Therefore, specific risk management measures are required for these pests.

4.8 Kanzawa spider mite

Tetranychus kanzawai (EP, WA)

Tetranychus kanzawai is not present in Western Australia and is a pest of regional quarantine concern for that state.

Tetranychus kanzawai was included in the final import policy for table grapes from China (Biosecurity Australia 2011a), from Korea (Biosecurity Australia 2011b) and from Japan (Department of Agriculture 2014). In these existing policies, the unrestricted risk estimate for *T. kanzawai* was assessed as exceeding Australia's ALOP and therefore specific risk management measures are required for this pest.

The likelihood of establishment and spread of *T. kanzawai* in Western Australia for table grapes from India will be comparable to previous assessments. These likelihoods relate specifically to events that occur in Australia and are principally independent of the importation pathway. The consequences of *T. kanzawai* are also independent of the importation pathway.

Tetranychus kanzawai has a wide host range and the likelihood of distribution for this pest for table grapes from India would be comparable to that for table grapes from the previous export areas.

The department considered factors affecting the likelihood of importation for *T. kanzawai* for table grapes from India and those previously assessed. The likelihood of importation for *T. kanzawai* for table grapes from India would be comparable to that in the previous assessments.

In addition, the department has also reviewed the latest literature and no new information is available that would significantly change the risk ratings for importation, distribution, establishment, spread and consequences as set out for *T. kanzawai* in the existing policies.

4.8.1 Unrestricted risk estimate

The unrestricted risk estimate for *T. kanzawai* for table grapes from India is comparable to the estimates in previous assessments, exceeding Australia's ALOP. Therefore, specific risk management measures are required for this pest.

4.9 Black rot

***Guignardia bidwellii* (EP)**

Guignardia bidwellii was included in the final import policy for table grapes from China (Biosecurity Australia 2011a) and from Japan (Department of Agriculture 2014). In these existing policies, the unrestricted risk estimate for *G. bidwellii* was assessed as exceeding Australia's ALOP and therefore specific risk management measures are required for this pest.

The likelihood of establishment and spread of *G. bidwellii* in Australia for table grapes from India will be comparable to previous assessments. These likelihoods relate specifically to events that occur in Australia and are principally independent of the importation pathway. The consequences of *G. bidwellii* are also independent of the importation pathway.

Guignardia bidwellii has a wide host range and the likelihood of distribution for this pest for table grapes from India would be comparable to that for table grapes from the previously assessed export areas.

The department considered factors affecting the likelihood of importation for *G. bidwellii* for table grapes from India and those previously assessed. The likelihood of importation for *G. bidwellii* for table grapes from India would be comparable to that in the previous assessments.

In addition, the department has also reviewed the latest literature and no new information is available that would significantly change the risk ratings for importation, distribution, establishment, spread and consequences as set out for *G. bidwellii* in the existing policies.

4.9.1 Unrestricted risk estimate

The unrestricted risk estimate for *G. bidwellii* for table grapes from India is comparable to the estimates in previous assessments, exceeding Australia's ALOP. Therefore, specific risk management measures are required for this pest.

4.10 Brown rot

***Monilinia fructigena* (EP)**

Monilinia fructigena was included in several existing import policies, for example in the policies for apples from China (Biosecurity Australia 2010), for table grapes from China (Biosecurity Australia 2011a) and for table grapes from Japan (Department of Agriculture 2014). In these existing policies, the unrestricted risk estimate for *M. fructigena* was assessed as exceeding Australia's ALOP and therefore specific risk management measures are required for this pest.

The likelihood of establishment and spread of *M. fructigena* in Australia for table grapes from India will be comparable to previous assessments. These likelihoods relate specifically to events that occur in Australia and are principally independent of the importation pathway. The consequences of *M. fructigena* are also independent of the importation pathway.

Monilinia fructigena has a wide host range and the likelihood of distribution for this pest for table grapes from India would be comparable to that for table grapes from the previously assessed export areas.

The department considered factors affecting the likelihood of importation for *M. fructigena* for table grapes from India and those previously assessed. The likelihood of importation for *M. fructigena* for table grapes from India would be comparable to that in the previous assessments.

In addition, the department has also reviewed the latest literature and no new information is available that would significantly change the risk ratings for importation, distribution, establishment, spread and consequences as set out for *M. fructigena* in the existing policies.

4.10.1 Unrestricted risk estimate

The unrestricted risk estimate for *M. fructigena* for table grapes from India is comparable to the estimates in previous assessments, exceeding Australia's ALOP. Therefore, specific risk management measures are required for this pest.

4.11 Grapevine leaf rust

***Phakopsora euvitis* (EP)**

Phakopsora euvitis was included in the existing import policies for table grapes from China (Biosecurity Australia 2011a), Korea (Biosecurity Australia 2011b) and Japan (Department of Agriculture 2014). In these existing policies, the unrestricted risk estimate for *P. euvitis* was assessed as exceeding Australia's ALOP and therefore specific risk management measures are required for this pest.

The likelihood of establishment and spread of *P. euvitis* in Australia for table grapes from India will be comparable to previous assessments. These likelihoods relate specifically to events that occur in Australia and are principally independent of the importation pathway. The consequences of *P. euvitis* are also independent of the importation pathway.

Even though the main import windows differ between table grapes from the previous export areas and India, tissues susceptible to infection by *P. euvitis* will be available during the expected import window for table grapes from India as well as during the import windows for table grapes from the previous export areas. Therefore, the likelihood of distribution after arrival in Australia of *P. euvitis* will be comparable to that for table grapes from the previous export areas.

The department considered factors affecting the likelihood of importation for *P. euvitis* for table grapes from India and those previously assessed. The likelihood of importation for *P. euvitis* for table grapes from India would be comparable to that in the previous assessments.

In addition, the department has also reviewed the latest literature and no new information is available that would significantly change the risk ratings for importation, distribution, establishment, spread and consequences as set out for *P. euvitis* in the existing policies.

4.11.1 Unrestricted risk estimate

The unrestricted risk estimate for *P. euvitis* for table grapes from India is comparable to the estimates in previous assessments, exceeding Australia's ALOP. Therefore, specific risk management measures are required for this pest.

4.12 *Phomopsis* cane and leaf spot

Phomopsis viticola (EP, WA)

Phomopsis viticola is not present in Western Australia and is a pest of regional quarantine concern for that state.

Phomopsis viticola was included in several existing import policies, for example for table grapes from Chile (Biosecurity Australia 2005b), China (Biosecurity Australia 2011a), California to Western Australia (DAFF 2013) and Japan (Department of Agriculture 2014). In these existing policies, the unrestricted risk estimate for *P. viticola* was assessed as achieving Australia's ALOP and therefore no specific risk management measures are required for this pest.

The likelihood of establishment and spread of *P. viticola* in Western Australia for table grapes from India will be comparable to previous assessments. These likelihoods relate specifically to events that occur in Western Australia and are principally independent of the importation pathway. The consequences of *P. viticola* are also independent of the importation pathway.

The likelihood of distribution was recently reassessed for table grapes from California to take account of new information available as well as the differences in the expected import window compared to that assessed previously. Similar to table grapes from California, the main import window for table grapes from India occurs during a period when Australian grapevines are considered less susceptible to infection and climatic conditions in most areas of Western Australia are warm and dry and not conducive to disease development. Therefore, the likelihood of distribution for *P. viticola* for table grapes from India will be comparable to that for table grapes from California to Western Australia.

The department considered factors affecting the likelihood of importation for *P. viticola* for table grapes from India and those previously assessed. The likelihood of importation for *P. viticola* for table grapes from India would be comparable or at least not higher than the highest rating in the previous assessments. Also, if the likelihood of importation is assessed as 'high' (the possible highest rating) for *P. viticola* for table grapes from India, the unrestricted risk estimate will still achieve Australia's ALOP.

In addition, the department has also reviewed the latest literature and no new information is available that would significantly change the risk ratings for importation, distribution, establishment, spread and consequences as set out for *P. viticola* in the existing policies.

4.12.1 Unrestricted risk estimate

The unrestricted risk estimate for *P. viticola* for table grapes from India is comparable to the estimates in previous assessments, achieving Australia's ALOP. Therefore, no specific risk management measures are required for this pest.

Assessments for quarantine pests for which some of the likelihood and/or consequence ratings are adopted from previous assessments

4.13 Fruit flies

***Bactrocera correcta* (EP), *Bactrocera dorsalis* (EP)**

Bactrocera correcta (guava fruit fly) and *Bactrocera dorsalis* (Oriental fruit fly) belong to the fruit fly family Tephritidae which is a group considered to be among the most damaging pests of horticultural crops (Kapoor 2002; White & Elson-Harris 1992). The fruit fly species assessed here have been grouped together because of their related biology and taxonomy, and they are predicted to pose a similar risk and to require similar mitigation measures. Unless explicitly stated, the term 'fruit fly' is used to refer to both species and the information presented is considered as applicable to both species.

Bactrocera correcta is considered a potential pest in India where it often occurs sympatrically with serious pest species such as *B. dorsalis* and *B. zonata* and is a major threat to guava (Kapoor 2002; White & Elson-Harris 1992). In a survey of grapevine pests in India, Mani (1992) recorded *B. correcta* from grape (*Vitis vinifera*).

Bactrocera dorsalis has been recorded from table grapes in China (Chu & Tung 1996; Ye & Liu 2005) and has been observed to attack undamaged grapes in laboratory studies in Japan (Iwaizumi, Kumagai & Katsumata 1994). There are no reports in the literature of *B. dorsalis* attacking grapes in other countries or in India.

Both *B. correcta* and *B. dorsalis* has four life stages: egg, larva, pupa and adult. Eggs are laid below the skin of the host fruit. Hatched larvae feed within the fruit. Pupation occurs in the soil under the host plant (CABI 2012; Christenson & Foote 1960; White & Elson-Harris 1992). They can produce several generations a year, depending on the temperature (CABI 2012; Christenson & Foote 1960).

The risk scenario of concern for *B. correcta* and *B. dorsalis* is the presence of eggs and developing larvae within table grapes.

Bactrocera correcta was assessed in the existing import policy for mangoes from India (Biosecurity Australia 2008). *Bactrocera dorsalis* was assessed in several existing import policies, for example, in the import policies for longan and lychee fruit from China and Thailand (DAFF 2004a) and for table grapes from China (Biosecurity Australia 2011a). The assessment of *B. correcta* and *B. dorsalis* presented here builds upon these previous assessments.

Differences in horticultural practices, climatic conditions and the prevalence of the pests between previously assessed export areas and India make it necessary to reassess the likelihood that *B. correcta* and *B. dorsalis* will be imported into Australia with table grapes from India.

Both *B. correcta* and *B. dorsalis* have a wide range of hosts and the likelihood of distribution after arrival in Australia of these species with table grapes from India would be comparable to that for table grapes from China (Biosecurity Australia 2011a).

The likelihood of establishment and of spread of *B. correcta* and *B. dorsalis* in Australia will be comparable regardless of the fresh fruit commodity in which these species are imported into

Australia, as these likelihoods relate specifically to events that occur in Australia and are independent of the importation pathway. The consequences of *B. correcta* and *B. dorsalis* are also independent of the importation pathway.

In addition, the department has reviewed the latest literature and no new information is available that would significantly change the risk ratings for distribution, establishment, spread and consequences as set out for these species in the existing policies. Therefore, the likelihood ratings for *B. correcta* and *B. dorsalis* in the existing policies will be adopted here.

4.13.1 Likelihood of entry

The likelihood of entry is considered in two parts, the likelihood of importation and the likelihood of distribution, which consider pre-border and post-border issues, respectively.

Likelihood of importation

The likelihood that *B. correcta* and *B. dorsalis* will arrive in Australia with the importation of table grapes from India is: **Very low**.

The following information provides supporting evidence for this assessment.

- Both *B. correcta* and *B. dorsalis* are widely distributed in India. *Bactrocera correcta* is recorded from Andhra Pradesh, Bihar, Goa, Gujarat, Haryana, Himachal Pradesh, Punjab, Karnataka, Kerala, Maharashtra, Tamil Nadu, Uttar Pradesh and West Bengal (CABI 2012). *Bactrocera dorsalis* is recorded from Andhra Pradesh, Assam, Bihar, Delhi, Goa, Himachal Pradesh, Punjab, Jammu and Kashmir, Karnataka, Kerala, Madhya Pradesh, Maharashtra, Manipur, Orissa, Rajasthan, Sikkim, Tamil Nadu, Uttar Pradesh and West Bengal (CABI 2012; Drew & Hancock 1994; Kapoor 2002; Satarkar et al. 2009). Grapes are commercially grown in many of these areas, for example Maharashtra, Karnataka, Andhra Pradesh, Tamil Nadu and Punjab (DPP 2007).
- While both *B. correcta* and *B. dorsalis* have a wide host range, grapes do not appear to be a preferred host for these two fruit fly species. *Bactrocera dorsalis* was recorded in table grapes in China, where it is considered a minor pest of grapevine (Chu & Tung 1996; Ye & Liu 2005). This species has also been observed to attack undamaged grapes in laboratory studies in Japan (Iwaizumi, Kumagai & Katsumata 1994). There are no reports of *B. dorsalis* attacking grapes in India or other countries other than China. In a survey of grapevine pests in India, Mani (1992) recorded *B. correcta* from grape (*Vitis vinifera*). There have been no other reports that associate *B. correcta* with grapevines in India or in other countries.
- *Bactrocera dorsalis* undergoes hibernation during winter in northern India, but is active throughout the year in the southern part of India (Verghese et al. 2002). The overwintering and hibernation patterns of *B. correcta* are assumed to be similar.
- Table grapes are produced in the northern, central, and southern areas of India (DPP 2007).
- The main harvest season of table grapes in India is from February to the end of April (DAFF 2010; DPP 2012). Small volumes of table grapes may be harvested at other times of the year. There is potential for these fruit fly species to be active and could be present in Indian table grape vineyards during the harvest season.

- Studies in China, on other hosts, have demonstrated that *B. correcta* adults will oviposit at temperatures as low as 18 degrees Celsius (Liu & Ye 2009). Temperatures during the harvest season for Indian table grapes are likely to exceed 18 degrees Celsius (Shikhamany 2001).
- Damage caused by fruit flies consists of punctures of the host tissue by adults during oviposition and feeding by the larvae within the fruit pulp (Christenson & Foote 1960; Ye & Liu 2005). This allows for secondary infection from fungi or bacteria, which cause extensive rotting and dropping of fruit (Mau & Martin Kessing 2007).
- In the absence of blemishes and damage to the skin, eggs and early instar larvae, if present, are unlikely to be detected during picking, sorting and quality inspection. However, fruit that show obvious signs of attack or tissue decay, particularly those with secondary infection by fungi and/or bacteria, are likely to be removed from the export pathway during harvesting, sorting and packing processes.
- Cold temperature treatments of 1.7 degrees Celsius for 14 days or 1.0 degrees Celsius for 13 days killed third instar larvae (the most cold tolerant life-stage) in naturally-infested citrus and longans (Wu 2005). Armstrong *et al.* (1995) demonstrated that cold temperature treatment at 1.1 degrees Celsius for 12 days was sufficient to kill *B. dorsalis* eggs and larvae in carambolas. There is no specific research data available on the lethal effects of cold storage on *B. correcta*.
- Grapes are usually pre-cooled to less than 4 degrees Celsius after packing and cold stored at 0–2 degrees Celsius at 90–95 per cent humidity after palletizing until shipment (DPP 2009). Conditions are maintained at 0–2 degrees Celsius and 90–95 per cent humidity throughout transport (DAFF 2010; DPP 2009, 2012). These storage and transportation conditions, without a specific cold treatment, are likely to be only sub-lethal to *B. correcta* and *B. dorsalis* eggs and larvae.

The wide distribution of *B. correcta* and *B. dorsalis* in India including in the main grape production areas, the possibility that some infested fruit without obvious signs of infestation may escape detection during harvesting, sorting and packing processes, moderated by the information that there are no reports of *B. dorsalis* attacking grapes in India, the single report of *B. correcta* in grapes in India is from 1999 and not since and there have been no reports of *B. correcta* in grape in any other countries, support a likelihood estimate for importation of ‘very low’

Likelihood of distribution

The likelihood of distribution for *B. correcta* and *B. dorsalis* has been adopted from the assessment of *B. dorsalis* for table grapes from China (Biosecurity Australia 2011a), that is **Moderate**.

Overall likelihood of entry

The overall likelihood of entry is determined by combining the likelihood of importation with the likelihood of distribution using the matrix of rules shown in Table 2.2.

The likelihood that *B. correcta* and *B. dorsalis* will enter Australia as a result of trade in table grapes from India and be distributed in a viable state to a susceptible host is: **Very low**.

4.13.2 Likelihood of establishment and spread

The likelihood of establishment and of spread for *B. correcta* and *B. dorsalis* has been adopted from the assessment of *B. dorsalis* for longan and lychee from China and Thailand (DAFF 2004a). The ratings from the previous assessment are:

| | |
|-----------------------------|-------------|
| Likelihood of establishment | High |
| Likelihood of spread | High |

4.13.3 Overall likelihood of entry, establishment and spread

The overall likelihood of entry, establishment and spread is determined by combining the likelihoods of entry, of establishment and of spread using the matrix of rules shown in Table 2.2.

The overall likelihood that *B. correcta* and *B. dorsalis* will enter Australia as a result of trade in table grapes from India, be distributed in a viable state to a susceptible host, establish in Australia and subsequently spread within Australia is: **Very low**.

4.13.4 Consequences

The consequences of *B. correcta* and *B. dorsalis* in Australia have been adopted from the previous assessment of *B. dorsalis* for longan and lychee from China and Thailand (DAFF 2004a), that is: **High**.

4.13.5 Unrestricted risk estimate

Unrestricted risk is the result of combining the likelihoods of entry, establishment and spread with the outcome of overall consequences. Likelihoods and consequences are combined using the risk estimation matrix shown in Table 2.5.

Unrestricted risk estimate for *Bactrocera correcta* and *Bactrocera dorsalis*

| | |
|---|----------|
| Overall likelihood of entry, establishment and spread | Very low |
| Consequences | High |
| Unrestricted risk | Low |

As indicated, the unrestricted risk estimate for *Bactrocera correcta* and *Bactrocera dorsalis* has been assessed as 'low', which exceeds Australia's ALOP. Therefore, specific risk management measures are required for these pests.

4.14 Leaf rolling moth

*Archips machlopi*s

*Archips machlopi*s (leaf rolling moth) belongs to the Tortricidae or the leafroller family. Leaf rolling moth larvae are known to damage fruit of several economic plant species by chewing large holes that usually allows entry to fungi that cause fruit rot (CABI 2012).

*Archips machlopi*s has frequently been misidentified in the literature as *Archips micaceana* (Meijerman & Ulenberg 2011; Robinson, Tuck & Shaffer 1994; Rose & Pooni 2004; Tuck 1990). *Archips micaceana* is found in China, Hong Kong, South Vietnam, Burma and Northern Thailand. *Archips machlopi*s is found in Pakistan, Nepal, India, Burma, Thailand, North Vietnam, China (Jiangxi), Malaysia, Sumatra and Java (Meijerman & Ulenberg 2011; Tuck 1990).

While older literature reported that *A. micaceana* was present in India (Puttarudriah, Kataoihallimath & Chandrasekhar 1961), recent literature refers the leaf rolling moth present in India as *A. machlopi*s (Rose & Pooni 2004).

*Archips machlopi*s has four life stages: egg, larva (caterpillar), pupa and adult (moth) (Puttarudriah, Kataoihallimath & Chandrasekhar 1961).

Adults of this species lay eggs on their hosts, or on glass surfaces in enclosures, in varying egg masses of 7 to 58. The eggs are held together by a glutinous material and take about eight days to hatch (Puttarudriah, Kataoihallimath & Chandrasekhar 1961).

When fully grown, the larva is about 20 millimetres long (Puttarudriah, Kataoihallimath & Chandrasekhar 1961).

The webbing or silken shelters are also used as a pupation site. The pupae are broad, around 10 millimetres long and 3 millimetres wide and take about a week to develop (Puttarudriah, Kataoihallimath & Chandrasekhar 1961).

Adult moths are inactive during the day, but will fly away when disturbed. They become active after dusk and the males are attracted to bright light (Puttarudriah, Kataoihallimath & Chandrasekhar 1961). Egg laying commences during the night, approximately two days after the moths emerge from the pupae (Puttarudriah, Kataoihallimath & Chandrasekhar 1961).

No information was found about the number of generations per year *A. machlopi*s can produce. However, a related species, *Archips podana*, has one generation per year in northern and central Europe, two generations per year in the south of the Republic of Belarus and Ukraine, and three generations per year in the Caucasus and Transcaucasia (CABI 2014; Meijerman & Ulenberg 2000; Ovsyannikova & Grichanov 2009a).

The risk scenario of concern for *Archips machlopi*s is the presence of larvae on imported grape bunches.

Archips micaceana was assessed in the existing import policy for table grapes from China (Biosecurity Australia 2011a). *Archips machlopi*s has similar biology to *A. micaceana* and the two species are predicted to pose a similar risk and to require similar mitigation measures. The assessment of *A. machlopi*s presented here builds upon the previous assessment for *A. micaceana*.

Differences in horticultural practices, climatic conditions and the prevalence of the pests between the previously assessed export areas and India make it necessary to reassess the likelihood that *A. machlopi*s will be imported into Australia with table grapes from India.

Similar to *A. micaceana*, *A. machlopi*s has a wide range of hosts and the likelihood of distribution after arrival in Australia of *A. machlopi*s will be comparable to that for *A. micaceana* for table grapes from the previous export area (Biosecurity Australia 2011a).

The likelihood of establishment and of spread of *A. machlopi*s in Australia will be comparable regardless of the fresh fruit commodity in which this species is imported into Australia, as these likelihoods relate specifically to events that occur in Australia and are independent of the importation pathway. The consequences of *A. machlopi*s are also independent of the importation pathway.

In addition, the department has reviewed the latest literature and no new information is available that would significantly change the risk ratings for distribution, establishment, spread and consequences as set out for *A. micaceana* in the existing policy. Therefore, those likelihood ratings and consequences estimate will be adopted for this assessment.

4.14.1 Likelihood of entry

The likelihood of entry is considered in two parts, the likelihood of importation and the likelihood of distribution, which consider pre-border and post-border issues, respectively.

Likelihood of importation

The likelihood that *A. machlopi*s will arrive in Australia with the importation of table grapes from India is: **Low**.

The following information provides supporting evidence for this assessment.

- *Archips machlopi*s is present in Karnataka, Kerala, Punjab, Uttaranchal and Himachal Pradesh (Puttarudriah, Kataoihallimath & Chandrasekhar 1961; Rose & Pooni 2004; Varma 1984). Karnataka is India's second biggest grape producing state (APEDA 2015). Grapes are also grown commercially in Punjab (APEDA 2015; DPP 2007).
- Within Karnataka, *A. machlopi*s has been reported on grapevine, including on grape bunches, in Kenchanahalli, Bangalore city, Byatarayanapura and Yelahanka, with noticeable damage occurring during November and December (Puttarudriah, Kataoihallimath & Chandrasekhar 1961).
- Puttarudriah (1961) reports that larvae were found in harvested grape bunches brought to a domestic market.
- The larvae feed under thin webbing on the epidermis of the leaves as well as on parts of the grape bunch, namely on the rachis, pedicels and both immature and mature grape berries (Puttarudriah, Kataoihallimath & Chandrasekhar 1961).
- Feeding by the larvae on the rachis and pedicels causes the berries to become unattached from the bunch and shrivel up (Puttarudriah, Kataoihallimath & Chandrasekhar 1961). Feeding by the larvae on berries causes cavities to form at the base of the affected berries (Puttarudriah, Kataoihallimath & Chandrasekhar 1961). Cavities usually allow entry for

fungi that cause fruit rot (CABI 2012). Affected bunches with many shrivelled berries, many berries with cavities and/or rotten berries are likely to be detected and removed from the export pathway during harvesting, sorting and packing processes.

- Larvae of *A. machlopi*s are of a noticeable size, about 20 millimetres long when fully grown (Puttarudriah, Kataoihallimath & Chandrasekhar 1961), and silk webbing and frass are likely to be present on affected bunches. This increases the likelihood of affected bunches being detected and removed from the export pathway during harvesting, sorting and packing processes.
- No information was found about where on the host plant eggs of *A. machlopi*s are laid. Moths of the Tortricidae family, for example *Lobesia botrana*, *Epiphyas postvittana*, *Archips podana*, *Platynota stultana*, *Eupoecilia ambiguella* and *Sparganothis pilleriana*, often lay their eggs on the leaves, shoots, buds or on or near flower clusters of grapevine (Bentley et al. 2008; INRA 1998; Loch 2007; Zalom, Varela & Cooper 2011). Eggs of *Argyrotaenia franciscana* are laid on any smooth surface of the grapevine plant such as upper leaf surfaces, stems, canes or berries (Flaherty et al. 1992). Second generation of *E. ambiguella* and second and third generation of *L. botrana* lay their eggs on the berries (INRA 1998; Zalom, Varela & Cooper 2011). Egg masses or young larvae, if present on grape bunches at harvest, would be more difficult to detect.
- Packed grapes are transported in cold humidified storage to ensure grape quality is maintained. In India, grapes are usually pre-cooled to less than 4 degrees Celsius after packing and cold stored at 0–2 degrees Celsius at 90–95 per cent humidity after palletizing until shipment (DPP 2009). Conditions are maintained at 0–2 degrees Celsius and 90–95 per cent humidity throughout transport (DAFF 2010; DPP 2009, 2012). Leafroller larvae can survive cold conditions experienced during refrigerated transport, but survival rate, for example for *Platynota stultana*, decreases to around 6 per cent after two weeks at less than 1 degree Celsius (Yokoyama & Miller 2000).

The records of *A. machlopi*s causing damage in some vineyards in grape production regions in India, moderated by the information that affected grape bunches are likely to be detected and removed from the export pathway due to the conspicuous nature of damage caused by this pest and the limited ability of leafroller larvae to survive more than two weeks of cold storage, support a likelihood estimate for importation of 'low'.

Likelihood of distribution

The likelihood of distribution for *A. machlopi*s has been adopted from the assessment of *A. micaceana* for table grapes from China (Biosecurity Australia 2011a), that is **Moderate**.

Overall likelihood of entry

The overall likelihood of entry is determined by combining the likelihood of importation with the likelihood of distribution using the matrix of rules shown in Table 2.2.

The likelihood that *A. machlopi*s will enter Australia as a result of trade in table grapes from India and be distributed in a viable state to a susceptible host is: **Low**.

4.14.2 Likelihood of establishment and spread

The likelihood of establishment and of spread for *A. machlopi*s has been adopted from the assessment for *A. micaceana* for table grapes from China (Biosecurity Australia 2011a). The ratings from the previous assessment are:

| | |
|-----------------------------|-------------|
| Likelihood of establishment | High |
| Likelihood of spread | High |

4.14.3 Overall likelihood of entry, establishment and spread

The overall likelihood of entry, establishment and spread is determined by combining the likelihoods of entry, of establishment and of spread using the matrix of rules shown in Table 2.2.

The overall likelihood that *A. machlopi*s will enter Australia as a result of trade in table grapes from India, be distributed in a viable state to a susceptible host, establish in Australia and subsequently spread within Australia is: **Low**.

4.14.4 Consequences

The consequences of *A. machlopi*s have been adopted from the assessment for *A. micaceana* for table grapes from China (Biosecurity Australia 2011a), that is: **Moderate**.

4.14.5 Unrestricted risk estimate

Unrestricted risk is the result of combining the likelihoods of entry, establishment and spread with the outcome of overall consequences. Likelihoods and consequences are combined using the risk estimation matrix shown in Table 2.5.

Unrestricted risk estimate for *Archips machlopi*s

| | |
|---|----------|
| Overall likelihood of entry, establishment and spread | Low |
| Consequences | Moderate |
| Unrestricted risk | Low |

As indicated, the unrestricted risk estimate for *Archips machlopi*s has been assessed as 'low', which is above Australia's ALOP. Therefore, specific risk management measures are required for this pest.

4.15 Citrus pyralid

Cryptoblabes gnidiella (EP)

Cryptoblabes gnidiella (citrus pyralid) is part of the Pyralidae family, which is a large and ubiquitous family of insects which are common pests of cultivated plants and stored products (Nielsen & Common 1991). *Cryptoblabes gnidiella* is native to the Mediterranean region, but has since spread throughout Europe, Africa, tropical and subtropical areas of the Americas, New Zealand and parts of Asia (CABI 2012; Carter 1984).

Adult *C. gnidiella* have a body length of around 7.5 millimetres and a wing expanse of 14 millimetres (Singh & Singh 1995). *Cryptoblabes gnidiella* lays about 100 eggs on the fruit or on the foliage of host plants and eggs hatch in around 4–7 days (Carter 1984). In India, eggs laid on sorghum are reported to hatch in around 3–6 days (Singh & Singh 1995). There are five larval instars, with the duration of the larval period being 13 days (CABI 2012). Fully grown larvae are around 12 millimetres long (CABI 2012). The pupae are around 7 millimetres in length, with the pupal stage lasting eight days (CABI 2012). The period from egg to adult is around 27 days (CABI 2012). The pre-ovipositional period lasts a full day after mating, with most eggs laid during the first night (Wysoki, Yehuda & Rosen 1993). There are three or four generations a year in southern Europe, up to five in North Africa (Carter 1984) and nine in India (Singh & Singh 1995).

Cryptoblabes gnidiella is capable of causing direct damage to a range of host plants, with larval feeding on foliage and fruits causing direct crop losses. It is a polyphagous species associated with about 60 plant species in more than 30 families (Bagnoli & Lucchi 2001). Its commercial hosts include *Allium sativum* (garlic), *Annona muricata* (soursop), *Citrus* spp., *Coffea* spp., *Ficus* spp., *Gossypium* spp., *Macadamia* spp., *Malus* spp., *Mangifera indica* (mango), *Oryza sativa* (rice), *Persea americana* (avocado), *Phaseolus* spp., (beans), *Prunus* spp., *Punica granatum* (pomegranate), *Pyrus* spp., *Ricinus communis* (castor bean), *Saccharum officinarum* (sugarcane), *Sorghum bicolor* (sorghum), *Vaccinium* sp. (blueberry), *Vitis* spp. (grapevine) and *Zea mays* (maize) (Bagnoli & Lucchi 2001; CABI 2012; Silva & Mexia 1999; Wysoki 1986; Yehuda, Wysoki & Rosen 1991).

Larvae of *C. gnidiella* mainly attack the fruit of host plants, but also feed on foliage, bark and twigs (DPP 2012; Liotta & Mineo 1964). Larvae are often found in association with infestations of other pests. For example on citrus, *C. gnidiella* larvae are often found where there is infestation of the mealybug *Planococcus citri* and on grapes they are often found following attack by the European grapevine moth, *Lobesia botrana* (Carter 1984). *Cryptoblabes gnidiella* is attracted to honeydew created by mealybugs (Swirski et al. 1980; Zimmerman 1958).

In India, *C. gnidiella* is most active from March to November and overwinters in the pupal stage with the onset of cold weather (Singh & Singh 1995). There are reports of *C. gnidiella* causing serious damage to hybrid sorghum in India (Sharma, Singh & Nwanze 1997; Singh & Singh 1995). There are no reports of *C. gnidiella* attacking grapevine in India. However, *C. gnidiella* has been recorded as a pest of grapevine in Israel (Harari et al. 2007), Brazil (Bisotto-de-Oliveira et al. 2007) and Italy (Bagnoli & Lucchi 2001). Where it is found in grapevine, *C. gnidiella* larvae can feed on dry flower parts, berry stalks, berry juice as well as sound grapes, they feed mainly on the rachis and stalks of the growing green grape bunches (Bagnoli & Lucchi 2001).

The risk scenario of concern for *C. gnidiella* is that larvae and eggs may be imported with grape bunches.

Cryptoblabes gnidiella was assessed in the existing import policy for sweet oranges from Italy (Biosecurity Australia 2005a). The assessment of *C. gnidiella* presented here builds upon this previous assessment.

Differences in commodities, horticultural practices, climatic conditions and the prevalence of *C. gnidiella* between the previous export area and India make it necessary to reassess the likelihood that *C. gnidiella* will be imported into Australia with table grapes from India.

Cryptoblabes gnidiella has multiple hosts and the likelihood of distribution after arrival in Australia of *C. gnidiella* for table grapes from India will be comparable to that for sweet oranges from Italy.

The likelihood of establishment and of spread of *C. gnidiella* in Australia will be comparable regardless of the fresh fruit commodity in which this species is imported into Australia, as these likelihoods relate specifically to events that occur in Australia and are principally independent of the importation pathway. The consequences of *C. gnidiella* are also independent of the importation pathway.

In addition, the department has reviewed the latest literature and no new information is available that would significantly change the risk ratings for distribution, establishment, spread and consequences as set out for *C. gnidiella* in the existing policy. Therefore, those likelihood ratings and consequence estimates will be adopted for this assessment.

4.15.1 Likelihood of entry

The likelihood of entry is considered in two parts, the likelihood of importation and the likelihood of distribution, which consider pre-border and post-border issues, respectively.

Likelihood of importation

The likelihood that *C. gnidiella* will arrive in Australia with the importation of table grapes from India is: **Very low**.

The following information provides supporting evidence for this assessment.

- *Cryptoblabes gnidiella* is reported present in Karnataka, Maharashtra and Uttar Pradesh (CABI 2012; DPP 2012; Zhang 1994). Maharashtra and Karnataka are the two major grape growing states of India (APEDA 2015).
- *Cryptoblabes gnidiella* is reported as a pest of grapevine in Israel (Harari et al. 2007), Brazil (Bisotto-de-Oliveira et al. 2007) and Italy (Bagnoli & Lucchi 2001).
- In India, *C. gnidiella* is recorded as a serious pest of sorghum and also attacks *Azolla* spp. (aquatic ferns), but there are no reports of this pest associated with grapevine in India (Sasmal & Kulshreshtha 1985; Singh & Singh 1995).
- Where it is found in grapevine, larvae of *C. gnidiella* are often found in association with *Lobesia botrana* and the control of *C. gnidiella* in vineyards can be achieved through the

effective control of *L. botrana* (Bagnoli & Lucchi 2001). *Lobesia botrana* is currently not recorded in India.

- Larvae of *C. gnidiella* can feed on many parts of grapevine including dry flower parts, berry stalk, berry juice as well as sound grape, but they mainly feed on the rachis and stalks of growing green grape bunches (Bagnoli & Lucchi 2001). It is likely that the feeding of larvae on rachis and stalks of growing green grape bunches will result in berries/bunches not developing normally and/or dropping prematurely.
- In addition to direct damage through feeding of larvae on grape bunches, *C. gnidiella* also causes indirect damage including bunch decay due to fungal infection (Harari et al. 2007; Silva & Mexia 1999).
- Affected grape bunches with several berries not fully developed and/or missing are likely to be detected and removed from the export pathway during the harvesting, sorting and packing processes. Similarly, bunches with decay from fungal infection are likely to be removed from the export pathway.
- Fully grown larvae of *C. gnidiella* are 12 millimetres long (Singh & Singh 1995) and are likely to be detected and removed from the export pathway during harvesting, sorting and packing processes.
- *Cryptoblabes gnidiella* is reported to lay around 100 eggs on the fruit or foliage of host plants and eggs hatch in 4-7 days (Carter 1984). Eggs are between 0.7 and 0.4 millimetres and first instar larvae are one millimetre in size (Ioriatti, Lucchi & Varela 2012). Egg masses or young larvae, if present on grape bunches at harvest, would be more difficult to detect.
- The period from egg to adult in *Cryptoblabes gnidiella* is only around 27 days with eggs laid on sorghum in India reported to hatch in around 3–6 days (CABI 2012; Singh & Singh 1995). Due to the short development times of *C. gnidiella* life stages, the period when eggs and young larvae may be present on grape bunches is expected to be very short. Grapes are usually stored and transported at low temperatures to prolong shelf life. Low temperature is expected to delay egg and larval development. Therefore, it could be expected that some, but not all, eggs and young larvae, if present on exported grape bunches, are unlikely to develop much further before arriving in Australia. Those developed further would likely be detected at routine inspection on arrival.

The presence of *C. gnidiella* in the major grape growing states in India, moderated by the lack of reports of *C. gnidiella* affecting grapevine in India and the fact that, if affected, the affected grape bunches are likely to be removed from the export pathway, support a likelihood estimate for importation of 'very low'.

Likelihood of distribution

The likelihood of distribution for *C. gnidiella* has been adopted from the assessment of *C. gnidiella* for sweet oranges from Italy (Biosecurity Australia 2005a), that is **Low**.

Overall likelihood of entry

The overall likelihood of entry is determined by combining the likelihood of importation with the likelihood of distribution using the matrix of rules shown in Table 2.2.

The likelihood that *C. gnidiella* will enter Australia as a result of trade in table grapes from India and be distributed in a viable state to a susceptible host is: **Very low**.

4.15.2 Likelihood of establishment and spread

The likelihood of establishment and of spread for *C. gnidiella* has been adopted from the assessment of *C. gnidiella* for sweet oranges from Italy (Biosecurity Australia 2005a). The ratings from the previous assessment are:

| | |
|-----------------------------|-------------|
| Likelihood of establishment | High |
| Likelihood of spread | High |

4.15.3 Overall likelihood of entry, establishment and spread

The overall likelihood of entry, establishment and spread is determined by combining the likelihoods of entry, of establishment and of spread using the matrix of rules shown in Table 2.2.

The overall likelihood that *C. gnidiella* will enter Australia as a result of trade in table grapes from India, be distributed in a viable state to a susceptible host, establish in Australia and subsequently spread within Australia is: **Very low**.

4.15.4 Consequences

The consequences of *C. gnidiella* have been adopted from the assessment of *C. gnidiella* for sweet oranges from Italy (Biosecurity Australia 2005a), that is: **Moderate**.

4.15.5 Unrestricted risk estimate

Unrestricted risk is the result of combining the likelihoods of entry, establishment and spread with the outcome of overall consequences. Likelihoods and consequences are combined using the risk estimation matrix shown in Table 2.5.

Unrestricted risk estimate for *Cryptoblabes gnidiella*

| | |
|---|----------|
| Overall likelihood of entry, establishment and spread | Very low |
| Consequences | Moderate |
| Unrestricted risk | Very low |

As indicated, the unrestricted risk estimate for *Cryptoblabes gnidiella* has been assessed as 'very low', which achieves Australia's ALOP. Therefore, no specific risk management measures are required for this pest.

4.16 Bitter rot

***Greeneria uvicola* (EP, WA)**

Greeneria uvicola is not present in Western Australia and is a pest of regional quarantine concern for that state.

Bitter rot of grapevine is caused by the fungus *Greeneria uvicola*. The disease occurs on many *Vitis* spp. including *Vitis vinifera*, *V. labrusca*, *V. aestivalis*, *V. bourquina*, *V. rotundifolia* and *V. munsoniana* (Farr et al. 2001; Longland & Sutton 2008; Ridings & Clayton 1970) under warm and humid conditions (Farr et al. 2001; McGrew 1988). Bitter rot disease is, however, more severe on muscadine grapes (*V. rotundifolia*) (McGrew 1988). Under experimental conditions, the fungus has also been shown to infect wounded fruit of apple, cherry, strawberry, peach, blueberry and banana causing fruit rot (Ridings & Clayton 1970). However, *G. uvicola* is not known to cause problems on horticultural crops other than grapes.

In Australia, *G. uvicola* is known to be present in New South Wales (NSW) and Queensland (Qld) (Castillo-Pando et al. 2001; Castillo-Pando, Steel & Somers 1999; Plant Health Australia 2001; Sergeeva, Nair & Spooner-Hart 2001) but has not been recorded in Western Australia (DAWA 2006a; Taylor 2012) and is a pest of quarantine concern for that state.

The fungus can infect young shoots, leaves, tendrils, peduncle, rachis, pedicels and fruit of grapevine (Ellis 2008; Kummuang et al. 1996b). It has also been isolated from dormant canes, wood and bark (Castillo-Pando et al. 2001; Emmett 2006). *Greeneria uvicola* has been reported to cause girdling of shoots, flecking of young leaves, stems, shoots and individual flower buds (Kummuang et al. 1996b; McGrew 1988; Momol, Ritchie & Dankers 2007; Tashiro 1992).

Although *G. uvicola* can infect many different tissues of grapevine, the disease mainly damages fruit, particularly if rainy weather persists into the harvest season (Farr et al. 2001).

There are mixed reports on at what developmental stage berries are susceptible to infection. The incidence of bitter rot disease for muscadine grapes (*V. rotundifolia*) on non-sprayed vines was reported to be more severe on young berries and decreased drastically thereafter (Kummuang et al. 1996b). The authors stated that bitter rot symptoms had already been observed on some flower buds. Steel *et al.* (2012) reported on their inoculation study that inflorescences of Chardonnay grapes (*V. vinifera*) were also susceptible to infection by *G. uvicola*, and infection of inflorescences at mid-flowering led to berry rot at veraison. There are also reports to suggest that grapes of several *V. vinifera* cultivars become more susceptible to infection after veraison (Steel 2007; Steel, Greer & Savocchia 2007). In inoculation studies conducted over two years using three *V. vinifera* cultivars, it was reported that the susceptibility of grapes increased from bloom until veraison in one year, and from bloom until two weeks before veraison in the other year (Longland & Sutton 2008).

Reports on the timing of first symptoms on berries also vary. For muscadine grapes on non-sprayed field-grown vines, the development of symptoms varies between different muscadine cultivars and vineyard locations, but disease symptoms were most prevalent on all cultivars at the young berry stage (Kummuang et al. 1996b). The authors also reported that *G. uvicola* was isolated from symptomless berries, especially those late in the growing season (Kummuang et al. 1996b). McGrew (1988) and Momol *et al.* (2007) reported that greenish

brown lesions can be found on young muscadine berries as well as blight of pedicels, which causes the young berries to shrivel and break off. However, these same authors also stated, but did not mention on what type of grapes, that *G. uvicola* invades corky lenticular warts which form on the pedicel in the spring (shortly after flowering) but remains latent until the berry reaches maturity. The fungus then invades the pedicel and moves into the berry, where conidia are produced within four days (McGrew 1988; Momol, Ritchie & Dankers 2007). It is unknown if these reports (McGrew 1988; Momol, Ritchie & Dankers 2007) were based on field (vineyard) observation or results of inoculation studies. Longland and Sutton (2008) reported on their inoculation studies, where grapes were inoculated from bloom until two weeks before harvest, that symptoms were not observed until just before harvest.

Greeneria uvicola overwinters on mummified berries, damaged shoot tips, infected senescent and fallen leaves, and necrotic bark (Farr et al. 2001; Kummuang et al. 1996a; Momol, Ritchie & Dankers 2007; Smith 2012). The optimum temperature for infection is reported to be around 28–30 degrees Celsius (McGrew 1988; Momol, Ritchie & Dankers 2007; Ridings & Clayton 1970; Sutton & Gibson 1977; Taylor 2012). Even though one author stated that transmission of the fungus is via air-borne conidia (Sutton & Gibson 1977), most authors agree that conidia of *G. uvicola* are spread by rain splash (Ellis 2008; Kummuang et al. 1996a; MAFF 2008; Smith 2012).

The risk scenario of concern for *G. uvicola* is that symptomless infected grape bunches may be imported into Western Australia.

Greeneria uvicola was included in the existing import policy for table grapes from Japan (Department of Agriculture 2014). The assessment of *G. uvicola* presented here builds on this existing policy.

Differences in horticultural practices, climatic conditions and the prevalence of the pest between previously assessed export area (Japan) and India make it necessary to reassess the likelihood that *G. uvicola* will be imported into Western Australia with table grapes from India.

Due to the differences in the main import window and the expected import volume between table grapes from Japan and table grapes from India, the likelihood of distribution of *G. uvicola* after arrival in Western Australia with table grapes from India is reassessed here.

The likelihood of establishment and of spread of *G. uvicola* in Western Australia will be comparable regardless of the fresh fruit commodity in which this species is imported into Western Australia, as these likelihoods relate specifically to events that occur in Western Australia and are independent of the importation pathway. The consequences of *G. uvicola* are also independent of the importation pathway.

In addition, the department has reviewed the latest literature and no new information is available that would significantly change the risk ratings for establishment, spread and consequences as set out for *G. uvicola* in the existing policy. Therefore, those risk ratings will be adopted for this assessment.

4.16.1 Likelihood of entry

The likelihood of entry is considered in two parts, the likelihood of importation and the likelihood of distribution, which consider pre-border and post-border issues, respectively.

Likelihood of importation

The likelihood that *G. uvicola* will arrive in Western Australia with the importation of table grapes from India is: **High**.

The following information provides supporting evidence for this assessment.

- *Greeneria uvicola* has been recorded in Andhra Pradesh, Bihar (Reddy & Reddy 1983) and Karnataka (Ullasa & Rawal 1986). Andhra Pradesh and Karnataka are commercial grape production areas expected to export grapes to Australia (DPP 2009).
- *Greeneria uvicola* infects grape clusters (McGrew 1988). On young berries, symptoms first develop as brown lesions (Milholland 1991) or flecks (Kummuang et al. 1996b). Severe infection can cause blight on young berries and pedicels which causes young berries to shrivel and drop (Kummuang et al. 1996b; McGrew 1988; Momol, Ritchie & Dankers 2007).
- On maturing berries, the fungus causes brownish, water-soaked lesions, with concentric rings of spore bodies, which rapidly spread and eventually cover the entire berry (Ellis 2008; Momol, Ritchie & Dankers 2007; Taylor 2012). Black, raised acervuli form on the decaying fruit which can cause the epidermis and cuticle to rupture (McGrew 1988; Momol, Ritchie & Dankers 2007). Some infected berries soften and detach easily from the bunch, particularly in wet weather, whilst others continue to dry and shrivel (McGrew 1988; Momol, Ritchie & Dankers 2007; Taylor 2012; Ullasa & Rawal 1986). Grape bunches with several berries missing, or with several shrivelled berries, are likely to be discarded at harvesting or packing processes.
- Symptoms of infection are easily recognised on the berries and are reported to develop on healthy berries one week after contact with fungal spores and in less time on damaged fruit (Castillo-Pando, Steel & Somers 1999; Ellis 2008). However, one study which pinned bitter-rotted berries onto healthy bunches did not result in infection of adjacent non-wounded berries (Ridings & Clayton 1970). Infected grape berries/bunches showing obvious symptoms are likely to be removed from the export pathway during harvesting or packing processes. It has also been reported that grapes inoculated with *G. uvicola* from bloom to two weeks before harvest did not show symptoms until just close to harvest (Longland & Sutton 2008). Some authors report that *G. uvicola* invades pedicels of grapes in the spring (shortly after flowering) but remains latent until the berry reaches maturity (McGrew 1988; Momol, Ritchie & Dankers 2007). The fungus then invades the berries, where conidia are produced within four days (McGrew 1988). Kummuang *et al.* (1996b) also reported that *G. uvicola* was isolated from symptomless berries, especially those late in the growing season. Infected grape bunches without or with only mild symptoms at harvest may escape detection and be picked and packed for export.
- The fungus can invade any injured tissue of *Vitis* spp. plants (McGrew 1988). Injury to mature, healthy berries due to bird and insect damage or cracking of berries due to rain can allow conidial infection and lead to rapid spread of the disease (McGrew 1988; Momol, Ritchie & Dankers 2007). Damaged grape berries/bunches are likely to be removed from the export pathway during harvesting or packing processes.
- The varieties known to be naturally infected in India are Anab-e-Shahi, Angur Kalan, Black Champa, Gulabi, Jaos Beli, Kali Sahabi, Khandari, Pandri Sahebi, Selection 94, Thompson

Seedless and Taifi Rosovi (Reddy & Reddy 1983). Some of these varieties are likely to be exported to Australia (DPP 2007, 2009).

- Measures used to control *G. uvicola* in India include pruning of infected canes (NHB 2011).
- Bitter rot symptoms develop quickly on mature berries. It could be expected that any berries with latent infection that were picked and packed for export via sea freight would show symptoms by the time they arrive in Western Australia. Grape bunches showing symptoms would be detected during routine inspection on arrival. However, grapes are usually stored at low temperatures to prolong shelf life. Information on the time required for symptoms to develop under cold storage conditions could not be found, but it is likely that symptoms will develop more slowly under low temperatures. Grapes via air freight may show no or mild symptoms at the time they arrive in Western Australia. Grape bunches without symptoms, or with only minor symptoms, may not be detected at routine inspection on arrival.

The possibility for some late infected berries to show no or mild symptoms and the uncertainty about the development of symptoms at low temperatures support a likelihood estimate for importation of 'high'.

Likelihood of distribution

The likelihood that *G. uvicola* will be distributed within Western Australia in a viable state as a result of the processing, sale or disposal of table grapes from India and subsequently transfer to a susceptible part of a host is: **Low**.

The following information provides supporting evidence for this assessment.

- Imported grapes are intended for human consumption. Distribution of the imported grapes would be for retail sale.
- As grapes are easily damaged during handling (Mencarelli, Bellincontro & DiRenzo 2005), packed grapes may not be processed or handled again until they arrive at the retailers. Therefore, pathogens in packed grapes are unlikely to be detected during transportation and distribution to retailers.
- Bitter rot symptoms develop quickly on mature berries. It could be expected that infected berries would show symptoms by the time they arrive at the retailers. Grape bunches with obvious symptoms of infection would not be marketable and would not be sold. However, if grapes are transported at low temperatures, symptoms may develop more slowly. Grape bunches without symptoms, or with only minor symptoms, could be marketable and could be sold.
- Most fruit waste will be discarded into managed waste systems and will be disposed of in municipal tips and would therefore pose little risk of exposure to a suitable host.
- Consumers will discard small quantities of fruit waste in urban, rural and natural localities. Small amounts of fruit waste will be discarded in domestic compost. There is some potential for consumer waste being discarded near host plants, including commercially grown, household or wild host plants. If present in fruit waste, the pathogen would then need to be transferred to a susceptible host.

- The primary host of *G. uvicola* is *Vitis rotundifolia*, but other *Vitis* spp. are also susceptible including *V. vinifera*, *V. bourquina*, *V. labrusca* and *V. munsoniana* (Farr et al. 2001; Longland & Sutton 2008; Ridings & Clayton 1970). No other natural hosts are known. While it was reported more than 40 years ago that, under experimental conditions, *G. uvicola* can infect wounded fruit of apple, cherry, strawberry, peach, blueberry and banana (Ridings & Clayton 1970), there have been no reports found on natural infection on these plant species.
- In Western Australia, *Vitis* spp. are grown commercially and are also common garden plants (ABS 2009a; ATGA 2013; Kiri-ganai Research Pty Ltd 2006; Waldecks 2013).
- Commercial table grape vineyards in Western Australia are located near the Western Australian coast, extending from the Gascoyne region (including Carnarvon) to the South-West region (including Harvey, Donnybrook, Margaret River and Busselton) (DAWA 2006b). The main wine grape production spans from Gingin just north of Perth, extending through the south-west and across to the Porongurup Range near Mount Baker (DAFWA 2006).
- Even though one author reports that transmission of the fungus is via air-borne conidia (Sutton & Gibson 1977), most authors agree that conidia of *G. uvicola* are spread by rain splash (Ellis 2008; Kummuang et al. 1996a; MAFF 2008; Smith 2012). In wet conditions, conidia present on the surface of infected grape bunches could be transmitted via rain splash and wind-driven rain to susceptible nearby host plants.
- The fungus can infect young shoots, leaves, tendrils, peduncle, rachis, pedicels and fruit of grapevine (Ellis 2008; Kummuang et al. 1996b). It has also been isolated from dormant canes, wood and bark (Castillo-Pando et al. 2001; Emmett 2006). The fungus can invade any injured tissue of *Vitis* spp. plants (McGrew 1988).
- The main export season for table grapes from India to Australia will be from February to the end of April (DAFF 2010; DPP 2012) (the end of summer to mid autumn in Australia). However small volumes of table grapes may come in at other times of the year. Grapevines in Western Australia would be susceptible to infection during the expected export window. Other hosts of the assessed fungi may also be susceptible to infection during the expected export window.

The host susceptibility during the expected export window, moderated by the limited range of potential conidia dispersal via rain splash and the limited host range support a likelihood estimate for distribution of 'low'.

Overall likelihood of entry

The overall likelihood of entry is determined by combining the likelihood of importation with the likelihood of distribution using the matrix of rules shown in Table 2.2.

The likelihood that *G. uvicola* will enter Western Australia as a result of trade in table grapes from India and be distributed in a viable state to a susceptible host is: **Low**.

4.16.2 Likelihood of establishment and spread

The likelihood of establishment and of spread for *G. uvicola* within Western Australia has been adopted from the assessment of *G. uvicola* for table grapes from Japan (Department of Agriculture 2014). The ratings from the previous assessment are:

| | |
|-----------------------------|------------|
| Likelihood of establishment | Low |
| Likelihood of spread | Low |

4.16.3 Overall likelihood of entry, establishment and spread

The overall likelihood of entry, establishment and spread is determined by combining the likelihoods of entry, of establishment and of spread using the matrix of rules shown in Table 2.2.

The overall likelihood that *G. uvicola* will enter Western Australia as a result of trade in table grapes from India, be distributed in a viable state to a susceptible host, establish in Western Australia and subsequently spread within Western Australia is: **Very low**.

4.16.4 Consequences

The consequences of *G. uvicola* in Western Australia have been adopted from the assessment of *G. uvicola* for table grapes from Japan (Department of Agriculture 2014), that is: **Low**.

4.16.5 Unrestricted risk estimate

Unrestricted risk is the result of combining the likelihoods of entry, establishment and spread with the outcome of overall consequences. Likelihoods and consequences are combined using the risk estimation matrix shown in Table 2.5.

Unrestricted risk estimate for *Greeneria uvicola*

| | |
|---|------------|
| Overall likelihood of entry, establishment and spread | Very low |
| Consequences | Low |
| Unrestricted risk | Negligible |

As indicated, the unrestricted risk estimate for *Greeneria uvicola* has been assessed as 'negligible', which achieves Australia's ALOP. Therefore, no specific risk management measures are required for this pest.

4.17 Fruit rot

***Pestalotiopsis menezesiana* (EP, WA), *Pestalotiopsis uvicola* (EP, WA)**

Pestalotiopsis menezesiana and *Pestalotiopsis uvicola* are plant pathogenic fungi that cause fruit rot of grapevine (Mishra, Prakash & Misra 1974; Xu et al. 1999).

Pestalotiopsis menezesiana and *P. uvicola* are assessed together as the two species cause a similar disease and their biology is likely to be the same or very similar; and they are predicted to cause a similar risk and would be managed by similar mitigation measures if required. Unless explicitly stated, the information presented is considered as applicable to both species. In this section, the common name fruit rot is used to refer to both species. The scientific name is used when the information is about a specific species.

Pestalotiopsis menezesiana and *Pestalotiopsis uvicola* are not present in Western Australia and is a pest of regional quarantine concern for that state.

Pestalotiopsis menezesiana and *P. uvicola* are not known to be present in Western Australia and are pests of quarantine concern for that state. In Australia, *P. menezesiana* is known to be present in NSW (Plant Health Australia 2001; Sergeeva, Priest & Nair 2005) and *P. uvicola* in NSW and Qld (Plant Health Australia 2001).

On *Vitis* spp., both assessed fungi have mainly been reported on *Vitis vinifera* (Guba 1961; Kobayashi 2007). *Pestalotiopsis uvicola* has also been reported on *V. coignetia*, *V. indivisa* and *V. labrusca* (Farr & Rossman 2013b; Guba 1961; Kobayashi 2007). Both fungi have been reported on leaves, canes and fruit of *Vitis* spp. (Bissett 1982; Guba 1961; MAFF 2008; Mishra, Prakash & Misra 1974; Mundkur & Thirumalachar 1946; Nag Raj 1993; Sergeeva, Priest & Nair 2005). *Pestalotiopsis uvicola* has also been isolated from flowers, cankers and internal wood rot of grapevine, and has been associated with grapevine trunk disease (Sergeeva, Priest & Nair 2005; Úrbez-Torres et al. 2009; Úrbez-Torres et al. 2012).

In addition to *Vitis* spp., *P. menezesiana* has also been reported to cause leaf spot of kiwifruit (*Actinidia chinensis*) and plantain (*Musa paradisiaca*), and rot of cuttings of grape ivy (*Cissus rhombifolia*) (Bissett 1982; Huang et al. 2007; Park et al. 1997). *Pestalotiopsis uvicola* has been reported to cause leaf spot and stem blight of bay laurel (*Laurus nobilis*), stem blight of Kermadac pohutukawa (*Metrosideros kermadecensis*) and leaf spot of mango (*Mangifera indica*) and carob (*Ceratonia siliqua*) (Carrieri, Carotenuto & Lahoz 2013; Grasso & Granata 2008; Ismail, Cirvilleri & Polizzi 2013; Vitale & Polizzi 2005).

Pestalotiopsis menezesiana, like many other species of *Pestalotiopsis*, has also been reported on dead or dying plant material (Guba 1961; Nag Raj 1993) and both assessed fungi have been isolated as endophytes on conifer trees in China (Liu, Xu & Guo 2007; Liu et al. 2013).

Infection of *Pestalotiopsis* spp. can occur from a resting endophytic stage, mycelium, ascospores or conidium on healthy tissue (Maharachchikumbura et al. 2011). The infection develops into enlarging, circular to irregular lesions that contain either pycnidia or perithecia. Spores are then released to continue the infection (Maharachchikumbura et al. 2011). However, the sexual stage does not often develop and thus conidia (asexual spores) are thought to provide the inocula (Maharachchikumbura et al. 2011).

The risk scenario of concern for the assessed fungi is that symptomless infected grape bunches may be imported into Western Australia.

Pestalotiopsis menezesiana and *P. uvicola* were included in the existing import policy for table grapes from Japan (Department of Agriculture 2014). The assessment of *P. menezesiana* and *P. uvicola* presented here builds on this existing policy.

Differences in horticultural practices, climatic conditions and the prevalence of the pests between previously assessed export area (Japan) and India make it necessary to reassess the likelihood that *P. menezesiana* and/or *P. uvicola* will be imported into Western Australia with table grapes from India.

Differences in the main import window and the expected import volume between table grapes from Japan and table grapes from India make it necessary to reassess the likelihood of distribution of *P. menezesiana* and *P. uvicola* after arrival in Western Australia.

The likelihood of establishment and of spread of *P. menezesiana* and *P. uvicola* in Western Australia will be comparable regardless of the fresh fruit commodity in which these species are imported into Western Australia, as these likelihoods relate specifically to events that occur in Western Australia and are independent of the importation pathway. The consequences of *P. menezesiana* and *P. uvicola* are also independent of the importation pathway.

In addition, the department has reviewed the latest literature and no new information is available that would significantly change the risk ratings for establishment, spread and consequences as set out for *P. menezesiana* and *P. uvicola* in the existing policy. Therefore, those likelihood ratings will be adopted for this assessment.

4.17.1 Likelihood of entry

The likelihood of entry is considered in two parts, the likelihood of importation and the likelihood of distribution, which consider pre-border and post-border issues, respectively.

Likelihood of importation

The likelihood that *P. menezesiana* and/or *P. uvicola* will arrive in Western Australia with the importation of table grapes from India is: **Moderate**.

The following information provides supporting evidence for this assessment.

- *Pestalotiopsis menezesiana* and *P. uvicola* are reported present on grapevine in India (Farr & Rossman 2015). *Pestalotiopsis menezesiana* has been reported present on grape berries in Bihar in 1970 (Mishra, Prakash & Misra 1974). *Pestalotiopsis uvicola* has been reported on *Acacia* from Kerala (Mohanan et al. 2005). Neither of these states are major table grape producing states (APEDA 2015; DPP 2007, 2012). However, Bihar neighbours with Uttar Pradesh and Kerala neighbours with Tamil Nadu, both of which have commercial grape growing areas (DPP 2012).
- On grapevine, grape bunches are among tissues which can be infected by the assessed fungi (Bissett 1982; MAFF 2008; Mishra, Prakash & Misra 1974; Xu et al. 1999).
- When grape berries of different maturity stages were punctured and inoculated with *P. menezesiana*, the rates of infection were: 20.0 per cent for raw berries, 93.3 per cent for

semi-ripe berries and 55.3 per cent for fully ripe berries (Mishra, Prakash & Misra 1974). Symptoms of infection develop more quickly on mature berries. Rot symptoms were visible after four days for ripe berries and after nine days for semi-ripe berries (Mishra, Prakash & Misra 1974).

- Infection also occurred on uninjured berries inoculated with the assessed fungi (Mishra, Prakash & Misra 1974; Xu et al. 1999), but at lower infection rates compared to injured berries (Mishra, Prakash & Misra 1974). Inoculation onto healthy uninjured berries caused rot after four days at 25 degrees Celsius for *P. menezesiana* and after two weeks at the same temperature for *P. uvicola* (Xu et al. 1999), suggesting that the pathogenicity of *P. uvicola* on grape berries might be weaker than that of *P. menezesiana*.
- Inoculation studies with injured grape berries indicate that colony formation/growth of the assessed fungi and decay of berries seems to be highest at the temperature range of 20-30 degrees Celsius (Xu et al. 1999).
- Symptoms on grape clusters are obvious. In India, symptoms of *P. menezesiana* first appear near the peduncle when the fruit is about to ripen and cover the upper portion of the fruit within two days (Mishra, Prakash & Misra 1974). The lesions first appear water-soaked and then turn Sienna colour (yellow brown or reddish brown) with numerous acervuli (Mishra, Prakash & Misra 1974). Lesions are irregular and the acervuli are raised in severe cases (Mishra, Prakash & Misra 1974). The skin of the berry becomes brownish-black and leathery, and bunches become completely unmarketable (Mishra, Prakash & Misra 1974).
- Diseased grape clusters showing obvious symptoms are likely to be removed from the export pathway during harvesting and/or packing processes.
- In Japan, the assessed fungi have been isolated from healthy tissue of both mature and immature grape bunches in the vineyard and were also detected on damaged fruit in markets (Xu et al. 1999). Although the authors suggested that these fungi could potentially cause latent infection and a post-harvest disease of grapes (Xu et al. 1999) they did not report if latent infection still occurs at harvest or investigate the condition at harvest of the damaged fruit in markets where the assessed fungi were isolated from.
- As symptoms of the assessed fungi develop quickly on mature berries, it could be expected that any infected berries that were picked and packed for export via sea freight would show symptoms by the time they arrive in Western Australia. Grape bunches showing symptoms would be detected during routine inspection on arrival.
- However, grapes are usually stored and transported at low temperatures to prolong shelf life. Detailed information on the time for symptoms to develop under cold storage conditions could not be found, but the study by Xu *et al.* (1999) indicates that symptoms develop more slowly at low temperatures. Grape bunches without symptoms, or with only minor symptoms, may not be detected at routine inspection on arrival.

The possibility that the assessed fungi could be present on grape bunches without symptoms, or with only minor symptoms, at harvest, moderated by the fact that symptoms of the assessed fungi develop quickly on mature berries and that grape bunches showing obvious symptoms are likely to be removed from the export pathway, support a likelihood estimate for importation of 'moderate'.

Likelihood of distribution

The likelihood that *P. menezesiana* and/or *P. uvicola* will be distributed within Western Australia in a viable state as a result of the processing, sale or disposal of table grapes from India and subsequently transfer to a susceptible part of a host is: **Low**.

The following information provides supporting evidence for this assessment.

- Imported grapes are intended for human consumption. Distribution of the imported grapes would be for retail sale.
- As grapes are easily damaged during handling (Mencarelli, Bellincontro & DiRenzo 2005), packed grapes may not be processed or handled again until they arrive at the retailers. Therefore, pathogens in packed grapes are unlikely to be detected during transportation and distribution to retailers.
- It could be expected that infected berries would show symptoms by the time they arrive at the retailers. Grape bunches with obvious symptoms of infection would not be marketable and would not be sold. If grapes are transported at low temperatures, symptoms may develop more slowly. Grape bunches without symptoms, or with only minor symptoms, could be marketable and sold.
- Most fruit waste will be discarded into managed waste systems and will be disposed of in municipal tips and would therefore pose little risk of exposure to a suitable host.
- Consumers will discard small quantities of fruit waste in urban, rural and natural localities. Small amounts of fruit waste will be discarded in domestic compost. There is some potential for consumer waste being discarded near host plants, including commercially grown, household or wild host plants. If present in fruit waste, the assessed fungi would then need to be transferred to a susceptible host.
- *Pestalotiopsis menezesiana* and *P. uvicola* reproduce through conidia (Bissett 1982; Guba 1961). Conidia are produced at 13–28 degrees Celsius, with the most conidia produced at 22 degrees Celsius (Huang et al. 2007). Similar to other Coelomycetes with appendage bearing conidia, conidia of the genus *Pestalotiopsis* are dispersed by rain splash or wind-blown droplets (MAFF 2008; Maharachchikumbura et al. 2011; Nag Raj 1993).
- If present in fruit waste, the conidia would then need to be transferred from the fruit waste in water droplets to susceptible host tissue. This transmission is limited to a short distance for fruit waste on the ground.
- For both fungi, germination of conidia occurred at 10–33 degrees Celsius and no germination was observed at 35 degrees Celsius or higher (Xu et al. 1999). Optimum temperature for germination of conidia was 25 degrees Celsius for *P. menezesiana* and 23–25 degrees Celsius for *P. uvicola* (Xu et al. 1999).
- *Pestalotiopsis menezesiana* overwintered in diseased leaves of kiwifruit on the ground in Korea (Park et al. 1997).
- Members of the genus *Pestalotiopsis* are generally not very host specific (Maharachchikumbura et al. 2011). The known hosts of *P. menezesiana* include *Actinidia chinensis* (kiwifruit) (Park et al. 1997), *Cissus rhombifolia* (grape-ivy) (Bissett 1982), *Vitis vinifera* (grapevine) (Mishra, Prakash & Misra 1974; Xu et al. 1999) and

Musa paradisiaca (plantain) (Huang et al. 2007). The known hosts of *P. uvicola* include *Ceratonia siliqua* (carob) (Carrieri, Carotenuto & Lahoz 2013), *Laurus nobilis* (bay laurel) (Vitale & Polizzi 2005), *Macadamia integrifolia* (macadamia nut), *Mangifera indica* (mango) (Ismail, Cirvilleri & Polizzi 2013), *Metrosideros kermadecensis* (Kermadec pohutukawa) (Grasso & Granata 2008), *Vitis coignetia*, *V. indivisa*, *V. labrusca* and *V. vinifera* (Farr & Rossman 2013b; Guba 1961; Kobayashi 2007; Xu et al. 1999). Many of these hosts are grown in Western Australia, some of these are grown commercially such as grapevine, mango and kiwifruit.

- The main export season for table grapes from India to Australia will be from February to the end of April (DAFF 2010; DPP 2012) (the end of summer to mid autumn in Australia). However, small volumes of table grapes may come in at other times of the year. Grapevines in Western Australia would be susceptible to infection during the expected export window. Other hosts of the assessed fungi may also be susceptible to infection during the expected export window.
- In inoculation studies, *P. menezesiana* was able to form fungal colonies at temperatures as low as 5 degrees Celsius in four days (Xu et al. 1999). Cooling of grape bunches during transport and storage is unlikely to affect the viability of the assessed fungi.

The availability of host plants in Western Australia, moderated by the limited potential for dispersal of conidia via rain splash from fruit waste to a susceptible part of a host, the short time required for symptoms to develop on mature bunches and subsequent removal of such bunches from being sold, support a likelihood estimate for distribution of 'low'.

Overall likelihood of entry

The overall likelihood of entry is determined by combining the likelihood of importation with the likelihood of distribution using the matrix of rules shown in Table 2.2.

The likelihood that *P. menezesiana* and/or *P. uvicola* will enter Western Australia as a result of trade in table grapes from India and be distributed in a viable state to a susceptible host is: **Low**.

4.17.2 Likelihood of establishment and spread

The likelihood of establishment and spread for *P. menezesiana* and *P. uvicola* within Western Australia has been adopted from the assessment of *P. menezesiana* and *P. uvicola* for table grapes from Japan (Department of Agriculture 2014). The ratings from the previous assessment are:

| | |
|-----------------------------|-------------|
| Likelihood of establishment | High |
| Likelihood of spread | High |

4.17.3 Overall likelihood of entry, establishment and spread

The overall likelihood of entry, establishment and spread is determined by combining the likelihoods of entry, of establishment and of spread using the matrix of rules shown in Table 2.2.

The overall likelihood that *P. menezesiana* and/or *P. uvicola* will enter Western Australia as a result of trade in table grapes from India, be distributed in a viable state to a susceptible host, establish in Western Australia and subsequently spread within Western Australia is: **Low**.

4.17.4 Consequences

The consequences of *P. menezesiana* and *P. uvicola* in Western Australia have been adopted from the assessment of *P. menezesiana* and *P. uvicola* for table grapes from Japan (Department of Agriculture 2014), that is: **Low**.

4.17.5 Unrestricted risk estimate

Unrestricted risk is the result of combining the likelihoods of entry, establishment and spread with the outcome of overall consequences. Likelihoods and consequences are combined using the risk estimation matrix shown in Table 2.5.

| Unrestricted risk estimate for <i>Pestalotiopsis menezesiana</i> and <i>Pestalotiopsis uvicola</i> | |
|---|----------|
| Overall likelihood of entry, establishment and spread | Low |
| Consequences | Low |
| Unrestricted risk | Very low |

As indicated, the unrestricted risk estimate for *Pestalotiopsis menezesiana* and *Pestalotiopsis uvicola* has been assessed as 'very low', which achieves Australia's ALOP. Therefore, no specific risk management measures are required for these pests.

4.18 White rot

***Pilidiella castaneicola* (EP, WA), *Pilidiella diplodiella* (EP, WA)**

Pilidiella castaneicola (synonym *Coniella castaneicola*) and *Pilidiella diplodiella* (synonyms *Coniella diplodiella*, *Coniothyrium diplodiella*) are plant pathogenic fungi which cause white rot, also known as hail disease, of grapevine (Bisiach 1988; Kishi 1998; Yamato 1995).

Pilidiella castaneicola and *P. diplodiella* are assessed together as the two species cause the same disease and their biology is likely to be very similar, and they are predicted to pose a similar risk and would be managed by similar mitigation measures if required. Unless explicitly stated, the information presented is considered as applicable to both species. In this section, the common name white rot is used to refer to both species. The scientific name is used when the information is about a specific species.

Pilidiella castaneicola and *Pilidiella diplodiella* is not present in Western Australia and is a pest of regional quarantine concern for that state.

Pilidiella castaneicola and *P. diplodiella* are not known to be present in Western Australia and are pests of quarantine concern for that state. *Pilidiella castaneicola* is known to be present on a number of hosts, but not on grapevine, in NSW, NT, Qld and Vic. (Langrell, Glen & Alfnas 2008; Plant Health Australia 2001). *Pilidiella diplodiella* is known to be present on grapevine in NSW and Qld (Plant Health Australia 2001; Simmonds 1966). White rot of grapevine caused by *P. diplodiella* is rare in Australia and of little economic significance (Sergeeva 2010).

Pilidiella castaneicola and *P. diplodiella* affect peduncle, rachis, pedicel and berries of grapevine (Bisiach 1988; Kishi 1998; Yamato 1995). *Pilidiella diplodiella* is known to infect both young and mature grape berries (Lauber & Schuepp 1968). The assessed fungi are unable to infect intact grape berries directly (Bisiach 1988; Kishi 1998). Infection of intact berries occurs through the pedicel and progresses through the subepidermal layers of the berry (Bisiach & Viterbo 1973; Locci & Quaroni 1972). Peduncle, rachis and pedicel can be directly infected by the pathogens without wounding and symptoms progress down towards the berries (Bisiach & Viterbo 1973; Kishi 1998; Locci & Quaroni 1972). If conditions are favourable, the disease can also spread from an infected, injured berry through the pedicel to the rachis and lead to the decay of a major portion of the grape cluster (Bisiach & Viterbo 1973; Lauber & Schuepp 1968).

Pilidiella diplodiella is also known to cause cankers in nonlignified shoots of grapevine but it rarely infects leaves (Bisiach 1988).

The risk scenario of concern for *P. castaneicola* and/or *P. diplodiella* is that symptomless infected grape bunches may be imported into Western Australia.

Pilidiella castaneicola and *P. diplodiella* were included in the existing import policy for table grapes from Japan (Department of Agriculture 2014). The assessment of *P. castaneicola* and *P. diplodiella* presented here builds on this existing policy.

Differences in horticultural practices, climatic conditions and the prevalence of the pests between previously assessed export area (Japan) and India make it necessary to reassess the likelihood that *P. castaneicola* and/or *P. diplodiella* will be imported into Western Australia with table grapes from India.

Due to the differences in the main import window and the expected import volume between table grapes from Japan and table grapes from India, the likelihood of distribution of *P. castaneicola* and *P. diplodiella* after arrival in Western Australia with table grapes from India is reassessed here.

The likelihood of establishment and of spread of *P. castaneicola* and *P. diplodiella* in Western Australia will be comparable regardless of the fresh fruit commodity in which these species are imported into Western Australia, as these likelihoods relate specifically to events that occur in Australia and are independent of the importation pathway. The consequences of *P. castaneicola* and *P. diplodiella* are also independent of the importation pathway.

In addition, the department has reviewed the latest literature and no new information is available that would significantly change the risk ratings for establishment, spread and consequences as set out for *P. castaneicola* and *P. diplodiella* in the existing policy. Therefore, those risk ratings will be adopted for this assessment.

4.18.1 Likelihood of entry

The likelihood of entry is considered in two parts, the likelihood of importation and the likelihood of distribution, which consider pre-border and post-border issues, respectively.

Likelihood of importation

The likelihood that *P. castaneicola* and/or *P. diplodiella* will arrive in Western Australia with the importation of table grapes from India is: **Moderate**.

The following information provides supporting evidence for this assessment.

- *Pilidiella diplodiella* has been reported on *Vitis vinifera* and *Vitis* sp. in India (Farr & Rossman 2015) whereas *P. castaneicola* has only been reported on *Eucalyptus* sp. and *Quercus* sp. (Nag Raj 1993). However, in other countries, including Japan, *P. castaneicola* has also been reported on *Vitis* species (Kobayashi 2007; Nag Raj 1993; Yamato 1995).
- *Pilidiella castaneicola* has been reported present in Karnataka and Himachal Pradesh (Nag Raj 1993). *Pilidiella diplodiella* has been reported present in Bihar and Uttar Pradesh (CABI 2013). Karnataka is one of the major table grape producing states (DPP 2007, 2012). Uttar Pradesh is also known to produce table grapes (DPP 2012).
- White rot of grapevine caused by *P. diplodiella* is common in areas that are prone to hailstorms (Bisiach 1988). In the absence of hailstorms, summer rain followed by persistent high humidity combined with temperatures of 24–27 degrees Celsius can also lead to disease outbreaks (Bisiach 1988). These climatic conditions are expected to be available in some of the export production regions in India (see Figure 2). However, no reports were found on the economic significance of *P. diplodiella* on grapevine in India. Sutton and Waterstone (1964) note that white rot of grapevine caused by *P. diplodiella* is sporadic.
- A number of studies and reports indicate that infections and outbreaks of white rot caused by *P. diplodiella* often seem to occur before or at veraison. Bisiach (1988) notes that typical symptoms of white rot are found on grape clusters before veraison, a few days after a hailstorm. During a study conducted in Slovakia, *P. diplodiella* was isolated frequently from grape berries at early veraison and to a lesser extent at ripening before harvest (Mikusova et

al. 2012). A study by David and Rafaila (1966) shows that attack by *P. diplodiella* increases with the increase in sugar content in the grape berries up to 3.0–3.5 per cent and that above 8 per cent sugar content, no fructification occurs in the attacked area.

- Berries infected with *P. diplodiella* turn yellow and later blue, lose their turgor and become covered with brown to violet pycnidia, which, when mature, turn white/grey (Bisiach 1988; Lauber & Schuepp 1968). The berries dry out and fall to the ground at the end of the season (Bisiach 1988; Lauber & Schuepp 1968). Berries on infected immature clusters turn pale green, become limp and later turn brown (Bisiach 1988). Symptoms on peduncle, rachis and pedicel begin as small, pale brown, elongated depressions, which may spread in favourable conditions (Bisiach 1988). If a lesion occurs on the rachis, the proportion of the cluster below the lesion dries quickly (Bisiach 1988). Symptoms of *P. castaneicola* and *P. diplodiella* on grapevine differ only slightly (Yamato 1995).
- Grape bunches may become contaminated with conidia of *P. diplodiella* when contaminated soil is splashed onto the vine by heavy rain, hail or machinery (Bisiach 1988; Kishi 1998). Under favourable conditions, conidia will germinate on the grape bunch and initiate infection (Bisiach 1988).
- Incubation periods of the assessed fungi can vary with temperature, humidity, means of penetration and the tissue infected from three to eight days (Bisiach 1988; Bisiach & Viterbo 1973; Kishi 1998). Infection of grapevine by *P. diplodiella* is favoured by warm temperatures and high relative humidity (Bisiach 1988). Disease development of *P. diplodiella* occurs rapidly at temperatures of 24–27 degrees Celsius, slowly at temperatures below 15 degrees Celsius and only slightly above 34 degrees Celsius (Bisiach 1988; Locci & Quaroni 1972). Infection is negligible if the temperatures are below 15 degrees Celsius for 24–48 hours following a hailstorm (Bisiach 1988).
- Grape bunches with obvious symptoms of infection are likely to be removed during routine harvesting, grading and packing processes and would not be packed for export. Grape bunches without symptoms, or with only minor symptoms such as small lesions on peduncle, rachis or pedicel could still be exported.
- Grapes are usually stored and transported at low temperatures to prolong shelf life. Conidia of *P. diplodiella* germinate and initiate infection slowly at temperatures below 15 degrees Celsius (Bisiach 1988). As a result, symptoms will develop more slowly under low temperatures. Some infected grapes may exhibit no or mild symptoms at the time they arrive in Western Australia. Grape bunches without symptoms, or with only minor symptoms, may not be detected at routine inspection on arrival.

A possibility for some infected grape bunches showing no or mild symptoms, moderated by the lack of reports of the economic importance of the disease in India, the short time required for symptom development and the obvious symptoms of infection on berries, support a likelihood estimate for importation of 'moderate'.

Likelihood of distribution

The likelihood that *P. castaneicola* and/or *P. diplodiella* will be distributed within Western Australia in a viable state as a result of the processing, sale or disposal of table grapes from India and subsequently transfer to a susceptible part of a host is: **Moderate**.

The following information provides supporting evidence for this assessment.

- Imported grapes are intended for human consumption. Distribution of the imported grapes would be for retail sale.
- As grapes are easily damaged during handling (Mencarelli, Bellincontro & DiRenzo 2005), packed grapes may not be processed or handled again until they arrive at the retailers. Therefore, pathogens in packed grapes are unlikely to be detected during transportation and distribution to retailers.
- It could be expected that infected berries would show symptoms by the time they arrive at the retailers. Grape bunches with obvious symptoms of infection would not be marketable and would not be sold. However, if grapes are transported at low temperatures, symptoms may develop more slowly. Grape bunches without symptoms or with only minor symptoms could be marketable and could be sold.
- Most fruit waste will be discarded into managed waste systems and will be disposed of in municipal tips and would therefore pose little risk of exposure to a suitable host.
- Consumers will discard small quantities of fruit waste in urban, rural and natural localities. Small amounts of fruit waste will be discarded in domestic compost. There is some potential for consumer waste being discarded near host plants, including commercially grown, household or wild host plants. If present in fruit waste, the assessed fungi would then need to be transferred to a susceptible host.
- *Pilidiella castaneicola* has a variety of hosts including *Acer* sp., *Carya* sp., *Castanea* spp., *Eucalyptus* spp., *Fragaria* sp., *Liquidambar styracifolia* (sweet gum), *Metrosideros* sp., *Mangifera indica* (mango), *Quercus alba* (white oak), *Q. rubra* (red oak), *Quercus* sp., *Rhus copallina* (black sumac), *Rhus* sp., *Rosa rugosa-prostrata*, *Vitis cordifolia* and *V. vinifera* (Farr & Rossman 2012; Nag Raj 1993). Some of these hosts are widely distributed in Western Australia.
- *Vitis vinifera* is the principle host of *P. diplodiella* (Bisiach 1988; Van Niekerk et al. 2004). This fungus has also been reported to cause a disease on *Hibiscus sabdariffa* (Roselle) and *Artabotrys hexapetalos* (Ylang Ylang Vine) (Sánchez et al. 2011; Shreemali 1973). Single reports have been found for *P. diplodiella* on *Rosa* sp., *Geranium* sp. and *Anogeissus latifolia* (buttontree) in India and *Citrus aurantiifolia* (lime) in Mexico (Farr & Rossman 2013a; Singh & Sinch 1966). In Western Australia, *Vitis* spp. are grown commercially and are also common garden plants (ABS 2009a; Kiri-ganai Research Pty Ltd 2006; Waldecks 2013). Other possible hosts may also be available in Western Australia including *Hibiscus sabdariffa*, *Rosa* sp., *Geranium* sp. and *Citrus aurantiifolia*. *Hibiscus sabdariffa* is regarded an environmental weed and has widely naturalised in northern Western Australia (University of Queensland 2011).
- *Pycnidia* and conidia of *P. diplodiella* overwinter on dead plant material and in the soil in vineyards (Bisiach 1988). Dried pycnidia of *P. diplodiella* remain able to produce viable conidia for more than 15 years and released conidia remain viable for two to three years (Bisiach 1988). Pycnidia and conidia of the assessed fungi are likely to survive storage and transport.
- Conidia of the assessed fungi are dispersed over short distances by water splash from infected plant material or contaminated soil (Bisiach 1988; Kishi 1998; Sutton & Waterston

1964). In wet conditions, conidia present on the surface of infected grape bunches or fruit waste could be transmitted via rain splash and wind-driven rain to susceptible nearby host plants.

- Infection of grapevine by *P. diplodiella* is favoured by warm temperatures and high relative humidity (Bisiach 1988). Germination of conidia and development of infection progress rapidly at 24–27 degrees Celsius, slowly below 15 degrees Celsius and only slightly above 34 degrees Celsius (Bisiach 1988; Locci & Quaroni 1972). Infection is negligible if the temperatures are below 15 degrees Celsius for 24–48 hours following a hailstorm (Bisiach 1988). In laboratory studies, infection with *P. diplodiella* was stimulated by high relative humidity (90–100 per cent) (David & Rafaila 1966).
- The main export season for table grapes from India to Australia will be from February to the end of April (DAFF 2010; DPP 2012) (the end of summer to mid autumn in Australia). However small volumes of table grapes may come in at other times of the year. The assessed fungi can infect rachis, pedicel and berries of grapevine (Bisiach 1988; Kishi 1998; Yamato 1995). The fungus rarely infects grapevine leaves, but on some cultivars, it can also infect non lignified shoots (Bisiach 1988). Grapevines in Western Australia would be susceptible to infection during the expected export window. Other hosts of the assessed fungi may also be susceptible to infection during the expected export window.

The wide distribution of a number of hosts in Western Australia, the host susceptibility during the expected export window and the ability for pycnidia and conidia of at least one of the assessed fungi, *P. diplodiella*, to remain viable for a long period of time on dead plant material and in the soil, moderated by the limited range of potential conidia dispersal via rain splash, support a likelihood estimate for distribution of ‘moderate’.

Overall likelihood of entry

The overall likelihood of entry is determined by combining the likelihood of importation with the likelihood of distribution using the matrix of rules shown in Table 2.2.

The likelihood that *P. castaneicola* and/or *P. diplodiella* will enter Western Australia as a result of trade in table grapes from India and be distributed in a viable state to a susceptible host is: **Low**.

4.18.2 Likelihood of establishment and spread

The likelihood of establishment and of spread for *P. castaneicola* and/or *P. diplodiella* within Western Australia has been adopted from the assessment of *P. castaneicola* and *P. diplodiella* for table grapes from Japan (Department of Agriculture 2014). The ratings from the previous assessment are:

| | |
|-----------------------------|-----------------|
| Likelihood of establishment | Moderate |
| Likelihood of spread | Moderate |

4.18.3 Overall likelihood of entry, establishment and spread

The overall likelihood of entry, establishment and spread is determined by combining the likelihoods of entry, of establishment and of spread using the matrix of rules shown in Table 2.2.

The overall likelihood that *P. castaneicola* and/or *P. diplodiella* will enter Western Australia as a result of trade in table grapes from India, be distributed in a viable state to a susceptible host, establish in Western Australia and subsequently spread within Western Australia is: **Low**.

4.18.4 Consequences

The consequences of *P. castaneicola* and *P. diplodiella* in Western Australia have been adopted from the assessment of *P. castaneicola* and *P. diplodiella* for table grapes from Japan (Department of Agriculture 2014), that is: **Low**.

4.18.5 Unrestricted risk estimate

Unrestricted risk is the result of combining the likelihoods of entry, establishment and spread with the outcome of overall consequences. Likelihoods and consequences are combined using the risk estimation matrix shown in Table 2.5.

Unrestricted risk estimate for *Pilidiella castaneicola* and *Pilidiella diplodiella*

| | |
|---|----------|
| Overall likelihood of entry, establishment and spread | Low |
| Consequences | Low |
| Unrestricted risk | Very low |

As indicated, the unrestricted risk estimate for *Pilidiella castaneicola* and *Pilidiella diplodiella* has been assessed as 'very low' which achieves Australia's ALOP. Therefore, no specific risk management measures are required for these pests.

4.19 Tobacco necrosis viruses

Tobacco necrosis virus A (TNV-A), Tobacco necrosis virus D (TNV-D) and Tobacco necrosis virus Nebraska isolate or related viruses (EP)

The taxonomy of 'tobacco necrosis virus' (TNV) has been revised. *Tobacco necrosis virus A* (TNV-A) and *Tobacco necrosis virus D* (TNV-D) have been recognised as distinct species in the Necrovirus genus (Coutts et al. 1991; Meulewaeter, Seurinck & Van Emmelo 1990), as have *Chenopodium necrosis virus* (ChNV) and *Olive mild mosaic virus* (OMMV), which were previously considered TNV isolates (Cardoso et al. 2005; Tomlinson et al. 1983). TNV isolates from Nebraska and Toyama (TNV-NE and TNV-Toyama) represent another species in the genus, as yet not officially recognised (Saeki et al. 2001; Zhang, French & Langenberg 1993) and molecular sequence data indicates some other necroviruses called 'tobacco necrosis viruses' are also distinct species (NCBI 2009).

The risk scenario of concern for tobacco necrosis viruses (TNVs) is that symptomless infected fruit might enter Australia and result in the establishment of one of them.

TNVs were assessed in many existing import policies, for example, for apples from China (Biosecurity Australia 2010) and table grapes from China (Biosecurity Australia 2011a). The assessment of TNVs presented here builds on these existing policies.

Differences in commodity, commercial production practices, climate conditions and the prevalence of pests between previous export areas and India make it necessary to reassess the likelihood that TNVs will be imported into Australia with table grapes from China.

TNVs have a wide range of hosts and the likelihood of distribution after arrival in Australia of TNVs with table grapes from India will be comparable to that for table grapes or other fruits from the previously assessed export area (Biosecurity Australia 2010, 2011a).

The likelihood of establishment and spread of TNVs in Australia will be comparable regardless of the fresh fruit commodity in which TNVs is imported into Australia, as these likelihoods relate specifically to events that occur in Australia and are independent of the importation pathway. The consequences of TNVs are also independent of the importation pathway. However, the department has reassessed the consequences of a TNV species outbreak in Australia in light of the taxonomic changes and additional information.

The department has reviewed the latest literature and no new information is available that would significantly change the risk ratings for distribution, establishment and spread as set out for *Tobacco necrosis viruses* in the existing policy. Therefore, those risk ratings will be adopted for this assessment.

The consequences of a TNV species outbreak in Australia have been reviewed in light of the taxonomic changes and additional information and analysis.

4.19.1 Likelihood of entry

The likelihood of entry is considered in two parts, the likelihood of importation and the likelihood of distribution, which consider pre-border and post-border issues, respectively.

Likelihood of importation

The likelihood that *Tobacco necrosis viruses* will arrive in Australia with the importation of table grapes from India is: **Low**.

Supporting information for this assessment is provided below:

- TNV was first reported in India in Madras and Tamil Nadu (Ramachandraiah, Venkatarathnam & Sulochana 1979). The virus isolate was able to cause typical TNV symptoms of leaf mottling, necrotic leaf lesions and ringspots in cluster bean, French bean and cowpea. The virus strain was tentatively designated as TNV-D (Ramachandraiah, Venkatarathnam & Sulochana 1979).
- Grapes are commercially grown in Madras and Tamil Nadu (DPP 2007), however, no information was found about the incidence or distribution of TNVs in grapevines in India.
- A strain of TNV was found naturally infecting several grapevine cultivars in South Africa (Cesati & Van Regenmortel 1969). The taxonomy, incidence and global distribution of the grapevine-infecting TNVs are not known.
- The strain of TNV found in grapevine in South Africa is graft-transmissible and spreads systemically in grapevine (Cesati & Van Regenmortel 1969). The virus is likely to be present in grape bunches.
- TNVs can infect a few species systemically (Kassanis 1970; Uyemoto 1981). Detectable systemic infection only occurs with certain combinations of host species and TNV species or strains (Brunt & Teakle 1996; Kassanis 1970; Uyemoto 1981). Some TNV species and strains may not infect grapevine and some may infect grapevines but not systemically and may not be in grape bunches.
- Some fruit species infected with TNV may not show adverse effects (Nemeth 1986). TNV usually causes necrotic lesions (Kassanis 1970), but no record was found indicating that infected grapevine showed symptoms.
- Grapes are usually pre-cooled to less than 4 degrees Celsius after packing and cold stored at 0–2 degrees Celsius at 90–95 per cent humidity after palletizing until shipment (DPP 2009). Conditions are maintained at 0–2 degrees Celsius and 90–95 per cent humidity throughout transport (DAFF 2010; DPP 2009, 2012). TNVs are likely to survive conditions during transport and storage.

The possible systemic infection in grapevine, moderated by the information that some TNV species and strains may not infect grapevine or infect grapevine but not systemically and the lack of reports of TNVs in grapevine in India, support a likelihood estimate for importation of 'low'.

Likelihood of distribution

The likelihood of distribution for TNVs has been adopted from the assessment of TNVs for apples and table grapes from China (Biosecurity Australia 2010, 2011a), that is: **Moderate**.

Overall likelihood of entry

The overall likelihood of entry is determined by combining the likelihood of importation with the likelihood of distribution using the matrix of rules shown in Table 2.2.

The likelihood that *Tobacco necrosis viruses* will enter Australia as a result of trade in table grapes from India and be distributed in a viable state to a susceptible host is: **Low**.

4.19.2 Likelihood of establishment and spread

The likelihood of establishment and of spread for TNVs has been adopted from the assessment of TNVs for apples from China (Biosecurity Australia 2010). The ratings from the previous assessments are:

| | |
|-----------------------------|-------------|
| Likelihood of establishment | High |
| Likelihood of spread | High |

4.19.3 Overall likelihood of entry, establishment and spread

The overall likelihood of entry, establishment and spread is determined by combining the likelihoods of entry, of establishment and of spread using the matrix of rules shown in Table 2.2.

The likelihood that TNVs will enter Australia as a result of trade in table grapes from India, be distributed in a viable state to a susceptible host, establish in Australia and subsequently spread within Australia is: **Low**.

4.19.4 Consequences

The consequences of the establishment of TNVs in Australia have been estimated according to the methods described in Table 2.3:

Based on the decision rules described in Table 2.4, that is, where the consequences of a pest with respect to one or more criteria are 'D', the overall consequences are estimated to be **Low**.

| Criterion | Estimate and rationale |
|----------------------------------|--|
| Direct | |
| Plant life or health | <p>D— significant at the district level</p> <p>Among the hosts in which TNVs cause disease, carrot, potato and strawberry are the most economically important in Australia. In 2009/10, the estimated value of the carrot, potato and strawberry crops is \$176m, \$614m and \$308m respectively (HAL 2012).</p> <p>TNVs cause sporadic diseases in vegetable and ornamental crops in some years (Kassanis 1970; Nemeth 1986; Smith et al. 1988; Uyemoto 1981; Zitikaite & Staniulis 2009). No reports of adverse effects on fruit trees have been found (Nemeth 1986). A disease in trembling aspen (<i>Populus tremuloides</i>) may be caused by TNVs (Hibben, Bozarth & Reese 1979).</p> <p>TNVs cause rusty root disease of carrot, Augusta disease of tulip, stipple streak disease of common bean, and necrosis diseases of cabbage, cucumber, soybean and zucchini and ABC disease of potato (Smith et al. 1988; Uyemoto 1981; Xi et al. 2008; Zitikaite & Staniulis 2009).</p> <p>Losses between 20 per cent and 50 per cent have been reported in glasshouse grown cucumbers and in tulips (CABI 2014). Lower losses probably occur more frequently than such high losses. No estimates of losses in carrot, potato and strawberry have been found but it is possible that substantial losses occur sometimes. Symptomless viral infections of plants, in general, may cause no yield loss, but they may cause yield losses as high as 15 per cent (Bos 1999; Gibbs & Harrison 1976).</p> <p>Naturally infected vegetable crops show a range of symptoms, including spots, flecks, streaks, necrosis and stunting. In strawberry in the Czech Republic, TNV has caused dwarfing and leaf and root necrosis (Martin & Tzanetakis 2006).</p> <p>Stipple streak disease has been reported in Qld causing small yield losses (Teakle 1988), but no reports of TNVs causing other diseases in Australia have been found, suggesting that the combinations of virus strain, vector biotype and host plant cultivar that result in disease have not occurred in Australia.</p> <p>Strains have been distinguished by various characteristics, including the symptoms they cause, their host ranges and genetic sequences (Kassanis 1970). The diseases recorded in common bean and cucumber are probably caused by distinct TNV strains (Brunt & Teakle 1996; Zitikaite & Staniulis 2009). The TNV strains detected in apple caused lesions in tests with cowpea (<i>Vigna sinensis</i>) and <i>Chenopodium quinoa</i> (Uyemoto & Gilmer 1972), but no report of further investigation of their disease causing potential was found.</p> <p>A satellite virus replicates with some strains of TNV (Kassanis 1970; Uyemoto 1981) but no report has been found indicating greater disease when the satellite virus is present.</p> <p>Because the wide host range of TNVs and their chytrid vectors it is possible that some native plants will be susceptible, although no supporting evidence was found.</p> |
| Other aspects of the environment | <p>A— Indiscernible at the local level</p> <p>There are no known direct consequences of these species on other aspects of the environment.</p> |
| Indirect | |
| Eradication, control | <p>D— significant at the district level</p> <p>Virus control measures in fields are limited and eradication may not be possible unless an outbreak is detected at an early stage. If detected at an early stage, an outbreak might be controlled or eradicated by removing host plants and deep burying or incinerating potentially infected plant material, then leaving the fields fallow and controlling weed hosts. Further action might be required including cropping with non-host species and altering and lengthening crop rotations. Resistant cultivars may be planted, if they are available (CABI 2014). Establishment and spread in a glasshouse may be controlled by reducing or eliminating <i>Oplidium</i> infestation of soil by chemical treatment or by heating by composting or soil pasteurisation (Asjes & Blom-Barnhoorn 2002; CABI 2014). This may add significantly to costs. TNVs tolerate temperatures as high as 95 degrees Celsius (Brunt & Teakle 1996), so the temperatures achieved by composting and pasteurisation may not eliminate the viruses. Propagation of virus free plants and careful sanitation may reduce the chance of outbreaks (CABI 2014; Smith et al. 1988).</p> |

| Criterion | Estimate and rationale |
|----------------------------------|---|
| Domestic trade | C— minor significance at the district level Australian states are unlikely to set up restrictions on interstate trade if a foreign TNV becomes established unless it causes significant disease, which is unlikely. |
| International trade | C— minor significance at the district level If a damaging foreign TNV became established in Australia, additional restrictions might be introduced on the international trade of some vegetables or ornamentals that might lead to the loss of markets and some industry adjustment. |
| Environmental and non-commercial | A— Indiscernible at the local level There are no known indirect consequences of these species on the environment. |

4.19.5 Unrestricted risk estimate

Unrestricted risk is the result of combining the likelihoods of entry, establishment and spread with the outcome of overall consequences. Likelihoods and consequences are combined using the risk estimation matrix shown in Table 2.5.

| Unrestricted risk estimate for <i>Tobacco necrosis viruses</i> | |
|--|----------|
| Overall likelihood of entry, establishment and spread | Low |
| Consequences | Low |
| Unrestricted risk | Very low |

As indicated, the unrestricted risk estimate for *Tobacco necrosis viruses* has been assessed as 'very low', which achieves Australia's ALOP. Therefore, no specific risk management measures are required for this pest.

Assessments for quarantine pests for which the likelihood and consequence ratings have been determined in a previous assessment

4.20 Spotted wing drosophila

Drosophila suzukii (EP)

The quarantine risks posed by *Drosophila suzukii* from all countries and for all commodities, including table grapes, were previously assessed in the Final pest risk analysis (PRA) report for *Drosophila suzukii* (Department of Agriculture 2013b). Therefore, there is no need to reassess this pest here. A summary of pest information and previous assessments from the final PRA report for *D. suzukii* is provided here.

In India, *Drosophila suzukii indicas* has been found in Kashmir (Hauser, Gaimari & Damus 2009), northern India (Toda 1991) and Uttar Pradesh (Chamoli & Pauri region) at approximately 5000 feet (1524 metres) (Singh & Negi 1989) or at 6000 feet (1800 metres) above sea level (Singh & Bhatt 1988). *Drosophila suzukii* has also been recorded from Mysore in southern India at altitude (680 metres and above) where it is collected infrequently (Guruprasad, Hegde & Krishna 2010).

Drosophila suzukii preferentially oviposit on ripe fruit but will also oviposit on unripe and overripe fruit (Brewer et al. 2012; Kanzawa 1939; Lee et al. 2011). Larvae feeding on very acidic fruit fail to complete development (Kanzawa 1935). In its native and introduced range, *D. suzukii* has been recorded to cause damage to a range of fruits including grapes, cherry, blueberry and red bayberry, mulberries, peaches, plums, strawberries and various caneberries.

On grapes, oviposition trials on wine and table grapes have shown that fully-ripe table grapes can be attacked (Atallah et al. 2014; Maiguashca et al. 2010; Saguez, Lasnier & Vincent 2013). Damaged fruit with low sugar levels will be oviposited in but larvae develop poorly and fail to pupate (Maiguashca et al. 2010). Kanzawa (1939) recorded that different grape varieties sustained different levels of attack and considered skin thickness was the factor that limited oviposition. Oviposition of *D. suzukii* has been reported on a number of grape varieties/cultivars which are 100 per cent *V. vinifera*, such as Gros Coleman, Muscat of Alexandria, Muscat of Hamburg, Foster's seeding Rose de Italy, Kyoshin (Kanzawa 1939), Thompson Seedless (Lee et al. 2011), Black Manuka and Perlette (WSUE 2010). Reports of oviposition on grape varieties/cultivars which are 100 per cent *Vitis labrusca* have not been found. There have been reports of a number of grape varieties/cultivars not being attacked by *D. suzukii*, some of these are 100 per cent *Vitis vinifera* (for example Koshu, Chasselas de Fontainbleau, Golden champion and White Malaga), some are 100 per cent *Vitis labrusca* (for example Concord, Eaton, Niagara and Hostess seedling) (Kanzawa 1939), and some are hybrids between *V. vinifera* and *V. labrusca* for which percentage of *V. vinifera* as parentage range from 25 per cent (for example Early Campbell) (Maiguashca et al. 2010) to 75 per cent (for example Brighton) (Kanzawa 1939).

When *D. suzukii* is given a choice between several host fruits (for example raspberry, cherry, strawberry, grape), grape ('Thompson Seedless') were the least preferred host on undamaged fruit (Atallah et al. 2014; Lee et al. 2011).

During the 1930s in Japan, *D. suzukii* was trapped in vineyards at high levels and there are reports of damage as high as 80 per cent (Kanzawa 1939). More recently there have been reports of outbreaks of *D. suzukii* on grapes in Hokkaido (CFIA 2010).

The risk scenario of concern for *D. suzukii* is the presence of the larvae in mature bunches of grapes.

4.20.1 Overall likelihood of entry, establishment and spread

Based on the Final pest risk analysis (PRA) report for *Drosophila suzukii* (Department of Agriculture 2013b) the overall likelihood that *D. suzukii* will enter Australia as a result of trade in table grapes (*Vitis vinifera*) from India, be distributed in a viable state to a susceptible host, establish in Australia and subsequently spread within Australia is: **Moderate**.

The final PRA for *D. suzukii* (Department of Agriculture 2013b) recognises that the importation risk of *D. suzukii* on table grape pathway could be different for particular varieties and/or cultivars. The importation risk and hence the overall likelihood of entry, establishment and spread are likely to be lower for commercial quality grapes of varieties and/or cultivars of *V. vinifera* or hybrids demonstrated to be poor hosts for oviposition by *D. suzukii*.

4.20.2 Consequences

Based on the Final pest risk analysis (PRA) report for *Drosophila suzukii* (Department of Agriculture 2013b) the potential consequences of the establishment of *D. suzukii* in Australia are: **High**.

4.20.3 Unrestricted risk estimate

Based on the Final pest risk analysis (PRA) report for *Drosophila suzukii* (Department of Agriculture 2013b) the unrestricted risk estimate for *D. suzukii* has been assessed as 'high', which exceeds Australia's ALOP. Therefore, specific risk management measures are required for this pest.

4.21 Pest risk assessment conclusions

Key to Tables 4.5 to 4.8 (starting next page)

Genus species (EP): pests for which policy already exists. The outcomes of previous assessments and/or reassessments in this IRA are presented in Tables 4.5 to 4.8

Genus species (Acronym for state/territory): state/territory in which regional quarantine pests have been identified

Likelihoods for entry, establishment and spread

| | |
|-----|---|
| N | negligible |
| EL | extremely low |
| VL | very low |
| L | low |
| M | moderate |
| H | high |
| EES | overall likelihood of entry, establishment and spread |

Assessment of consequences from pest entry, establishment and spread

| | |
|-----|--|
| PLH | plant life or health |
| OE | other aspects of the environment |
| EC | eradication, control |
| DT | domestic trade |
| IT | international trade |
| ENC | environmental and non-commercial |
| A-G | consequence impact scores are detailed in section 2.2.3 |
| A | Indiscernible at the local level |
| B | Minor significance at the local level |
| C | Significant at the local level |
| D | Significant at the district level |
| E | Significant at the regional level |
| F | Significant at the national level |
| G | Major significance at the national level |
| URE | unrestricted risk estimate. This is expressed on an ascending scale from negligible to extreme |

Table 4.5 Summary of unrestricted risk estimates for quarantine pests associated with table grapes from India for which a full pest risk assessment is conducted

| Pest name | Likelihood of | | | | | | Consequences | | | | | | URE | |
|--|---------------|--------------|---------|---------------|--------|-----|--------------|----|----|----------|----|-----|-----|---------|
| | Entry | | | Establishment | Spread | EES | Direct | | | Indirect | | | | Overall |
| | Importation | Distribution | Overall | | | | PLH | OE | EC | DT | IT | ENC | | |
| Bacteria | | | | | | | | | | | | | | |
| <i>Xanthomonas campestris</i> <i>pv. viticola</i> | H | VL | VL | M | M | VL | D | A | D | D | D | B | L | N |
| Viruses | | | | | | | | | | | | | | |
| <i>Tomato black ring virus</i> | M | VL | VL | VL | M | EL | E | A | E | D | D | B | M | N |

Table 4.6 Summary of unrestricted risk estimates for quarantine pests associated with table grapes from India for which the URE outcome is adopted from previous assessments

| Pest name | Likelihood of | | | | | | Consequences | | | | | | URE | |
|--|---|--------------|---------|---------------|--------|-----|--------------|----|----|----------|----|-----|-----|---------|
| | Entry | | | Establishment | Spread | EES | Direct | | | Indirect | | | | Overall |
| | Importation | Distribution | Overall | | | | PLH | OE | EC | DT | IT | ENC | | |
| Phylloxera [Hemiptera: Phylloxeridae] | | | | | | | | | | | | | | |
| <i>Daktulosphaira vitifoliae</i> (EP) | The URE outcome of exceeding Australia's ALOP from existing policy has been adopted | | | | | | | | | | | | | |
| Soft scale [Hemiptera: Coccidae] | | | | | | | | | | | | | | |
| <i>Parthenolecanium corni</i> (EP, WA) | The URE outcome of achieving Australia's ALOP from existing policy has been adopted | | | | | | | | | | | | | |

| Pest name | Likelihood of | | | | | Consequences | | | | | | URE | | |
|--|---|--------------|---------|---------------|--------|--------------|--------|----|----|----------|----|-----|---------|-----|
| | Entry | | | Establishment | Spread | EES | Direct | | | Indirect | | | Overall | |
| | Importation | Distribution | Overall | | | | PLH | OE | EC | DT | IT | | | ENC |
| Mealybugs [Hemiptera: Pseudococcidae] | | | | | | | | | | | | | | |
| <i>Planococcus ficus</i> (EP) | The URE outcome of exceeding Australia's ALOP from existing policy has been adopted | | | | | | | | | | | | | |
| <i>Planococcus lilacinus</i> (EP) | | | | | | | | | | | | | | |
| <i>Planococcus minor</i> (EP) | | | | | | | | | | | | | | |
| <i>Rastrococcus iceryoides</i> (EP) | | | | | | | | | | | | | | |
| Plume moth [Lepidoptera: Pterophoridae] | | | | | | | | | | | | | | |
| <i>Platyptilia ignifera</i> (EP) | The URE outcome of achieving Australia's ALOP from existing policy has been adopted | | | | | | | | | | | | | |
| Thrips [Thysanoptera: Thripidae] | | | | | | | | | | | | | | |
| <i>Retithrips syriacus</i> (EP) | The URE outcome of exceeding Australia's ALOP from existing policy has been adopted | | | | | | | | | | | | | |
| <i>Rhipiphorothrips cruentatus</i> (EP) | | | | | | | | | | | | | | |
| Spider mite [Trombidiformes: Tetranychidae] | | | | | | | | | | | | | | |
| <i>Tetranychus kanzawai</i> (EP, WA) | The URE outcome of exceeding Australia's ALOP from existing policy has been adopted | | | | | | | | | | | | | |
| Fungi | | | | | | | | | | | | | | |
| <i>Guignardia bidwellii</i> (EP) | The URE outcome of exceeding Australia's ALOP from existing policy has been adopted | | | | | | | | | | | | | |
| <i>Monolinia fructigena</i> (EP) | The URE outcome of exceeding Australia's ALOP from existing policy has been adopted | | | | | | | | | | | | | |
| <i>Phakopsora euvtitis</i> (EP) | The URE outcome of exceeding Australia's ALOP from existing policy has been adopted | | | | | | | | | | | | | |
| <i>Phomopsis viticola</i> (EP, WA) | The URE outcome of achieving Australia's ALOP from existing policy has been adopted | | | | | | | | | | | | | |

Table 4.7 Summary of unrestricted risk estimates for quarantine pests associated with table grapes from India for which some of the likelihood and/or consequence ratings are adopted from previous assessments

| Pest name | Likelihood of | | | | | | Consequences | | | | | | URE | |
|--|---------------|--------------|---------|---------------|--------|-----|--------------|----|----|----------|----|-----|-----|---------|
| | Entry | | | Establishment | Spread | EES | Direct | | | Indirect | | | | Overall |
| | Importation | Distribution | Overall | | | | PLH | OE | EC | DT | IT | ENC | | |
| | | | | | | | | | | | | | | |
| Fruit fly [Diptera: Tephritidae] | | | | | | | | | | | | | | |
| <i>Bactrocera correcta</i> (EP) | VL | M | VL | H | H | VL | | | | | | H | L | |
| <i>Bactrocera dorsalis</i> (EP) | | | | | | | | | | | | | | |
| Tortricid moth [Lepidoptera: Tortricidae] | | | | | | | | | | | | | | |
| <i>Archips machlopiis</i> | L | M | L | H | H | L | | | | | | M | L | |
| Pyralid moth [Lepidoptera: Pyralidae] | | | | | | | | | | | | | | |
| <i>Cryptoblabes gnidiella</i> (EP) | VL | L | VL | H | H | VL | | | | | | M | VL | |
| Fungi | | | | | | | | | | | | | | |
| <i>Greeneria uvicola</i> (EP, WA) | H | L | L | L | L | VL | | | | | | L | N | |
| <i>Pestalotiopsis menezesiana</i> (EP, WA) | M | L | L | H | H | L | | | | | | L | VL | |
| <i>Pestalotiopsis uvicola</i> (EP, WA) | | | | | | | | | | | | | | |
| <i>Pilidiella castaneicola</i> (EP, WA) | M | M | L | M | M | L | | | | | | L | VL | |
| <i>Pilidiella diplodiella</i> (EP, WA) | | | | | | | | | | | | | | |
| Viruses | | | | | | | | | | | | | | |
| <i>Tobacco necrosis viruses</i> (EP) | L | M | L | H | H | L | D | A | D | C | C | A | L | VL |

Table 4.8 Summary of unrestricted risk estimates for quarantine pests associated with table grapes from India for which the likelihood and consequence ratings have been determined in a previous assessment

| Pest name | Likelihood of | | | | | Consequences | | | | | | URE | | |
|---|---------------|--------------|---------|---------------|--------|--------------|--------|----|----|----------|----|-----|---------|-----|
| | Entry | | | Establishment | Spread | EES | Direct | | | Indirect | | | Overall | |
| | Importation | Distribution | Overall | | | | PLH | OE | EC | DT | IT | | | ENC |
| | | | | | | | | | | | | | | |
| Spotted wing drosophila [Diptera: Drosophilidae] | | | | | | | | | | | | | | |
| <i>Drosophila suzukii</i> (EP) | | | | | | | | | | | | | | |
| | | | | | M | | | | | | | H | H | |

5 Pest risk management

This chapter provides information on the management of quarantine pests identified with an unrestricted risk exceeding Australia's appropriate level of protection (ALOP). The recommended phytosanitary measures are described in this chapter.

5.1 Pest risk management measures and phytosanitary procedures

Pest risk management evaluates and selects options for measures to reduce the risk of entry, establishment or spread of quarantine pests for Australia where they have been assessed to have an unrestricted risk above Australia's ALOP. In calculating the unrestricted risk, existing commercial production practices in India have been considered, as have post-harvest procedures and the packing of fruit.

In addition to India's existing commercial production practices for table grapes and minimum border procedures in Australia, specific pest risk management measures, including operational systems, are recommended to achieve Australia's ALOP.

In this chapter, the Australian Government Department of Agriculture and Water Resources has identified risk management measures that may be applied to consignments of table grapes sourced from India. Finalisation of the quarantine conditions may be undertaken with input from the Australian states and territories as appropriate.

5.1.1 Pest risk management for quarantine pests

The pest risk analysis identified the quarantine pests listed in Table 5.1 as having an unrestricted risk above Australia's ALOP. Therefore, phytosanitary measures are required to manage the risks posed by these pests.

Table 5.1 Phytosanitary measures recommended for quarantine pests for table grapes from India

| Pest | Common name | Measures |
|---|--------------------------|---|
| Arthropods | | |
| <i>Planococcus ficus</i> (EP) | Grapevine mealybug | Visual inspection and, if detected, remedial action a (for example methyl bromide fumigation) |
| <i>Planococcus lilacinus</i> (EP) | Coffee mealybug | |
| <i>Planococcus minor</i> (EP) | Pacific mealybug | |
| <i>Rastrococcus iceryoides</i> (EP) | Downey snowline mealybug | |
| <i>Tetranychus kanzawai</i> (EP, WA) | Kanzawa spider mite | |
| <i>Archips machlopiis</i> | Leaf rolling moth | |
| <i>Retithrips syriacus</i> (EP) | Black vine thrips | |
| <i>Rhipiphorothrips cruentatus</i> (EP) | Grapevine thrips | |
| <i>Bactrocera correcta</i> (EP) | Guava fruit fly | Area freedom b |
| <i>Bactrocera dorsalis</i> (EP) | Oriental fruit fly | OR Fruit treatment known to be effective against all life stages of <i>Bactrocera correcta</i> and <i>Bactrocera dorsalis</i> (for example irradiation OR cold disinfestation treatment) |
| <i>Drosophila suzukii</i> (EP) | Spotted wing drosophila | Area freedom b OR Systems approach OR Fruit treatment known to be effective against all life stages of <i>Drosophila suzukii</i> (for example irradiation OR SO ₂ /CO ₂ fumigation followed by cold treatment) |
| <i>Daktulosphaira vitifoliae</i> (EP) | Grapevine phylloxera | Area freedom b OR Fruit treatment known to be effective against all life stages of <i>Daktulosphaira vitifoliae</i> (for example sulphur pad treatment) |
| Pathogens | | |
| <i>Guignardia bidwellii</i> (EP) | Black rot | Area freedom b |
| <i>Monilinia fructigena</i> (EP) | Brown rot | OR |
| <i>Phakopsora euvitis</i> (EP) | Grapevine leaf rust | Systems approach |

a Remedial action by DAC may include: withdrawing the consignment from export to Australia or applying approved treatment of the consignment to ensure that the pest is no longer viable.

b Area freedom may include pest free areas, pest free places of production or pest free production sites.

(EP) Species has been assessed previously and import policy already exists.

(WA) Pests of regional concern for Western Australia only.

Risk management measures recommended here build on the existing policies for the import of table grapes from California (AQIS 1999, 2000; Biosecurity Australia 2006a; DAFF 2013), Chile (Biosecurity Australia 2005b), China (Biosecurity Australia 2011a), Korea (Biosecurity Australia 2011b), and Japan (Department of Agriculture 2014), which include most of the pests identified in Table 5.1. Among these existing policies, considerable trade in table grapes from California has taken place since 2002, demonstrating that the measures implemented are effective.

The risk management measure options recommended in this final report are consistent with the existing policies and include:

- visual inspection and, if detected, remedial action for spider mites, mealybugs, moths and thrips
- area freedom or fruit treatment known to be effective against all life stages of fruit flies
- area freedom, systems approach approved by the Australian Government Department of Agriculture and Water Resources or fruit treatment known to be effective against all life stages of spotted wing drosophila
- area freedom or fruit treatment known to be effective against all life stages of grapevine phylloxera
- area freedom or a systems approach approved by the Australian Government Department of Agriculture and Water Resources for black rot, brown rot and grapevine leaf rust.

Management for *Tetranychus kanzawai*, *Planococcus ficus*, *Planococcus lilacinus*, *Planococcus minor*, *Rastrococcus iceryoides*, *Archips machlopiis*, *Retithrips syriacus* and *Rhipiphorothrips cruentatus*

To manage the risks from *Tetranychus kanzawai* (Kanzawa spider mite); *Planococcus ficus* (grapevine mealybug), *Planococcus lilacinus* (coffee mealybug), *Planococcus minor* (Pacific mealybug) and *Rastrococcus iceryoides* (downey snowline mealybug); *Archips machlopiis* (leaf rolling moth); *Rhipiphorothrips cruentatus* (grapevine thrips) and *Retithrips syriacus* (black vine thrips), the department recommends visual inspection and, if detected, remedial action as a measure for these pests. The objective of the recommended visual inspection is to ensure that any consignments of table grapes from India infested with these pests are identified and subjected to appropriate remedial action. The appropriate remedial action will reduce the risk associated with these pests to at least 'very low', which would achieve Australia's ALOP.

Pre-export visual inspection and, if detected, remedial action by DAC

All table grape consignments for export to Australia must be inspected by the Department of Agriculture and Cooperation of India (DAC) and found free of these quarantine arthropod pests. Export lots or consignments found to contain any of these pests must be subject to remedial action. Remedial action may include withdrawing the lots or consignments from export to Australia or, if available, applying approved treatment to the export lots or consignments to ensure that the pest is no longer viable.

Management for *Drosophila suzukii*

Options recommended to manage this pest in the Final pest risk analysis report for *Drosophila suzukii* (Department of Agriculture 2013b) are area freedom, irradiation, systems approach, or fruit treatment known to be effective against all life stages of *D. suzukii*.

Area freedom

The requirements for establishing pest free areas or pest free places of production are set out in ISPM 4: *Requirements for the establishment of pest free areas* (FAO 1995) and ISPM 10: *Requirements for the establishment of pest free places of production and pest free production sites* (FAO 1999).

Should India wish to use area freedom as a measure to manage the risk posed by *D. suzukii*, DAC would need to provide Australia with a submission demonstrating area freedom for consideration by the Australian Government Department of Agriculture and Water Resources.

If area freedom from *D. suzukii* could be demonstrated for any areas in India, the likelihood of importation of this pest with table grapes sourced from those areas would be reduced to at least 'extremely low'. The restricted risk would then be reduced to at least 'very low', which would achieve Australia's ALOP.

Systems approach

A systems approach that uses the integration of different risk management measures, at least two of which act independently, and which cumulatively achieve the required level of phytosanitary protection could be used to reduce the risk of *D. suzukii* being imported into Australia with consignments of table grapes. More information on a systems approach is set out in ISPM 14: *The use of integrated measures in a systems approach for pest risk management* (FAO 2002).

A systems approach to address the risks posed by *D. suzukii* on table grapes could be based on a combination of fruit protection for example fruit bagging, vineyard preventative measures and monitoring, and pest control with post-harvest measures. The approach could be used to progressively reduce the risk of infested fruit being imported into Australia with consignments of table grapes.

Should India wish to use a systems approach as a measure to manage the risk posed by *D. suzukii*, DAC would need to submit to Australia a proposal outlining components of the system and how these components will address the risks posed by this pest. The Australian Government Department of Agriculture and Water Resources will consider the effectiveness of any system proposed by DAC.

Treatment of fruit

A treatment that is known to be effective against all life stages of *D. suzukii* is a measure that might be applied to manage the risk posed by this pest in imports of host fruits.

Treatment of fruit, with suitable efficacy, would reduce the likelihood of importation of infested fruit to at least 'extremely low'. The restricted risk would then be reduced to at least 'very low',

which would achieve Australia's ALOP. Treatments of fruit will need to be applied offshore to ensure that any live adult flies in consignments of fruit do not enter Australia.

Treatment options that might be applied to manage the risk posed by *D. suzukii* in imports of table grapes include:

Irradiation

Irradiation treatment is considered a suitable measure option for *D. suzukii* (Follett, Swedman & Price 2014). The Australian Government Department of Agriculture and Water Resources recommends a treatment schedule of 150 gray minimum absorbed dose, consistent with ISPM 28 Annex 7: *Irradiation treatment for fruit flies of the family Tephritidae (generic)* (FAO 2009). Although lower doses (78 gray) have been shown to induce sterility of all immature life stages associated with fruit, adults can successfully emerge from irradiated pupae. The detection of a sterilised *D. suzukii* adult post border would result in regulatory actions. A dose of 150 gray would make adult emergence from irradiated fruit an unlikely event.

Methyl bromide fumigation

Preliminary methyl bromide fumigation trials have shown 100 per cent mortality on all life stages. Methyl bromide fumigation of exported fruit might be used as a stand-alone treatment to achieve Australia's ALOP. However, before methyl bromide could be recommended as a permanent phytosanitary measure for *D. suzukii* in table grapes, a complete efficacy treatment proposal by a proponent country would need to be reviewed and accepted by the Australian Government Department of Agriculture and Water Resources.

Combined sulphur dioxide (SO₂)/carbon dioxide (CO₂) fumigation followed by cold disinfestation treatment

The Australian Government Department of Agriculture and Water Resources reviewed the efficacy data in support of a combination treatment of SO₂/CO₂ fumigation followed by a cold disinfestation treatment (listed below), and considered it suitable to manage the risk of *D. suzukii* in table grapes (*Vitis vinifera*).

- 6 per cent carbon dioxide (CO₂) and 1 per cent sulphur dioxide (SO₂) by volume for 30 minutes, at a pulp temperature of 15.6 degrees Celsius or greater, followed by;
- A cold treatment for 6 days or more at a pulp temperature of -0.50 degrees Celsius plus or minus 0.50 degrees Celsius.

OR

- 6 per cent carbon dioxide (CO₂) and 1 per cent sulphur dioxide (SO₂) by volume for 30 minutes, at a pulp temperature of 15.6 degrees Celsius or greater, followed by;
- A cold treatment for twelve days or more at a pulp temperature of 0.9 degrees Celsius plus or minus 0.50 degrees Celsius.

Additional post-treatment security measures may be required to limit post-treatment contamination by flies that are attracted to ripe fruit.

Alternative treatments

Other treatments, demonstrated to be effective against all life stages of *D. suzukii* for table grapes, will be considered by the Australian Government Department of Agriculture and Water Resources if proposed by DAC.

Management for *Bactrocera correcta* and *Bactrocera dorsalis*

The Australian Government Department of Agriculture and Water Resources recommends the options of area freedom, irradiation, or cold disinfestation treatment as measures to reduce the risks associated with *Bactrocera correcta* and *Bactrocera dorsalis*.

Area freedom

Area freedom is a measure that might be applied to manage the risk posed by guava fruit fly and Oriental fruit fly. The requirements for establishing pest free areas or pest free places of production are set out in ISPM 4: Requirements for the *establishment of pest free areas* (FAO 1995) and ISPM 10: *Requirements for the establishment of pest free places of production and pest free production sites* (FAO 1999) and more specifically in ISPM 26: *Establishment of pest free areas for fruit flies (Tephritidae)* (FAO 2015b).

Should India wish to use area freedom as a measure to manage the risk posed by *B. correcta* and *B. dorsalis*, DAC would need to provide Australia with a submission demonstrating area freedom for consideration by the Australian Government Department of Agriculture and Water Resources.

If area freedom from *Bactrocera correcta* and/or *Bactrocera dorsalis* can be demonstrated for any areas in India, the likelihood of importation of the pest species with table grapes sourced from these areas would be reduced to at least 'extremely low'. The restricted risk would then be reduced to at least 'very low', which would achieve Australia's ALOP.

Monitoring and trapping of fruit flies in the specific table grape export vineyards and packing houses of India would be required.

Under the area freedom option, the Department of Agriculture and Cooperation, India would be required to notify the Australian Government Department of Agriculture and Water Resources of a detection of any fruit fly species (Tephritidae) of economic importance in the regions within 48 hours. The Australian Government Department of Agriculture and Water Resources would then assess the species and number of individual flies detected and the circumstances of the detection, before advising DAC of the action to be taken. If fruit flies are detected at offshore pre-shipment inspection or on-arrival inspection, trade would be suspended immediately, pending the outcome of an investigation.

For table grapes sourced from outside the recognised fruit fly pest free areas, fruit treatment known to be effective against all life stages of *B. correcta* and *B. dorsalis*, for example cold disinfestation treatment or irradiation must be undertaken.

Cold disinfestation treatments

Cold disinfestation treatments is considered a suitable measure option for *B. dorsalis* and *B. correcta*. Cold treatments can be conducted pre-export in India or in-transit.

Treatment regimes consistent with the *USDA Treatment Manual* (USDA 2015) for *B. dorsalis* on a range of commodities are recommended for the disinfestation of *B. dorsalis* and *B. correcta* on table grapes. Cold treatments which are effective for *B. dorsalis* are considered to be effective for *B. correcta* as *B. correcta* is more sensitive to cold than *B. dorsalis* (Liu & Ye 2009).

It is recommended that where cold disinfestation treatment is utilised, it must be carried out according to the specifications below:

- 0.99 degrees Celsius or below for 15 days, or
- 1.38 degrees Celsius or below for 18 days

Irradiation

Irradiation treatment is considered a suitable measure option for *B. correcta* and *B. dorsalis*. The treatment schedule of 150 gray minimum absorbed dose is specified in ISPM 28 Annex 7: *Irradiation treatment for fruit flies of the family Tephritidae (generic)* (FAO 2009).

Alternative measures

Measures for *B. dorsalis* and *B. correcta* could also include other equivalent measures, subject to the provision and acceptance of suitable efficacy data.

Management for *Daktulosphaira vitifoliae*

The Australian Government Department of Agriculture and Water Resources proposes area freedom or fruit treatment known to be effective against all life stages of *D. vitifoliae* such as sulphur pad treatment.

Area freedom

Area freedom is a measure that might be applied to manage the risk posed by *D. vitifoliae*. The requirements for establishing pest free areas or pest free places of production are set out in ISPM 4: *Requirements for the establishment of pest free areas* (FAO 1995) and ISPM 10: *Requirements for the establishment of pest free places of production and pest free production sites* (FAO 1999).

Should India wish to use area freedom as a measure to manage the risk posed by *D. vitifoliae*, DAC would need to provide Australia with a submission demonstrating area freedom for consideration by the Australian Government Department of Agriculture and Water Resources.

If area freedom from *D. vitifoliae* could be demonstrated for any areas in India, the likelihood of importation of this pest with table grapes sourced from those areas would be reduced to at least 'extremely low'. The restricted risk would then be reduced to 'negligible', which would achieve Australia's ALOP.

Treatment of fruit

Treatment that is known to be effective against all life stages of *D. vitifoliae* is a measure that might be applied to manage the risk posed by *D. vitifoliae* with table grapes sourced from areas infested or affected by this pest.

Treatment options that might be applied to manage the risk posed by *D. vitifoliae* in imports of table grapes include:

Sulphur pads

Commercial sulphur pads with proven efficacy against *D. vitifoliae* packed inside the plastic liner in all cartons of table grapes for export could be used to manage the risk posed by this pest. The sulphur pads must be a registered product containing a minimum of 970 grams per kilogram anhydrous sodium metabisulphite used at the rate specified on the label (DPIPWE Tasmania 2015; DPIF 2013; Queensland Government 2014).

The inclusion of sulphur pads in all cartons of table grapes for export is to reduce the survival of *D. vitifoliae* associated with packed table grapes and the likelihood of introduction to at least 'very low'. The restricted risk would then be reduced to at least 'very low', which would achieve Australia's ALOP.

Combined SO₂/CO₂ fumigation

The Australian Government Department of Agriculture and Water Resources reviewed the efficacy data in support of a combination treatment of SO₂/CO₂ fumigation (listed below) and considered it suitable to manage the risk of *D. vitifoliae*

- 6 per cent carbon dioxide (CO₂) and 1 per cent sulphur dioxide (SO₂) by volume for 30 minutes, at a pulp temperature of 15.6 degrees Celsius or greater.

Additional post-treatment security measures may be required to limit post-treatment contamination by this pest.

Treatment of table grapes with combined SO₂/CO₂ fumigation would reduce the likelihood of introduction of infested fruit to at least 'extremely low'. The restricted risk would then be reduced to at least 'negligible', which would achieve Australia's ALOP.

Management for *Guignardia bidwellii*, *Monilinia fructigena* and *Phakopsora euvitis*

The Australian Government Department of Agriculture and Water Resources recommends area freedom or a systems approach as measures for *Guignardia bidwellii*, *Monilinia fructigena* and *Phakopsora euvitis*.

Area freedom

Area freedom is a measure that might be applied to manage the risk posed by these pathogens. The requirements for establishing pest free areas or pest free places of production are set out in ISPM 4: *Requirements for the establishment of pest free areas* (FAO 1995) and ISPM 10: *Requirements for the establishment of pest free places of production and pest free production sites* (FAO 1999).

Should India wish to use area freedom as a measure to manage the risk posed by these pathogens, DAC would need to provide Australia with a submission demonstrating area freedom for consideration by the Australian Government Department of Agriculture and Water Resources.

If area freedom from these pathogens could be demonstrated for any areas in India, the likelihood of importation of these pathogens with table grapes sourced from those areas would be reduced to at least 'extremely low'. The restricted risk would then be reduced to at least 'very low', which would achieve Australia's ALOP.

Systems approach

A systems approach that uses the integration of different risk management measures, at least two of which act independently, and which cumulatively achieve the required level of phytosanitary protection could be used to reduce the risk of these pathogens being imported to Australia with consignments of table grapes. More information on a systems approach is set out in ISPM 14: *The use of integrated measures in a systems approach for pest risk management* (FAO 2002).

The Australian Government Department of Agriculture and Water Resources considers a systems approach could be based on area of low pest prevalence, a combination of fruit protection for example fruit bagging, vineyard preventative measures and monitoring, and pest control with post-harvest measures. The approach could be used to progressively reduce the risk of infested table grapes being imported to Australia.

Should India wish to use a systems approach as a measure to manage the risk posed by these pathogens, DAC would need to submit a proposal outlining components of the system and how these components will address the risks posed by these pathogens. The Australian Government Department of Agriculture and Water Resources will consider the effectiveness of any system proposed by DAC.

5.1.2 Consideration of alternative measures

Consistent with the principle of equivalence detailed in ISPM 11: *Pest risk analysis for quarantine pests* (FAO 2013), the Australian Government Department of Agriculture and Water Resources will consider any alternative measure proposed by DAC, providing that it achieves Australia's ALOP. Evaluation of such measures or treatments will require a technical submission from DAC that details the proposed treatment and including data from suitable treatment trials to demonstrate efficacy.

5.2 Operational system for the maintenance and verification of phytosanitary status

A system of operational procedures is necessary to maintain and verify the phytosanitary status of table grapes from India. This is to ensure that the proposed risk management measures have been met and are maintained.

Details of the operational system, or equivalent, will be determined by agreement between the Australian Government Department of Agriculture and Water Resources and DAC before the commencement of trade.

5.2.1 Provision for traceability

A system of traceability to source vineyards

The objectives of this recommended procedure are to ensure that:

- table grapes are sourced only from vineyards producing commercial quality fruit
- vineyards from which table grapes are sourced can be identified so investigation and corrective action can be targeted rather than applying it to all contributing vineyards in the event that live pests are intercepted.

It is recommended that DAC establish a system to enable traceability back to the vineyards where table grapes for export to Australia are sourced from. The Department of Agriculture and Cooperation, India would be responsible for ensuring that export table grape growers are aware of pests of quarantine concern to Australia and control measures. The records of the pest control programme would need to be made available to the Australian Government Department of Agriculture and Water Resources, if requested.

Registration of packing house and treatment providers and auditing of procedures

The objectives of this recommended procedure are to ensure that:

- table grapes are sourced only from DAC registered packing houses, processing commercial quality fruit
- reference to the packing house and the vineyard source (by name or a number code) are clearly stated on cartons of table grapes destined for export to Australia for trace back and auditing purposes.

It is recommended that export packing houses and treatment providers (if applicable) are registered with DAC before the commencement of harvest each season. The list of registered packing houses and treatment providers must be kept by DAC, and would need to be made available to the Australian Government Department of Agriculture and Water Resources, if requested.

DAC would be required to audit the registered providers at the beginning of each export season to ensure that packing houses and treatment facilities are suitably equipped to carry out the specified phytosanitary activities and treatments. Records of DAC audits would be made available to the Australian Government Department of Agriculture and Water Resources, if requested.

Packing houses will be required to identify individual vineyards with a unique identifying system and identify fruit from individual vineyards by marking cartons or pallets with a unique number or identification provided by DAC.

Where table grapes undergo fruit treatment prior to export, this process could only be undertaken by the treatment providers that have been registered with and audited by DAC for the purpose.

DAC must immediately suspend exports of table grapes to Australia from packing houses/treatment providers found to be non-compliant and must notify the Australian Government Department of Agriculture and Water Resources of the suspension.

Suspended packing houses/treatment providers may only be re-instated for processing of table grapes for export to Australia when DAC and the Australian Government Department of Agriculture and Water Resources are satisfied that non-compliance issues have been adequately addressed.

5.2.2 Packaging and labelling

The objectives of this recommended procedure are to ensure that:

- table grapes proposed for export to Australia and all associated packaging is not contaminated by quarantine pests or regulated articles
 - regulated articles are any items other than table grapes. Regulated articles may include plant, plant product, soil and any other organisms, object or material capable of harbouring or spreading pests, deemed to require phytosanitary measures, particularly where international transportation is involved
 - in this report, table grapes is defined as table grape bunches or clusters, which include peduncles, rachises, laterals, pedicels and berries (Pratt 1988), but not other plant parts (section 1.2.2)
- unprocessed packing material (which may vector pests identified as not being on the pathway and pests not known to be associated with table grape bunches) is not imported with the table grapes
- all wood material used in packaging of table grapes complies with the Australian Government Department of Agriculture and Water Resources conditions
- secure packaging is used during storage and transport to Australia and must meet Australia's general import conditions for fresh fruits and vegetables, available on the Australian Government Department of Agriculture and Water Resources website
- the packaged table grapes are labelled with the vineyard source (by name or a number code), packing house registration number for the purposes of trace back
- the phytosanitary status of table grapes must be clearly identified.

5.2.3 Specific conditions for storage and movement

The objectives of this recommended procedure are to ensure that:

- table grapes for export to Australia that have been treated and/or inspected are kept secure and segregated at all times from any fruit for domestic or other markets, untreated/non-cleared product, product to prevent mixing or cross-contamination
- the quarantine integrity of the commodity during storage and movement is maintained.

5.2.4 Freedom from trash

All table grapes for export must be free from trash (for example, stem and leaf material, seeds, soil, animal matter/parts or other extraneous material) and foreign matter. Freedom from trash

will be confirmed by the inspection procedures. Export lots or consignments found to contain trash and/or foreign matter should be withdrawn from export unless approved remedial action is available and applied to the export consignments and then re-inspected.

5.2.5 Pre-export phytosanitary inspection and certification by DAC

The objectives of this recommended procedure are to ensure that:

- all consignments have been inspected in accordance with official procedures for all visually detectable quarantine pests and other regulated articles (including soil, animal and plant debris) at a standard 600 unit sampling rate per phytosanitary certificate, or equivalent, whereby one unit is one bunch of table grapes
- an international phytosanitary certificate (IPC) is issued for each consignment upon completion of pre-export inspection and treatment to verify that the relevant measures have been undertaken offshore
- each IPC includes:
 - a description of the consignment (including orchard registration number or reference code and packing house details)
 - details of disinfestation treatments (for example irradiation or pre-shipment/in-transit cold treatment) which include date, temperature, dose, duration, and/or treatment certificate (as appropriate)
- an additional declaration that *'The fruit in this consignment has been produced in India in accordance with the conditions governing entry of fresh table grapes to Australia and inspected and found free of quarantine pests and regulated articles'*
- another statement may be required, where irradiation is used. Additional information about live pests detected in the consignments during NPPO inspection must also be included on the IPC, as the treatment is performed following NPPO inspection on the shipment.

5.2.6 Verification by the Australian Government Department of Agriculture and Water Resources

The objectives of this recommended procedure are to ensure that:

- all consignments comply with Australian import requirements
- consignments are as described on the phytosanitary certificate and quarantine integrity has been maintained.

To verify that phytosanitary status of consignments of table grapes from India meets Australia's import conditions, it is recommended that the Australian Government Department of Agriculture and Water Resources complete a verification inspection of all consignments of table grapes. It is recommended that the department randomly sample 600 fruit per phytosanitary certificate.

The detection of any quarantine pest or regulated article for Australia would require suitable remedial action.

5.2.7 Remedial action(s) for non-compliance

The objectives of remedial action(s) for non-compliance are to ensure that:

- any quarantine risk is addressed by remedial action, as appropriate
- non-compliance with import requirements is addressed, as appropriate.

Any consignment that fails to meet Australia's import conditions must be subject to a suitable remedial treatment, if one is available, re-exported from Australia, or destroyed.

Separate to the corrective measures mentioned, there may be other breach actions necessary depending on the specific pest intercepted and the risk management strategy put in place against that pest in the protocol.

If product repeatedly fails inspection, the Australian Government Department of Agriculture and Water Resources reserves the right to suspend the export programme and conduct an audit of the risk management systems. The programme will recommence only when the department is satisfied that appropriate corrective action has been taken.

5.3 Responsibility of competent authority

The Department of Agriculture and Cooperation (DAC) of India is the designated NPPO under the International Plant Protection Convention (IPPC).

The NPPO's responsibilities include:

- inspecting plants and plant products moving in international trade
- issuing certificates relating to phytosanitary condition and origin of consignments of plants and plant products
- ensuring that all relevant agencies participating in this programme meet the recommended service and certification standards and recommended work plan procedures
- ensuring that administrative processes are established to meet the requirements of the programme.

5.3.1 Use of accredited personnel

Operational components and the development of risk management procedures may be delegated by DAC to an accredited agent under an agency arrangement as appropriate. This delegation must be approved by the Australian Government Department of Agriculture and Water Resources. DAC is responsible for auditing delegated risk management procedures.

The accrediting authority must provide DAC with the documented criteria upon which accreditation is based and this must be available for audit by DAC. The Australian Government Department of Agriculture and Water Resources may verify the accrediting system before the commencement of trade.

5.4 Uncategorized pests

If an organism, including contaminant pests, is detected on table grapes either in India or on-arrival in Australia that has not been categorised, it will require assessment by the Australian Government Department of Agriculture and Water Resources to determine its quarantine status and whether phytosanitary action is required. Assessment is also required if the detected species was categorised as not likely to be on the import pathway. If the detected species was categorised as on the pathway but assessed as having an unrestricted risk that achieves Australia's ALOP due to the rating for likelihood of importation, then it would require reassessment. The detection of any pests of quarantine concern not already identified in the analysis may result in remedial action and/or temporary suspension of trade while a review is conducted to ensure that existing measures continue to provide the appropriate level of protection for Australia.

5.5 Review of processes

5.5.1 Verification of protocol

Prior to the first season of trade, the Australian Government Department of Agriculture and Water Resources will verify the implementation of agreed import conditions and phytosanitary measures including registration, operational procedures and treatment providers, where applicable. This may involve representatives from the Australian Government Department of Agriculture and Water Resources visiting areas in India that produce table grapes for export to Australia.

5.5.2 Review of policy

The Australian Government Department of Agriculture and Water Resources reserves the right to review the import policy after the first year of trade or when there is reason to believe that the pest or phytosanitary status in India has changed.

DAC must inform the Australian Government Department of Agriculture and Water Resources immediately on detection in India of any new pests of table grapes that are of potential quarantine concern to Australia or of a significant change in the application of existing commercial practices considered in this report.

5.6 Meeting Australia's food laws

Imported food for human consumption must comply with the requirements of the *Imported Food Control Act 1992*, as well as Australian state and territory food laws. Among other things, these laws require all food, including imported food, to meet the standards set out in the Australia New Zealand Food Standards Code (the Code).

The Australian Government Department of Agriculture and Water Resources administers the *Imported Food Control Act 1992*. This legislation provides for the inspection and control of imported food using a risk-based border inspection program, the Imported Food Inspection Scheme. More information on this inspection scheme, including the testing of imported food, is available from the [department's website](#).

Food Standards Australia New Zealand (FSANZ) is responsible for developing and maintaining the Code, including Standard 1.4.2 - Agvet chemicals. This standard is available on the [Federal Register of Legislation](#) or through the [FSANZ website](#).

Standard 1.4.2 and Schedules 20 and 21 of the Code set out the maximum residue limits (MRLs) and extraneous residue limits (ERLs) for agricultural or veterinary chemicals that are permitted in food, including imported food.

Standard 1.1.1 of the Code specifies that a food must not have, as an ingredient or a component, a detectable amount of an Agvet chemical or a metabolite or a degradation product of the Agvet chemical; unless expressly permitted by the Code.

Anyone may apply to change the Code whether they are an individual, organisation or company. The application process, including the explanation of establishment of MRLs in Australia, is described at the [FSANZ website](#).

6 Conclusion

The findings of this final report for a non-regulated analysis of existing policy for table grapes from India are based on a comprehensive scientific analysis of relevant literature.

The Australian Government Department of Agriculture and Water Resources considers that the risk management measures recommended in this report will provide an appropriate level of protection against the pests identified as associated with the trade of table grapes from India.

Appendix A Initiation and categorisation for pests of table grapes from India

The steps in the initiation and categorisation processes are considered sequentially, with the assessment terminating at 'Yes' for column 3 (except for pests that are present, but under official control and/or pests of regional concern) or the first 'No' for columns 4, 5 or 6.

Details of the method used in this risk analysis are given in Section 2: Method for pest risk analysis.

This pest categorisation table does not represent a comprehensive list of all the pests associated with the entire plant of an imported commodity. Reference to soilborne nematodes, soilborne pathogens, wood borer pests, root pests or pathogens, and secondary pests have not been listed as they are not directly related to the export pathway of fresh table grapes and would be addressed by Australia's current approach to contaminating pests.

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|--------------------------------------|--------------------------|--|--|-------------------------------------|-------------------------------|
| ARTHROPODS | | | | | | |
| Coleoptera | | | | | | |
| <i>Adoretus bicolor</i> (Brenske, 1900) [Scarabaeidae] Cockchafer beetle | Yes (Ahmed, Kumar & Dharmaraju 1977) | No records found | No Although adults can puncture the berries, they mainly feed on the foliage and at night (Ahmed, Kumar & Dharmaraju 1977). They are not likely to be feeding on grapes at the time of harvest. | Assessment not required | Assessment not required | No |
| <i>Adoretus versutus</i> Harold, 1869 [Scarabaeidae] Rose beetle | Yes (CABI 2012) | No records found | No Larvae feed underground on roots of vine and surrounding vegetation; adults feed on foliage (CABI 2012). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|-------------------------|--|---|---|--|--------------------------------------|
| <i>Carpophilus humeralis</i> (Fabricius, 1758) [Nitidulidae] Pineapple sap beetle | Yes (DPP 2007) | Yes (Hossain & Williams 2003; Poole 2010) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Cerosterna scabrator</i> (Fabricius, 1781) [Cerambycidae] Babul-root boring longicorn | Yes (DPP 2007) | No records found | No The adult beetles feed on the outer bark and lay eggs on the trunk of the vine, or stem and the larvae bore directly into the stem (NHB 2009). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| <i>Chlorophorus annularis</i> (Fabricius, 1787) [Cerambycidae] Bamboo longhorn beetle | Yes (CABI 2012) | Yes (McKeown 1947) NSW, Qld, Vic (Plant Health Australia 2001) Not known to be present in WA (Poole 2010). | No Larvae of this species attack roots and stems, while adult beetles feed on flowers (Walker 2008). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| <i>Melolontha melolontha</i> (Linnaeus, 1758) [Scarabaeidae] White grub cockchafer | Yes (DPP 2007) | No records found | No Larvae feed underground on roots of vine and surrounding vegetation; adults feed on foliage (CABI 2012). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|-------------------------|---|---|---|--|--------------------------------------|
| <i>Phyllophaga</i> spp. [Scarabaeidae] White grubs | Yes (DPP 2007) | No <i>Phyllophaga</i> is a very large genus of scarab beetles in the subfamily Melolonthinae. In Australia, the subfamily Melolonthinae is represented by the genera <i>Dermolepida</i> and <i>Lepidiota</i> , rather than <i>Phyllophaga</i> (CABI 2015). | No Larvae feed on roots of its hosts and adults feed on foliage and fruits of orchard trees (CABI 2012). However, no records have been found which associate this genus with grape bunches. | Assessment not required | Assessment not required | No |
| <i>Scelodonta strigicollis</i> (Motschulsky, 1866) [Eumolpidae] Grape flea beetle | Yes (Bournier 1977) | No records found | No Larvae feed underground on roots of vine and the adults feed on the leaves, specifically the buds (Bournier 1977). They have only been reported to scrape unripe berries (Kulkarni 1971) and would not be expected to be present at time of harvest for mature berries. | Assessment not required | Assessment not required | No |
| <i>Sinoxylon anale</i> (Lesne, 1897) [Bostrichidae] Auger beetle | Yes (Mathew 1987) | Yes (CSIRO 2005) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|--|---|---|---|--|--------------------------------------|
| <i>Sthenias grisator</i> (Fabricius, 1784) [Cerambycidae] Grapevine girdler, Long-horned beetle | Yes (NHB 2009) | No records found | No Larvae feed underground on roots of vine and surrounding vegetation, while the adults girdle around the main stem around 15 centimetres above ground level (NHB 2009). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| <i>Xylopsocus capucinus</i> (Fabricius, 1781) [Bostrichidae] False powderpost beetle | Yes (Woodruff, Gerberg & Spilman 2011) | Yes NSW, NT (Plant Health Australia 2001) Not known to be present in WA. | No Larvae feed on roots and adults bore into stems (Woodruff, Gerberg & Spilman 2011). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| <i>Xylothrips flavipes</i> (Illiger, 1801) [Bostrichidae] Auger beetle | Yes (Walker 2011) | Yes NSW, Qld, Vic., NT (Plant Health Australia 2001) Not known to be present in WA. | No Adult and larvae bore into wood, for example trunk of vine (Walker 2011). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| <i>Xylosandrus crassiusculus</i> (Motschulsky, 1866) [Bostrichidae] Asian ambrosia beetle | Yes (Keshavareddy, Vergheese & Reddy 2007) | No records found | No Adult and larvae bore into stems and trunks of host plant (CABI 2012). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|--|---------------------------------|--|--|--|--------------------------------------|
| <i>Xylosandrus compactus</i> (Eichhoff 1875) [Bostrichidae] Shot-hole borer, Black twig borer | Yes (Keshavareddy, Verghese & Reddy 2007) | No records found | No Adult and larvae bore into stems and trunks of vine (CABI 2012). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| Diptera | | | | | | |
| <i>Bactrocera correcta</i> (Bezzi 1916) [Tephritidae] Guava fruit fly | Yes (CABI 2012; Verghese et al. 2002) | No records found | Yes Has been known to infect grapes (Mani 1992). A survey conducted in India to examine natural pests affecting ripe grape berries identified the emergence of <i>Bactrocera correcta</i> in insect cages for rearing purposes (Mani 1992). | Yes Known to be present in Sri Lanka, Pakistan, Nepal and Thailand (White & Elson-Harris 1992). In India it forms part of the <i>B. dorsalis</i> complex, which is important in terms of the pest's distribution, diverse host range, rapid population build up and potential economic damage (Satarkar et al. 2009). | Yes In India, this potential pest often occurs with serious pest species such as <i>B. zonata</i> and <i>B. dorsalis</i> (Kapoor 2002). This pest complex is considered one of the most important in world agriculture (Satarkar et al. 2009). <i>Bactrocera correcta</i> is known to affect citrus, mango, sandalwood, guava, peach (White & Elson-Harris 1992) and grapes (Mani 1992). | Yes (EP) |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|---|---|---|---|---|--------------------------------------|
| <i>Bactrocera dorsalis</i> (Hendel, 1912) [Tephritidae] Oriental fruit fly | Yes (CABI 2012; Verghese et al. 2002; Verghese, Tandon & Stonehouse 2004) | No records found | Yes Damage caused by <i>B. dorsalis</i> consists of punctures of the host tissue by adults during oviposition. They lay their eggs under the skin of fruits, and larvae subsequently feed on the fruit pulp (Chu & Tung 1996; Ye & Liu 2005). | Yes <i>Bactrocera dorsalis</i> has significant potential to become established and spread through areas of Australia. This is best shown by an incursion of the closely allied papaya fruit fly (<i>B. papayae</i>) (Drew and Hancock, 1994) in north Queensland during the mid-1990s. | Yes <i>Bactrocera dorsalis</i> can utilise more than 150 fruit species (Waite 2009). It is considered one of the five most important pests of agriculture in South East Asia (Waterhouse 1993). Females oviposit into the fruit of hosts, eggs hatch inside the fruit and the larvae consume the fruit pulp (CABI 2012). | Yes (EP) |
| <i>Drosophila melanogaster</i> Meigen, 1830 [Drosophilidae] Common fruit fly, Vinegar fly | Yes (DPP 2007) | Yes (CSIRO 2005) NSW, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Drosophila suzukii</i> Matsumura 1931 [Drosophilidae] Spotted wing drosophila | Yes (Guruprasad, Hegde & Krishna 2010) | No records found | A pest risk assessment for <i>D. suzukii</i> will not be conducted in this risk analysis report for table grapes from India. There is existing policy for <i>D. suzukii</i> for all commodities, including table grapes, from all countries (Department of Agriculture 2013b). A summary of pest information and previous assessment is presented in Chapter 4 of this report. Further information on existing policy can be found in the 'Final pest risk analysis report for <i>Drosophila suzukii</i> ', published on 24 April 2013 (Department of Agriculture 2013b). | | | |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|------------------------------------|--|--|---|--|--------------------------------------|
| Hemiptera | | | | | | |
| <i>Aleurocanthus spiniferus</i> (Quaintance, 1903) [Aleyrodidae] Citrus blackfly | Yes (DPP 2007) | Yes NT, Qld (ABRS 2009; CABI 2015; Plant Health Australia 2001) No records found for WA. | No <i>Aleurocanthus spiniferus</i> is mainly associated with leaves (CABI & EPPO 1997a), primarily of citrus (Gyeltshen, Hodges & Hodges 2008). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| <i>Aleurocanthus woglumi</i> Ashby, 1915 [Aleyrodidae] Citrus blackfly | Yes (CABI 2012; CABI & EPPO 1997b) | No records found | No Although grapes are considered a minor host, <i>A. woglumi</i> is most commonly found on vegetative material, such as leaves and stems (CABI 2012; Plant Health Australia 2009b). Plant Health Australia (2009b) considered that this pest was not on the pathway. Eggs are laid on leaves and nymphs feed on the underside of leaves (CABI & EPPO 1997b). | Assessment not required | Assessment not required | No |
| <i>Aleurodicus dispersus</i> Russell, 1965 [Aleyrodidae] Spiralling whitefly | Yes (CABI 2012) | Yes NT, Qld (ABRS 2009; CSIRO 2005) | No Adults and nymphs only feed on leaves (CABI 2012). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|---|--|-----------------------------------|---|--|--------------------------------------|
| <i>Aleurolobus taeonabae</i> (Kuwana, 1911) [Aleyrodidae] Whitefly | No. Dubey and Ko (2009) reported <i>Aleurolobus taeonabae</i> as being present in India, however no actual records have been found. India has stated that <i>A. taeonabe</i> is absent from India (DPP 2012). Note: If the NPPO of India becomes aware of any records of this pest in India, it must inform Australia immediately and this pest categorisation may need to be reviewed accordingly. | No records found | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Aonidiella citrina</i> (Coquillett, 1891) [Diaspididae] Yellow scale | Yes (EPPO 2011) | Yes NSW, SA, Vic., WA (CABI 2012; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|---------------------------|--|--|---|--|--------------------------------------|
| <i>Aonidiella orientalis</i> (Newstead, 1894) [Diaspididae] Oriental yellow scale, Oriental scale | Yes (CABI 2012; DPP 2007) | Yes (CSIRO 2005) NT, Qld, WA (Plant Health Australia 2001). It is present in Northern Australia as far south as 24°S (Gladstone) (Astridge & Elder 2005). | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Amrasca biguttula biguttula</i> (Ishida, 1912) [Cicadellidae] Leafhopper | Yes (NRC 2013) | No records found | No The leafhopper <i>Amrasca biguttula biguttula</i> is associated with grapevine leaves in India (NRC 2013). It is associated with leaves of other hosts (CABI 2013) and no records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| <i>Aphis fabae</i> Scopoli, 1763 [Aphidae] Black bean aphid | Yes (DPP 2007) | No records found | No While this species attacks grapevine (USDA-APHIS 2002), it rests and feeds on leaves (Miles 1987) and is not associated with grape bunches (Ingels et al. 1998). | Assessment not required | Assessment not required | No |
| <i>Aphis gossypii</i> Glover, 1877 [Aphidae] Cotton aphid | Yes (DPP 2007) | Yes NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|-----------------------------------|--|--|---|--|--------------------------------------|
| <i>Aphis spiraecola</i> Patch, 1914 [Aphidae] Green citrus aphid | Yes (DPP 2007) | Yes NSW, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Arboridia viniferata</i> Sohi & Sandhu, 1971 [Cicadellidae] Grapevine cicadellid | Yes (Sohi, Bindra & Sohi 1975) | No records found | No This pest mainly attacks leaves (Sohi, Bindra & Sohi 1975). While individuals may at times be on grape bunches, they are likely to jump off bunches of grapes during harvesting and grading. | Assessment not required | Assessment not required | No |
| <i>Aspidiotus destructor</i> Signoret, 1869 [Diaspididae] Coconut scale | Yes (DPP 2007) | Yes NSW, NT, Qld, Vic, WA (Plant Health Australia 2001) WA (Poole 2010) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Ceroplastes rusci</i> (Linnaeus, 1758) [Coccidae] Fig wax scale | Yes (DPP 2007) | Yes NT (CSIRO 2005; Plant Health Australia 2001) Permitted into WA (Government of Western Australia 2014) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|-------------------------|--|-----------------------------------|---|--|--------------------------------------|
| <i>Chrysomphalus dictyospermi</i> [Diaspididae] Dictyospermum scale | Yes (Butani 1993) | Yes NSW, NT, Qld, SA (Plant Health Australia 2001) No records found for WA. However, WA does not require mitigation measures for this pest for other hosts (such as citrus, peach or nectarine fruit) from Australian states where this pest is present (DAFWA 2014). | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Coccus hesperidum</i> Linnaeus, 1758 [Coccidae] Brown soft scale | Yes (CABI 2012) | Yes ACT, NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|------------------------------|--|---|--|--|--------------------------------------|
| <i>Daktulosphaira vitifoliae</i> (Fitch, 1855) Synonym: <i>Viteus vitifoliae</i> (Fitch, 1855) [Phylloxeridae] Grapevine phylloxera | Yes (CABI 2012; DPP 2007) | Yes Present only in isolated areas of Vic. and NSW. The pest is under official control in these areas and strict quarantine conditions apply (NVHSC 2005; PGIBSA 2009). | Yes The first instar 'crawler' stage is the most dispersive stage and can be found on the soil surface and on the foliage or fruit of vines (Buchanan & Whiting 1991). | Yes <i>Daktulosphaira vitifoliae</i> is already established in small areas of Australia, where it is under official control (NVHSC 2008). In Australia, several generations develop in each growing season (NVHSC 2005). <i>Daktulosphaira vitifoliae</i> can be spread by human activities, notably movement of grapevine nursery stock and related products including soil associated with infested roots (for example carried on footwear or vehicle tyres). Harvesting machinery, other equipment and tools are also implicated with its spread (NVHSC 2005). The potential for spread on harvested table grapes is also a concern (Buchanan & Whiting 1991). | Yes <i>Daktulosphaira vitifoliae</i> only causes direct harm to grapevines (<i>Vitis</i> spp.). The only reliable control measure for <i>D. vitifoliae</i> is the complete removal of infested vines and their replacement with grapevines grown on resistant rootstock (Buchanan & Whiting 1991). | Yes (EP) |
| <i>Diaspidiotus perniciosus</i> (Comstock, 1881) [Diaspididae] San José scale | Yes (CABI 2012) | Yes NSW, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) WA (Poole 2010) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|-------------------------------|---|--|---|--|--------------------------------------|
| <i>Dysmicoccus brevipes</i> (Cockerell, 1893) [Pseudococcidae] Pineapple mealybug | Yes (Mani & Thontadarya 1987) | Yes NSW, NT, Qld, SA, WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Empoasca fabae</i> (Harris, 1841) Synonym: <i>Empoasca mali</i> (Baron, 1853) [Cicadellidae] Potato leaf hopper | Yes (Prasad 1960) | No records found | No <i>Empoasca fabae</i> can cause significant injury to vineyards, causing leaf cupping, reduced shoot growth, and leaf yellowing (Integrated Pest Management Center 2007; Isaacs 2007; Isaacs & van Timmeren 2009). Adults are very active, jumping or flying when disturbed. The immature forms, or nymphs run forward, backward or sideways when disturbed (Isaacs 2007). This pest is unlikely to remain on the host during harvesting. | Assessment not required | Assessment not required | No |
| <i>Empoasca vitis</i> (Göthe, 1875) [Cicadellidae] Smaller green leafhopper, Vine leaf hopper | Yes (DPP 2007) | No records found | No Attacks leaves and feeding causes scorching (CABI 2012; Pavan et al. 1998). This pest is unlikely to remain on the host during harvesting. | Assessment not required | Assessment not required | No |
| <i>Eulecanium tiliae</i> (Linnaeus, 1758) [Coccidae] Nut scale, Brown gooseberry scale | Yes (DPP 2007) | Yes (ALA 2013) Tas., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|--|--|--|---|--|--------------------------------------|
| <i>Ferrisia virgata</i> (Cockerell, 1893) [Pseudococcidae] Striped mealybug | Yes (DPP 2007) | Yes (CABI 2012; Poole 2010) NSW, NT, Qld, WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Halyomorpha halys</i> (Stål, 1855) [Pentatomidae] Brown marmorated stink bug | Yes (Nielsen, Hamilton & Matadha 2008) | No records found | No In grapes, <i>H. halys</i> adults suck sap from the fruit and the nymphs feed on leaves, stems and fruit (Zhang 2005). Pentatomid bugs are not likely to be carried by fruit because they characteristically drop from their hosts when disturbed, or fly off (Alcock 1971). | Assessment not required | Assessment not required | No |
| <i>Helopeltis antonii</i> Signoret, 1858 [Miridae] Tea bug | Yes (Devasahayam & Nair 1986) | No records found | No Feeds on tender leaves and developing fruits (Puttarudriah & Appanna 1955). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| <i>Hemiberlesia lataniae</i> (Signoret, 1869) Synonyms: <i>Aspidiotus lataniae</i> (Signoret, 1869); <i>Aspidiotus cydoniae</i> (Comstock, 1881) [Diaspididae] Latania scale, Quince scale | Yes (DPP 2007; Miller & Davidson 2005) | Yes (CSIRO 2005) NSW, NT, Qld, Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|---|---|---|---|--|--------------------------------------|
| <i>Icerya purchasi</i> (Maskell, 1876) [Monophlebidae] Cottony cushion scale | Yes (Kapur 1949; Verma, Chand & Kumar 2012) | Yes NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Icerya seychellarum</i> (Westwood, 1855) [Monophlebidae] Seychelles scale | Yes (DPP 2007) | Yes (ALA 2013; DAFWA 2008) NSW, NT, Qld, WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Jacobiasca lybica</i> (Bergevin & Zanon, 1922) Synonym: <i>Chlorita lybica</i> (Bergevin & Zanon 1922) [Cicadellidae] Cotton jassid | Yes (DPP 2007) | No records found | No Attacks leaves and feeding causes scorching (INRA 1997). This species is unlikely to remain on the host during harvesting. | Assessment not required | Assessment not required | No |
| <i>Maconellicoccus hirsutus</i> (Green, 1908) [Pseudococcidae] Pink hibiscus mealybug | Yes (DPP 2007; Mani & Thontadarya 1987) | Yes NT, Qld, Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Macrosiphum euphorbiae</i> (Thomas, 1878) [Aphididae] Potato aphid | Yes (DPP 2007) | Yes ACT, NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|------------------------|--|---|--|-------------------------------------|-------------------------------|
| <i>Myzus persicae</i> (Sulzer, 1776) [Aphididae] Green peach aphid | Yes (CABI & EPPO 1979) | Yes NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) <i>Myzus persicae</i> can vector <i>Broad bean wilt virus 2</i> (BBWV 2)(Zhou 2002), a potential virus of grapevine (Martelli 1999). BBWV 2 is present in India (CABI 2012; Mali, Vyanjane & Ekbote 1977) and is also present in NSW (Schwinghamer et al. 2007) and may be present in Qld (Plant Health Australia 2001), but is not known to occur in WA. Although <i>M. persicae</i> is present in Australia, the potential for <i>M. persicae</i> carrying BBWV 2. warrants further assessment for this species. | No Although reported from grapes in spring, <i>M. persicae</i> is likely to be present only as transients (Flaherty et al. 1992). Watson (1923) reported <i>M. persicae</i> on the leaves and tender stems of grapevine, but did not consider this species to be a berry feeder. <i>Myzus persicae</i> has been reported on grapevine flower clusters in California on one occasion (Flaherty et al. 1992). It has not been reported feeding on grape bunches but has been reported on the fruit of other hosts (Gildow et al. 2004). Also, <i>M. persicae</i> can only vector BBWV 2 for a maximum of two hours after feeding (Zhou 2002). No records have been found of virus acquisition from infected berries by <i>M. persicae</i> . | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|---|---|---|---|--|--------------------------------------|
| <i>Nipaecoccus nipae</i> (Maskell, 1893b) [Pseudococcidae] Coconut mealybug | Yes (Ben-Dov 2012c) | No records found | No This pest occurs on the foliage of its host plants (Ben-Dov 2012c). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| <i>Nipaecoccus viridis</i> (Newstead 1894) [Pseudococcidae] Spherical mealybug | Yes (DPP 2007; Mani & Thontadarya 1987) | Yes NT, Qld, WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Nysius niger</i> Baker, 1906 [Lygaeidae] Northern false chinch bug | Yes (DPP 2007) | No records found | No Species of <i>Nysius</i> pierce the leaves and buds with toxic saliva, which damages the leaves and buds and causes leaf fall (Bournier 1977). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| <i>Parasaissetia nigra</i> (Nietner, 1861) [Coccidae] Pomegranate scale | Yes (DPP 2007) | Yes NSW, NT, Qld, SA, Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|--|--|---|--|--|--------------------------------------|
| <i>Parthenolecanium corni</i> (Bouché, 1844) [Coccidae] European fruit lecanium, Plum scale, Peach scale | Yes (Bhagat, Ramzan & Farhan 1991; CABI 2012; DPP 2007) | Yes Tas. (Plant Health Australia 2001), Vic. (Snare 2006). Not known to be present in WA (Poole 2010). | Yes This species sucks sap from branches, leaves and fruit of grapevines (Zhang 2005). Due to their small size and habit of feeding in concealed areas on plant material and fruit, they are frequent invasive species (Miller et al. 2007). | Yes This pest is widely distributed in temperate and subtropical regions (Ben-Dov 2012a). This pest is highly polyphagous, attacking some 350 plant species placed in 40 families (Ben-Dov 2012a). Many of these host plants are available in Western Australia. | Yes This pest is highly polyphagous, attacking some 350 plant species placed in 40 families (Ben-Dov 2012a). It has been observed to cause heavy infestation and damage to <i>Vitis vinifera</i> in the Kasmir Valley (Bhagat, Ramzan & Farhan 1991) and is the most widespread and injurious soft scale in French vineyards (Sforza, Boudon-Padieu & Greif 2003). Trees infested with <i>P. lecanium</i> lose leaves and decrease their annual growth while heavy infestations lead to fungal growth on the honeydew secretions (David'yan 2008). This species also transmits viruses (Ben-Dov 2012a). | Yes (EP, WA) |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|------------------------------|---|---|--|---|--------------------------------------|
| <i>Parthenolecanium persicae</i> (Fabricius, 1776), [Coccidae] Peach scale, Grapevine scale | Yes (CABI 2012) | Yes NSW, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Perissopneumon ferox</i> Newstead, 1900 [Monophlebidae] | Yes (DPP 2007) | No records found | Yes Infests the fruit stalk, inflorescence and fruit (DPP 2012). | Yes Susceptible hosts (for example mango, citrus and neem) (Ben-Dov 2012b) are present in Australia. | No In an orchard in the Lucknow district, India, <i>P. ferox</i> heavily infested mango trees in 1980 (Srivastava & Vergheese 1985). <i>Perissopneumon ferox</i> has also been reported in the 1980s in India on custard apple and guava (Shukla & Tandon 1984; Tandon & Vergheese 1987). Since then, only limited evidence has been published of this species causing economic damage on any host. Therefore, this species is not considered further. | No |
| <i>Pinnaspis strachani</i> (Cooley, 1899) [Diaspididae] Lesser snow scale | Yes (CABI 2012; DPP 2007) | Yes NSW, NT, Qld, WA (Plant Health Australia 2001) SA, WA (Brookes 1964) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|---|---|-----------------------------------|---|--|--------------------------------------|
| <i>Planococcus citri</i> (Risso, 1813) [Pseudococcidae] Citrus mealybug | Yes (DPP 2007; Mani & Thontadarya 1987) | Yes NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|-------------------------|---------------------------------|---|---|--|--------------------------------------|
| <i>Planococcus ficus</i> (Signoret, 1875) [Pseudococcidae] Grapevine mealybug | Yes (Ben-Dov 2012c) | No records found | Yes Mealybugs occupy the main stems of the vines, but move to the new growth areas, such as leaves and grape bunches as the season progresses (Walton & Pringle 2004a). They have been known to accumulate in grape clusters (Millar et al. 2002). | Yes The grapevine mealybug can have up to four to six generations per year (Millar et al. 2002) and is very polyphagous, causing damage to plants in over 11 families (Ben-Dov 2012c). The grapevine mealybug occurs in many countries including Argentina, Brazil, Egypt, France, Mexico, Russia, South Africa and United States of America (Ben-Dov 2012c). Environments with climates similar to these regions exist in various parts of Australia, suggesting that <i>P. ficus</i> has the potential to establish and spread in Australia. | Yes <i>Planococcus ficus</i> is a key pest in vineyards worldwide (Ben-Dov 2012c; Millar et al. 2002; Walton & Pringle 2004b). This pest has the ability to destroy a grape crop, cause progressive weakening of vines through early leaf loss (Walton et al. 2006; Walton & Pringle 2004b). In the last decade, economic losses from this pest in Californian vineyards have increased dramatically (Millar et al. 2002). The pest is also a major transmitter of numerous viruses and diseases (Millar et al. 2002; Walton & Pringle 2004a). It also excretes large amounts of honeydew on grapes (Walton & Pringle 2004b). | Yes (EP) |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|---|---------------------------------|---|--|---|--------------------------------------|
| <i>Planococcus lilacinus</i> Cockerell, 1905 [Pseudococcidae] Coffee mealybug | Yes (CABI 2012; MacLeod 2006; Tandon & Verghese 1987) | No records found | Yes This mealybug has been commonly recorded on grape bunches in surveys conducted in 1983 in Bangalore, India (Tandon & Verghese 1987). | Yes <i>Planococcus lilacinus</i> is extremely polyphagous, and feeds on various tropical, sub-tropical and shade trees and crops including cocoa, guava, mango, citrus, potato, coffee, custard apple, tamarind and grapes (MacLeod 2006; Tandon & Verghese 1987). This species has been reported from tropical regions around the world as well as China and Japan (Ben-Dov 2012c). Environments with climates similar to these regions exist in various parts of Australia, suggesting that <i>P. lilacinus</i> has the potential to establish and spread in Australia. | Yes <i>Planococcus lilacinus</i> is extremely polyphagous and can feed on plants in over 35 families (MacLeod 2006), including many crops which are commercially grown in Australia. This species has been identified as a serious threat to grape crops in India (Tandon & Verghese 1987). | Yes (EP) |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|-------------------------|--|--|--|--|--------------------------------------|
| <p><i>Planococcus minor</i> (Maskell, 1897)</p> <p>Synonym: <i>Planococcus pacificus</i> Cox, 1981</p> <p>[Pseudococcidae]</p> <p>Pacific mealybug</p> | Yes (DPP 2007) | <p>Yes (Ben-Dov 2012c)</p> <p>NSW, NT, Qld, SA (Plant Health Australia 2001)</p> <p>Not known to be present in WA (Poole 2010)</p> | <p>Yes</p> <p>This pest has been recorded on grape bunches in India (Batra et al. 1987).</p> | <p>Yes</p> <p><i>Planococcus minor</i> is polyphagous attacking many wild and cultivated susceptible species; 250 host species in nearly 80 families are reported as hosts (Ben-Dov 2012c; Lit, Caasi-Lit & Calilung 1998; Sugimoto 1994; Venette & Davis 2004). Susceptible hosts are freely available in Western Australia, suggesting a high probability that a suitable host would be found.</p> <p>Many species of mealybugs are considered invasive, rapidly becoming established when introduced into new areas (Miller, Miller & Watson 2002).</p> | <p>Yes</p> <p><i>Planococcus minor</i> is a pest of many economically important species (Ben-Dov 2012c; Venette & Davis 2004). It has potential to cause economic damage if introduced into Western Australia.</p> | Yes (EP, WA) |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|-------------------------------|--|-----------------------------------|---|--|--------------------------------------|
| <i>Pseudaulacaspis pentagona</i> (Targioni Tozzetti, 1886) Synonym: <i>Diaspis pentagona</i> Targioni-Tozzetti, 1886 [Diaspididae] Mulberry scale | (Miller, Denno & Gimpel 2012) | Yes NSW, Qld (Plant Health Australia 2001) No records found for WA. However, WA does not require mitigation measures for this pest for other hosts (such as stonefruit) from Australian states where this pest is present (DAFWA 2014; Poole et al. 2011). | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Pseudococcus longispinus</i> (Targioni Tozzetti, 1867) [Pseudococcidae] Long-tailed mealybug | Yes (CABI 2012; DPP 2007) | Yes NSW, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|--|---|---|--|--|--------------------------------------|
| <i>Rastrococcus iceryoides</i> (Green, 1908) [Pseudococcidae] Downey snowline mealybug, Mango mealybug | Yes (Ben-Dov 2012c; CABI 2012; DPP 2007; Williams 1989). | No records found | Yes The pest has been recorded on grapevine in India (Williams 1989, 2004). <i>Rastrococcus iceryoides</i> can be spread by humans on infested planting material (DPP 2012). Therefore, this mealybug may be present on the stems of grape bunches. | Yes When introduced into new areas, <i>Rastrococcus</i> species become particularly injurious to tropical fruit trees and other crop plants (Williams 1989). This is one of the most widespread and polyphagous of all <i>Rastrococcus</i> species (Williams 2004). | Yes <i>Rastrococcus iceryoides</i> is one of the most polyphagous species of <i>Rastrococcus</i> , occurring on plants belonging to diverse botanical families. It has been recorded attacking over 60 genera of plants in 36 families, including <i>Vitis vinifera</i> (Ben-Dov 2012c; Williams 2004). | Yes (EP) |
| <i>Saissetia coffeae</i> (Walker, 1852) [Coccidae] Hemispherical scale | Yes (DPP 2007) | Yes NSW, NT, Qld, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Saissetia oleae</i> (Olivier, 1791) [Coccidae] Black scale | Yes (Ben-Dov 2013; Suresh & Mohanasundaram 1996) | Yes NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Singhius hibisci</i> (Kotinsky, 1907) [Aleyrodidae] Hibiscus whitefly | Yes (Dubey, Ko & David 2008) | No records found | No There is no information for this pest being on the export pathway (DPP 2012). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|------------------------------------|--|--|---|--|--------------------------------------|
| <i>Typhlocyba sp.</i> [Cicadellidae] Leaf hopper | Yes (NHB 2009) | Uncertain as not identified to species level | No The nymphs and adults suck sap from the underside of leaves (NHB 2009). This pest is unlikely to remain on the host during harvesting. | Assessment not required | Assessment not required | No |
| Hymenoptera | | | | | | |
| <i>Polistes dominula</i> (Christ, 1791) Synonym: <i>Polistes dominulus</i> (Christ) [Vespidae] Paper wasp | Yes (Buck, Marshall & Cheung 2008) | Yes WA (ABRS 2009; Plant Health Australia 2001) | No Although recorded as a pest of grapevine (Cranshaw, Larsen & Zimmerman 2011), it is believed that no stage of the wasp's life cycle would be present on the commodity after harvesting and grading. The larvae feed on insects (Cranshaw 2008) and adults feed on nectar (Cranshaw, Larsen & Zimmerman 2011). | Assessment not required | Assessment not required | No |
| <i>Vespula germanica</i> (Fabricius, 1793) [Vespidae] European wasp | Yes (Das & Gupta 1989) | Yes NSW, NT, SA, Tas., Vic. (Plant Health Australia 2001) <i>V. germanica</i> has been recorded in WA (Plant Health Australia 2001), but WA has a surveillance program in place to prevent the establishment of this pest (DAFWA 2015e). | No Although recorded as a pest of grapevine (Ward 2001), it is believed that no stage of the wasp's life cycle would be present on the commodity after harvesting and grading. The larvae feed on insects and meat (Department of Primary Industry 2011). Adults feed on nectar and ripe fruits but are aggressive when disturbed (Department of Primary Industry 2011; INRA 2012). | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|-------------------------|---|---|---|--|--------------------------------------|
| Lepidoptera | | | | | | |
| <i>Agrotis ipsilon</i> (Hufnagel, 1766) [Noctuidae] Black cutworm | Yes (CABI 2012) | Yes NSW, NT, Qld, SA, Tas., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Agrotis segetum</i> (Denis & Schiffermuller, 1775) [Noctuidae] Turnip moth | Yes (DPP 2007) | No records found | No Highly polyphagous, eggs are typically laid on soil; young larvae feed on foliage of plant; older larvae complete development mostly underground on roots (CABI 2012). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| <i>Ampelophaga rubiginosa</i> Bremer & Grey 1853 [Sphingidae] Hawkmoth, grape horn worm | Yes (DPP 2007) | No records found | No Sphingids generally feed only on foliage (Common 1990) and this species has been reported on grapevines (Pittaway & Kitching 2012). The larvae feed only on leaves of grapevine (Zhang 2005). | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|--|---------------------------------|---|---|---|--------------------------------------|
| <p><i>Archips machlopi</i>s (Meyrick, 1912) Synonym: <i>Cacoecia machlopi</i>s Meyrick, 1912) [Tortricidae] Leaf rolling moth, Bell moth</p> | <p>Yes (Puttarudriah, Kataoihallimath & Chandrasekhar 1961; Vanitha, Karupuchamy & Sivasubramanian 2011; Varma 1984) <i>Archips machlopi</i>s has frequently been misidentified in the literature as <i>Archips micaceana</i>. Previous reports of <i>A. micaceana</i> in India are likely to be misidentifications of <i>A. machlopi</i>s (Meijerman & Ulenberg 2011; Robinson, Tuck & Shaffer 1994; Rose & Pooni 2004; Tuck 1990).</p> | <p>No records found</p> | <p>Yes Table grapes are a host of <i>Archips machlopi</i>s (Puttarudriah, Kataoihallimath & Chandrasekhar 1961; Zhang 1994). <i>Archips machlopi</i>s caused damage to grapevines at Bangalore and Mysore in India where the larvae fed under thin webbing on the epidermis of the leaves, the main stalks of the bunch and the berries themselves and pupated within the webbing (Puttarudriah, Kataoihallimath & Chandrasekhar 1961).</p> | <p>Yes <i>Archips machlopi</i>s larvae feed on a wide range of plants including cereals, citrus, coffee, cotton, grapes, ornamental crops, eucalypts, pome fruits, strawberry, mango and tea (Robinson et al. 2010). Many of these species are present in Australia. This pest has been reported from India, Burma, Indonesia, Malaysia, Nepal, Pakistan, Sri Lanka, Thailand and Vietnam (Bharathie 1975; Puttarudriah, Kataoihallimath & Chandrasekhar 1961; Tuck 1990; Varma 1984). Environments with climates similar to these regions exist in various parts of Australia, suggesting that <i>A. machlopi</i>s has the potential to establish and spread in Australia.</p> | <p>Yes <i>Archips machlopi</i>s has caused damage to grapevines at Bangalore and Mysore in India (Puttarudriah, Kataoihallimath & Chandrasekhar 1961). This leafroller is polyphagous and causes considerable damage to eucalyptus seedlings (Varma 1984).</p> | <p>Yes</p> |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|-------------------------|--|---|---|--|--------------------------------------|
| <i>Autographa gamma</i> (Linnaeus, 1758) [Noctuidae] Silver-Y moth | Yes (DPP 2007) | No records found | No Larvae of this species scrape the skin from grapes and feed on the fruit contents (Abdullagatov & Abdullagatov 1986). However, larvae feed at night and shelter under leaves during the day (Venette et al. 2003). | Assessment not required | Assessment not required | No |
| <i>Conogethes punctiferalis</i> (Guenée, 1854) Synonym: <i>Dichrocrosis punctiferalis</i> Guenée, 1854 [Crambidae] Yellow peach moth, Castor capsule borer | Yes (DPP 2007) | Yes (Nielsen, Edwards & Rangsi 1996) NSW, NT, Qld, SA, WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Cossus cossus</i> (Linnaeus, 1758) [Cossidae] Goat moth | Yes (CABI 2012) | No records found | No Larvae bore within stems and trunks of grapevine (CABI 2012). Cossid moth larvae feed internally on the woody parts of plants (Grichanov 2009). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|--|---------------------------------|---|--|---|--------------------------------------|
| <i>Cryptoblabes gnidiella</i> (Millière, 1867) [Pyralidae] Citrus pyralid, honeydew moth | Yes. On sorghum and <i>Azolla</i> spp. (Harari et al. 2007; Sasmal & Kulshreshtha 1985; Sharma, Singh & Nwanze 1997) | No records found | Yes <i>Cryptoblabes gnidiella</i> has only been recorded on sorghum and <i>Azolla</i> spp. in India (Sasmal & Kulshreshtha 1985; Sharma, Singh & Nwanze 1997). In countries where <i>C. gnidiella</i> is associated with grapevine, the damage caused by this pest includes direct damage to the grape clusters and larval feeding within and among the grape berries (Harari et al. 2007). Juvenile stages can be found within grape clusters (Bisotto-de-Oliveira et al. 2007). The moth is native to the Mediterranean and it is believed to have spread throughout Europe on transported fruit (Mau & Martin Kessing 1992). | Yes <i>Cryptoblabes gnidiella</i> is an incredibly polyphagous pest (Silva & Mexia 1999). Hosts include <i>Citrus</i> spp., <i>Ficus</i> spp., <i>Malus</i> spp., <i>Prunus</i> spp., <i>Punica granatum</i> , <i>Vaccinium</i> sp. and <i>Vitis</i> spp. (Bagnoli & Lucchi 2001), which are available in Australia. <i>Cryptoblabes gnidiella</i> is native to the Mediterranean region but has since spread throughout Europe, Africa, tropical and subtropical areas of the Americas, New Zealand and parts of Asia (CABI 2012; Carter 1984). | Yes It has been reported in Tuscany as the most frequent and harmful Lepidopteran species in viticulture (Bagnoli & Lucchi 2001). It has also been introduced into Israel, and is now a major pest of avocados, grapes, pomegranates and citrus (Harari et al. 2007; Yehuda, Wysoki & Rosen 1991). | Yes (EP) |
| <i>Deilephila elpenor</i> (Linnaeus, 1758) [Sphingidae] Large elephant hawkmoth | Yes (DPP 2007) | No records found | No This species has been reported on grapevines (Pittaway & Kitching 2012). However, Sphingids generally feed only on foliage (Common 1990). They oviposit on leaves while larvae feed on leaves, or occasionally stems, and pupate in the soil (Australian Museum 2009). | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|--|--|-----------------------------------|---|--|--------------------------------------|
| <i>Eudocima fullonia</i> (Clerck, 1764) Synonyms: <i>Ophideres fullonica</i> Linnaeus, 1758; <i>Otheris fullonia</i> (Clerck, 1764) [Noctuidae] Fruit-piercing moth, Fruit sucking moth, Orange piercing moth | Yes (DPP 2007) | Yes (Nielsen, Edwards & Rangsi 1996) NSW, NT, Qld, WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Eupoecilia ambiguella</i> (Hübner, 1796) Synonym: <i>Clysia ambiguella</i> Hübner (1825) [Tortricidae] | No Specimens labelled as <i>E. ambiguella</i> in the British Museum of Natural History (dating from 1889, 1890) were re-identified as <i>E. turbinaris</i> by JD Bradley in 1957 (CABI & EPPO 1986). India has stated that <i>E. ambiguella</i> is absent from India (DPP 2012). No information can be found to associate <i>E. turbinaris</i> with grapes. | Assessment not required | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Hippotion celerio</i> (Linnaeus 1758) [Sphingidae] Grapevine hawk moth, Silver-striped hawk-moth | Yes (Pittaway & Kitching 2012) | Yes NSW, NT, Tas., Vic., WA (Common 1990; Plant Health Australia 2001; Poole 2010) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|-----------------------------|--|---|---|--|--------------------------------------|
| <i>Hyles livornica</i> (Esper, 1779) [Sphingidae] | Yes (Krizek 1991) | No records found | No Feeds on leaves (Alford 2007). In plague proportions, this pest feeds on everything, including fruit (Mijuskovic & Badulovic 1960). However, this pest under those conditions was at the later larval stage, which is about 6 centimetres long, before feeding on grapes (Mijuskovic & Badulovic 1960) and these larvae would be seen and removed during harvesting and packing procedures. | Assessment not required | Assessment not required | No |
| <i>Hyphantria cunea</i> Drury, 1770 [Actiidae] <i>Mulberry moth, Fall webworm</i> | Yes (CABI 2012) | No records found | No <i>Hyphantria cunea</i> larvae feed on foliage only (FAO 2007a; Grichanov & Ovsyannikova 2009). | Assessment not required | Assessment not required | No |
| <i>Mamestra brassicae</i> (Linnaeus, 1758) Synonym: <i>Barathra brassicae</i> Linnaeus, 1758 [Noctuidae] Cabbage moth | Yes (DPP 2007) | No records found | No Larvae of this species feed only on foliage of grapevines (Ovsyannikova & Grichanov 2009b) and hide on the ground during the day (Carter 1984). | Assessment not required | Assessment not required | No |
| <i>Oraesia emarginata</i> (Fabricius, 1794) [Noctuidae] Fruit-piercing moth, Smaller oraesia | Yes (Zaspel & Branham 2008) | Yes (Nielsen, Edwards & Rangsi 1996) Qld (Plant Health Australia 2001) Not known to be present in WA (Poole 2010). | No Though this species attacks grape berries (JSAEZ 1987), it feeds only at night and is not associated with grapevine during the day (Hattori 1969; Li 2004; MAFF 2008). | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|--------------------------------|---------------------------------|--|---|--|--------------------------------------|
| <i>Orgyia postica</i> Walker, 1855 [Lymantriidae] Cocoa tussock moth | Yes (CABI 2012; DPP 2007) | No records found | No Larvae feed on leaves and pupate on leaves and stems (MAF Biosecurity New Zealand 2009). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| <i>Pergesa acteus</i> (Cramer, 1779) [Sphingidae] Hawk moth | Yes (Pittaway & Kitching 2012) | No records found | No This species feeds on grapevines (JSAEZ 1987; Pittaway & Kitching 2012). However, Sphingids oviposit on leaves while larvae feed on leaves or occasionally stems and pupate in the soil (Australian Museum 2009; Common 1990). | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|--------------------------------------|---|---|---|--|--------------------------------------|
| <i>Platyptilia ignifera</i> Meyrick, 1908 [Pterophoridae] Large grape plume moth | Yes (Sidhu, Chandra & Pathania 2010) | No records found | Yes Larvae bore into grape berries and feed internally on the fruit (MAFF 2008; Zhang 1994). | Yes Hosts, <i>Vitis</i> spp. are present in Australia. This species has 2–3 generations per year (MAFF 2008). <i>Platyptilia ignifera</i> is recorded from Japan, Taiwan (Zhang 1994) and India (Sidhu, Chandra & Pathania 2010). Environments with climates similar to these regions exist in various parts of Australia, suggesting that <i>P. ignifera</i> has the potential to establish and spread in Australia. | Yes Infestation by larvae destroys grape berries (MAFF 2008). This species is considered an economic pest in its native range (MAFF 2008; Yano 1963). | Yes |
| <i>Spirama retorta</i> (Clerck, 1764) [Noctuidae] Owlet moth, Fruit-sucking moth | Yes (Roychoudhury & Joshi 2011) | No records found | No Adults feed on fruit at night; they are not associated with grape during the day (Li 2004). | Assessment not required | Assessment not required | No |
| <i>Spodoptera exigua</i> (Hübner) [Noctuidae] Beet army worm | Yes (Phadke, Rao & Pawar 1978) | Yes NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Spodoptera litura</i> Fabricius, 1775 [Noctuidae] Taro caterpillar | Yes (DPP 2007) | Yes NSW, NT, Qld, Tas., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|-----------------------------|---|---|---|--|--------------------------------------|
| <i>Stathmopoda auriferella</i> (Walker, 1864) Synonym: <i>Stathmopoda crocophanes</i> Meyrick, 1897 [Oecophoridae] Apple heliodinid | Yes (Robinson et al. 2010) | Yes This species is synonymous with <i>S. crocophanes</i> (Kasy 1973), which is present in Qld, NSW, Tas., SA, WA (CSIRO 2005; Meyrick 1897; Plant Health Australia 2001). | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Sylepta lunalis</i> (Guenee 1854) [Pyralidae] | Yes (Odak & Dhamdhare 1970) | No records found | No It is a leafroller and only affects the foliage (Hill 1987; Odak & Dhamdhare 1970). | Assessment not required | Assessment not required | No |
| <i>Theretra clotho</i> (Drury, 1773) Synonym: <i>Sphinx clotho</i> (Drury, 1773) [Sphingidae] Hawk moth | Yes (DPP 2007) | Yes One record from Qld (Plant Health Australia 2001) No records found for WA | No This species feeds on grapevine (Pittaway & Kitching 2006; Zhang 1994). However, Sphingids oviposit on leaves while larvae feed on leaves or occasionally stems and pupate in the soil (Australian Museum 2009). Adults feed on nectar (Common 1990). | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|--------------------------------|---|---|---|--|--------------------------------------|
| <i>Theretra latreillii lucasii</i> (Walker, 1856) [Sphingidae] | Yes (Pittaway & Kitching 2012) | Yes One record of <i>Theretra latreillii</i> from Qld and eight from WA (Plant Health Australia 2001). However, it is unclear whether the few records for Australia are the same species/subspecies as the one reported for India. | No This species feeds on grapevine (Pittaway & Kitching 2006; Zhang 1994). However, Sphingids oviposit on leaves while larvae feed on leaves or occasionally stems and pupate in the soil (Australian Museum 2009). Adults feed on nectar (Common 1990). | Assessment not required | Assessment not required | No |
| <i>Theretra oldenlandiae</i> Fabricius, 1775 [Sphingidae] Impatiens hawkmoth, Vine hawkmoth | Yes (DPP 2007) | Yes NSW, NT, Qld, SA, Vic, WA (CSIRO 2005; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Xestia c-nigrum</i> (Linnaeus, 1758) [Arctiidae] Spotted cutworm | Yes (CABI 2012; DPP 2007) | No records found | No Larvae feed on foliage close to ground level at night and shelter in litter on the ground during the day (Pfeiffer 2009; Washington State University 2008). They are unlikely to be associated with the fruit at harvest (day-time) (MAF Biosecurity New Zealand 2009; Washington State University 2008). | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|------------------------------|---|--|---|--|--------------------------------------|
| <i>Zeuzera coffeae</i> Nietner, 1861 [Cossidae] Coffee carpenter | Yes (DPP 2007) | No records found | No Larvae bore into stems and trunks of grapevine (CABI 2012). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| Orthoptera | | | | | | |
| <i>Anacridium rubrispinum</i> Bei-Bienko, 1948 [Acrididae] Red-spined tree locust | Yes (DPP 2007) | No records found | No Herbivore which attacks vegetative and flowering parts of trees and bushes including grapevines (CABI 2012). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| <i>Schistocerca gregaria</i> (Forskål, 1775) [Acrididae] Desert locust | Yes (CABI 2012) | No records found | No A general herbivore, attacking all soft parts of food plants (CABI 2012). This pest is 4–5 centimetres long (Davey 1954) and likely to be easily seen by pickers at harvest. | Assessment not required | Assessment not required | No |
| Thysanoptera | | | | | | |
| <i>Haplothrips tenuipennis</i> Bagnall, 1918 [Phlaeothripidae] | Yes (Verghese & Harish 2010) | Yes One record from 1980 in NSW on lucerne (Plant Health Australia 2001). No records found for WA | No This species feeds on buds, inflorescences and leaves of mango (Srivastava 1997) and pollinates sapodilla feeding on pollen, nectar and stigmatic exudations (Mickelbart 1996). On grapevine in India, it is mainly associated with tender leaves and flowers (Verghese & Harish 2010). | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|--|--|--|---|---|--------------------------------------|
| <i>Karnyothrips flavipes</i> Jones, 1912 [Phlaeothripidae] | Yes (Verghese & Harish 2010) | Yes. Qld (Plant Health Australia 2001) No records found for WA | No On grapevine in India, this species is mainly associated with tender leaves and flowers (Verghese & Harish 2010). It is a predatory thrips feeding on other arthropods on a wide variety of plants (Jaramillo et al. 2010; Pitkin 1976). | Assessment not required | Assessment not required | No |
| <i>Retithrips syriacus</i> (Mayet 1890) [Thripidae] Black vine thrips | Yes (CABI 2012; DPP 2007; Lal & Pillai 1981) | No records found | Yes This species is principally a pest of grapevine (Medina-Gaud & Franqui 2001). It has been recorded causing serious levels of berry scarring and heavy yield losses in India (Reddy 2006). | Yes Grapes are considered a major host for this species and it occurs in large numbers in Southern India (Mound 2005). It affects various crops such as cotton, roses and cassava, and has been observed breeding on <i>Ricinus sp.</i> and <i>Diospyros kaki</i> (Lal & Pillai 1981). Some of these hosts are present in Australia. <i>Retithrips syriacus</i> is found in Brazil, India, Iraq, Israel, Tunisia and USA (CABI 2012). Environments with climates similar to these regions exist in various parts of Australia, suggesting that <i>R. syriacus</i> has the potential to establish and spread in Australia. | Yes It has been listed as an insect species which is responsible for heavy losses in grapevines, affecting both the yield and aesthetic look of the grapes (Reddy 2006). <i>Retithrips syriacus</i> affects its hosts by defoliating and shrivelling the leaves, marring the fruit with scars and staining the fruit by abdomen droplets which contain faeces (Medina-Gaud & Franqui 2001). | Yes (EP) |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|--|---------------------------------|--|--|---|--------------------------------------|
| <i>Rhipiphorothrips cruentatus</i> Hood, 1919 [Thripidae] Grapevine thrips, Rose thrips | Yes (Bindra & Varma 1979; Bournier 1977; CABI 2012; DPP 2007; Lakra & Dahiya 2000) | No records found | Yes <i>Rhipiphorothrips cruentatus</i> is a polyphagous species feeding on the fruit, stems and leaves of various crops such as grape, guava, mango and jamun (Dahiya & Lakra 2001). The grape berries develop a corky layer and become brown (Kulkarni, Mani & Banerjee 2007). | Yes <i>Rhipiphorothrips cruentatus</i> is a polyphagous species attacking a number of commercial host plants (Dahiya & Lakra 2001) including cashew nut, sugarapple, mango, pomegranate and guava (CABI 2012), some of which are present in Australia. <i>Rhipiphorothrips cruentatus</i> is found in Afghanistan, Bangladesh, China, India, Oman, Myanmar, Pakistan, Sri Lanka and Thailand (CABI 2012). Environments with climates similar to these regions exist in various parts of Australia, suggesting that <i>R. cruentatus</i> has the potential to establish and spread in Australia. | Yes <i>Rhipiphorothrips cruentatus</i> is a serious pest of grapevine in the Punjab, sucking the sap from the lower surface of the leaves and causing russetting and scarring of grapes (Batra et al. 1980). | Yes (EP) |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|--|--|--|---|--|--------------------------------------|
| <i>Scirtothrips citri</i> (Moulton, 1909) [Thripidae] Citrus thrips, California citrus thrips | Yes (CABI 2012; Dadmal et al. 2001). On citrus (Dadmal et al. 2001) | No records found | No <i>Scirtothrips citri</i> has only been recorded on citrus in India (Dadmal et al. 2001). It is associated with grapevine, but grapevine is not a breeding host (CABI 2014). Records of <i>S. citri</i> on grapevine appear to be limited to the southern part of North America where it is considered to be a minor pest of grapevine (Cline 1986). This thrips seems to require access to soft green tissue (except for pupation), so only seedlings or cuttings are likely to carry the pest. Only young fruit are attacked. There is no direct evidence that this species has been spread beyond its native range by human activity (CABI & EPPO 1997c). No record of an association with grape bunches was found.(Cline 1986) | Assessment not required | Assessment not required | No |
| <i>Scirtothrips dorsalis</i> Hood, 1919 [Thripidae] Chilli thrips, Oriental tea thrips, Castor thrips, Strawberry thrips | Yes (CABI 2012) | Yes NSW, NT, Qld (Plant Health Australia 2001), WA (Government of Western Australia 2013) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Selenothrips rubrocinctus</i> (Giard 1901) [Thripidae] Red-banded thrips | Yes (CABI 2012) | Yes NSW, NT, Qld, SA, WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|------------------------------|---|---|---|--|--------------------------------------|
| <i>Streothrips arorai</i> Bhatti [Aeolothripidae] | Yes (Verghese & Harish 2010) | No records found | No This family of thrips feeds on floral tissues and insects (Mound 2009). | Assessment not required | Assessment not required | No |
| <i>Thrips hawaiiensis</i> (Morgan, 1913) [Thripidae] Banana flower thrips, Hawaiian flower thrips | Yes (DPP 2007) | Yes NSW, NT, Qld, SA, Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Thrips tabaci</i> Lindeman 1889 [Thripidae] Potato thrips, Onion thrips | Yes (DPP 2007) | Yes NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001; Poole 2010) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Xylaplothrips</i> sp. [Phlaeothripidae] | Yes (Verghese & Harish 2010) | Uncertain as not identified to species level. Some species of <i>Xylaplothrips</i> are present in Australia-ACT, NSW, WA (ABRS 2009), Qld, (ABRS 2009; Plant Health Australia 2001). | No Species of <i>Xylaplothrips</i> feed on fungi, other insects and grain seeds (Collyer 1976; ICRISAT 1985; Singh, Chakravorty & Varatharajan 2010). On grapevine in India, this species is mainly associated with tender leaves and flowers (Verghese & Harish 2010). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|--|---|-----------------------------------|---|--|--------------------------------------|
| Trombidiformes | | | | | | |
| <i>Brevipalpus californicus</i> (Banks, 1904) [Tenuipalpidae] Citrus flat mite | Yes (DPP 2007) | Yes NSW, NT, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Brevipalpus lewisi</i> McGregor, 1949 [Tenuipalpidae] Citrus flat mite, Grape bunch mite | Yes (Dhooria, Bhullar & Mallik 2005) | Yes NSW, SA, Vic., WA (Plant Health Australia 2001; Poole 2010) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Brevipalpus phoenicis</i> (Geijskes, 1930) [Tenuipalpidae] False spider mite | Yes (Rather 1999) | Yes NT, NSW, SA (Plant Health Australia 2001) WA (Poole 2010) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Calepitrimerus vitis</i> (Nalepa, 1905) [Eriophyidae] Grape leaf rust mite | Yes (DPP 2007) | Yes NSW, SA, Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Colomerus vitis</i> (Pagenstecher, 1857) [Eriophyidae] Grape gall mite | Yes (Rather 1999) | Yes NSW, SA, Vic., WA (Plant Health Australia 2001) Qld, Tas. (CSIRO 2005) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|------------------------------|---|--|---|--|--------------------------------------|
| <i>Eotetranychus truncatus</i> Estebanes and Baker 1968 [Tetranychidae] Spider mite | Yes (Rather 1999) | No records found | No This mite is known to mainly feed on leaves, which lead to leaf yellowing and premature leaf fall (Gupta & Dhooria 1972; Rather 1999). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| <i>Eutetranychus orientalis</i> (Klein, 1936) [Tetranychidae] Oriental spider mite | Yes (Rather 1999) | Yes NSW, NT, Qld, WA (Plant Health Australia 2001) Qld (Walter, Halliday & Smith 1995) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Oligonychus coffeae</i> (Nietner, 1861) [Tetranychidae] Tea red spider mite | Yes (Dhooria 2003; DPP 2007) | Yes NSW, NT, Qld (Plant Health Australia 2001) WA (Poole 2010) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Oligonychus mangiferus</i> (Rahman & Sapra, 1940) [Tetranychidae] Mango red spider mite | Yes (Nassar & Ghai 1981) | Yes Qld (Plant Health Australia 2001) WA (Poole 2010) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|---------------------------|---|--|---|--|--------------------------------------|
| <i>Oligonychus vitis</i> (Zaher & Shehata, 1965) [Tetranychidae] Grape spider mite | Yes (Rather 1999) | No records found | No Primarily feeds on foliage and lays eggs on the bases of leaf buds or in scars in wood. Larvae move towards leaves and are found on upper and lower surfaces of leaves and shoots (Gonzalez 1983). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| <i>Panonychus citri</i> (McGregor 1916) [Tetranychidae] Citrus red mite | Yes (CABI 2012; DPP 2007) | Yes NSW (Plant Health Australia 2001) NSW, SA (CSIRO 2005) Not known to be present in WA (Poole 2010). | No Though this species attacks grapevine (Migeon & Dorkeld 2012; Wu & Lo 1989), feeding occurs on leaves (Jeppson, Keifer & Baker 1975). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |
| <i>Panonychus ulmi</i> (Koch, 1836) [Tetranychidae] European red spider mite | Yes (DPP 2007) | Yes NSW, SA, Tas., Vic., WA (Plant Health Australia 2001) WA (Botha & Learmonth 2005; Poole 2010) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Polyphagotarsonemus latus</i> (Banks, 1904) [Tarsonemidae] Broad mite | Yes (DPP 2007) | Yes NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|-------------------------|---|--|---|---|--------------------------------------|
| <i>Tetranychus kanzawai</i> Kishida, 1927 Synonym: <i>Tetranychus hydrangea</i> [Tetranychidae] Kanzawa spider mite | Yes (DPP 2007) | Yes Qld, NSW (CSIRO 2005; Gutierrez & Schicha 1983; Navajas et al. 2001). Not known to be present in WA (Poole 2010). | Yes <i>Tetranychus kanzawai</i> mites and webbing are often found on the under surfaces of the leaves, but can occasionally attack and breed on grape berries (Ashihara 1996; CABI 2012; Ho & Chen 1994). | Yes Major hosts are groundnut, tea, pawpaw, citrus, soybean, peach, apple, cherry, aubergine, watermelon and grapevine (CABI 2012; Migeon & Dorkeld 2012; Moon et al. 2008), which are present in Western Australia. This species is recorded from China, Greece, India, Japan, Korea and Mexico (Migeon & Dorkeld 2006). It has also been introduced to, and has successfully established in, Queensland and NSW (Gutierrez & Schicha 1983). Environments with climates similar to these regions exist in various parts of Western Australia, suggesting that <i>T. kanzawai</i> has the potential to establish and spread in WA. | Yes <i>Tetranychus kanzawai</i> is a significant polyphagous pest subject to quarantine measures in many parts of the world (Navajas et al. 2001). | Yes (EP, WA) |
| <i>Tetranychus neocaledonicus</i> (Andre, 1933) [Tetranychidae] Vegetable spider mite | Yes (Rather 1999) | Yes NSW, NT, Qld, WA (Plant Health Australia 2001) WA (Poole 2010) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|-------------------------|--|-----------------------------------|---|--|--------------------------------------|
| <i>Tetranychus urticae</i> Koch, 1836 Synonym: <i>Tetranychus cinnabarinus</i> (Boisduval, 1867) [Tetranychidae] Two-spotted spider mite | Yes (DPP 2007) | Yes NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|------------------|--|--|--|-------------------------------------|-------------------------------|
| BACTERIA | | | | | | |
| <p><i>Candidatus</i> Phytoplasma asteris [16SrI Aster yellows group] Aster yellows phytoplasma Note: Phytoplasmas classified in subgroups 16SrI-A, 16SrI-B and 16SrI-C ('<i>Candidatus</i> Phytoplasma asteris'-related strains) are associated with grapevine yellows in several countries (Alma et al. 1996; Bianco et al. 1994; Davis et al. 1998). The strains related to 'Ca. Phytoplasma asteris' comprises of a large number of related phytoplasmas worldwide, representing the most diverse and widespread phytoplasma group (Lee et al. 2004). Although there is relatively high similarity in the 16S rDNA sequence, the strains in this group occupy diverse ecological niches and show substantial genetic variation (Firrao, Gibbs & Streten 2005).</p> | Yes (DPP 2007) | <p>No records found Australia has <i>Candidatus</i> Phytoplasma australiense (Australian grapevine yellows) and <i>Candidatus</i> Phytoplasma asteris-related (Buckland Valley grapevine yellows) (Streten & Gibb 2006).</p> | <p>Yes Infects the phloem, causing grapevine yellows (Weintraub & Jones 2010). Several molecularly distinct phytoplasma groups that cause grapevine yellows have been identified (Hren et al. 2009).</p> | <p>No The Aster yellows group are not seed transmissible (CABI 2012; Lee, Davis & Gundersen-Rindal 2000). The Aster yellows group is graft transmissible, long-distance dissemination may occur through propagative material (Belli, Bianco & Conti 2010; CABI 2012). Aster yellows are also vector transmissible, the primary vectors are not found in Australia and are unlikely to be transmitted from the fruit bunch to a suitable host plant (ABRS 2009; CABI 2012; Krüger et al. 2011; Plant Health Australia 2001; Weintraub & Beanland 2006; Wilson & Turner 2010).</p> | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|-----------------------------------|--|--|--|--|--------------------------------------|
| <i>Candidatus</i> Phytoplasma asteris subgroup C [16SrI-C Aster yellows group] Clover phyllody phytoplasma | Yes (DPP 2007; Singh et al. 1983) | No records found Australia has <i>Candidatus</i> Phytoplasma australiense (Australian grapevine yellows) and <i>Candidatus</i> Phytoplasma asteris-related (Buckland Valley grapevine yellows) (Streten & Gibb 2006). | Yes Infects the phloem, causing grapevine yellows (Weintraub & Jones 2010). Several molecularly distinct phytoplasma groups that cause grapevine yellows have been identified (Hren et al. 2009). | No The Aster yellows group are not seed transmissible (CABI 2012; Lee, Davis & Gundersen-Rindal 2000). The Aster yellows group are graft transmissible, long-distance dissemination may occur through propagative material (CABI 2012; Lee, Davis & Gundersen-Rindal 2000). The Aster yellows group are also vector transmissible, the primary vectors are not found in Australia and are unlikely to be transmitted from the fruit bunch to a suitable host plant (ABRS 2009; CABI 2012; Plant Health Australia 2001; Weintraub & Beanland 2006; Wilson & Turner 2010). | Assessment not required | No |
| <i>Pantoea agglomerans</i> (Beijerinck 1888) Gavini et al. 1989 Synonym: <i>Erwinia herbicola</i> (Lohnis 1911) Dye 1964 [Enterobacteriales: Enterobacteriaceae] Bacterial grapevine blight | Yes (Pathak & Verma 2009) | Yes NSW, Vic., WA (Plant Health Australia 2001; Wilkinson et al. 1994) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|---|---|-----------------------------------|---|--|--------------------------------------|
| <i>Pseudomonas syringae</i> pv. <i>syringae</i> van Hall 1902 [Pseudomonadales: Pseudomonadaceae] Bacterial canker | Yes (CABI 2012) | Yes NSW, NT, Qld, SA, Tas., Vic., WA (Bradbury 1986; McCormick & Hollaway 1999; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Pseudomonas viridiflava</i> (Burkholder 1930) Dowson 1939 [Pseudomonadales: Pseudomonadaceae] Bacterial leaf blight of tomato | Yes (Shekhawat, Gadewar & Chakrabarti 1999) | Yes NSW, Qld, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Rhizobium radiobacter</i> (Beijerinck & van Delden 1902) Young <i>et al.</i> 2001 Synonym: <i>Agrobacterium tumefaciens</i> (Smith and Townsend 1907) Conn1942 [Rhizobiales: Rhizobaeaceae] Crown gall | Yes (CABI 2012) | Yes NSW, Qld, SA, Tas., Vic. (Plant Health Australia 2001). WA (Shivas 1989). | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Rhizobium vitis</i> (Ophel & Kerr 1990) Young <i>et al.</i> 2001 Synonym: <i>Agrobacterium vitis</i> (Ophel & Kerr 1990) [Rhizobiales: Rhizobaeaceae] Crown gall of grapevine | Yes (CABI 2012) | Yes NSW, SA, Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|--|---------------------------------|--|---|--|--------------------------------------|
| <i>Xanthomonas campestris</i> pv. <i>viticola</i> (Nayudu 1972) Dye 1978 [Xanthomondales: Xanthomonadaceae] Grapevine bacterial canker disease, Leaf spot | Yes (Chand et al. 1999; Jambenal 2008; Trindade, Lima & Ferreira 2005) | No records found | Yes Infects leaves, shoots and berries. Berries can develop brown and black lesions; infected berries are small and shrivelled (Chand & Kishun 1990). | Yes Host plants include mango (Trindade, Lima & Ferreira 2005) and grapevine (Chand et al. 1999) which are both widely grown in Australia. This pathogen has spread and established in Brazil where it has become the most important bacterial disease of grapevines in the Sao Francisco region (Nascimento & Mariano 2004). | Yes The first incidence of this disease in India was in 1969. However, it became a major problem when it caused yield losses of up to 80 per cent in 1986–87 (Chand et al. 1999). It is now a regular problem in the major grape growing regions of India (Chand et al. 1999; Jambenal 2008). | Yes |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|---|---|---|--|--|--------------------------------------|
| <i>Xylella fastidiosa</i> Wells <i>et al.</i> (1987) Synonym: <i>Xylella fastidiosa</i> subsp. <i>fastidiosa</i> Wells <i>et al.</i> (1987) [Xanthomonadales: Xanthomonadaceae] Pierce's disease | Yes An isolated record from almonds in India in 1985 exists (Jindal & Sharma 1987). However, the phony peach chemical test the authors used to confirm the identity of the pathogen was based on a 1970s method. It is recommended that this record be confirmed by more modern methods (CABI 2012). No other records from India exist. India stated in their supplementary submission that <i>X. fastidiosa</i> is not present on table grapes in India (DPP 2012). | No records found | Yes It spreads systemically through xylem vessels and can be present where ever these tissues occur (Pearson & Goheen 1988). | No <i>Xylella fastidiosa</i> has been subject to rigorous assessment in context with the review of policy for the glassy winged sharpshooter, a vector of <i>X. fastidiosa</i> , in 2002 (Biosecurity Australia 2002) and with significant trade of table grapes into eastern Australian states since that time. Should new information suggest there is a change in the risk profile of this disease and/or its vectors, this would initiate a further review process to ensure appropriate measures are in place to reduce the risks posed to meet Australia's appropriate level of protection. | Assessment not required | No |
| CHROMALVEOLATA | | | | | | |
| <i>Phytophthora drechsleri</i> Tucker [Peronosporales: Pythiaceae] Fruit rot | Yes (Farr & Rossman 2012) | Yes ACT, NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001; Shepherd & Pratt 1973) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|--|---|---|---|--|--------------------------------------|
| <i>Plasmopara viticola</i> (Berk. & M. A. Curtis) Berl. & De Toni [Peronosporales: Peronosporaceae] Grapevine downy mildew | Yes (CABI 2012; DPP 2007) | Yes ACT, NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| FUNGI | | | | | | |
| <i>Alternaria alternata</i> (Fr.) Keissl. [Pleosporales: Pleosporaceae] Alternaria leaf spot | Yes (CABI 2012; DPP 2007) | Yes ACT, NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Alternaria tenuissima</i> (Nees) Wiltshire [Pleosporales: Pleosporaceae] Alternaria leaf spot | Yes (Farr & Rossman 2012) | Yes NSW, Qld, SA, Tas., Vic., WA (Pethybridge, Hay & Groom 2003; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Alternaria vitis</i> Cavara [Pleosporales: Pleosporaceae] Grapevine alternariosis | Yes (Farr & Rossman 2012; Suhag, Kaushik & Duhan 1982) | No records found | No <i>Alternaria vitis</i> only infects leaves (Suhag, Kaushik & Duhan 1982). | Assessment not required | Assessment not required | No |
| <i>Aplosporella beaumontiana</i> S. Ahmad [Botryosphaeriales: Botryosphaeriaceae] Sooty mould | Yes (Prakash & Raof 1985; Rajak & Pandey 1985) | No records found | No <i>Aplosporella beaumontiana</i> only infects dry stems and leaves (Damm, Fourie & Crous 2007; Prakash & Raof 1985; Rajak & Pandey 1985; Sutton 1980). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|---|---|-----------------------------------|---|--|--------------------------------------|
| <i>Aspergillus fumigatus</i> Fresen. Synonym: <i>Neosartorya fumigata</i> O'Gorman, H.T. Fuller & P.S. Dyer [Eurotiales: Trichocomaceae] | Yes (Farr & Rossman 2012) | Yes NSW, NT, Qld, WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Aspergillus niger</i> Tiegh. [Eurotiales: Trichocomaceae] Aspergillus ear rot, Black mould | Yes (CABI 2012; DPP 2007; Singh & Chohan 1974) | Yes ACT, NSW, NT, Qld, SA, Vic., WA (Plant Health Australia 2001; Shivas 1989) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Aspergillus terreus</i> Thom [Eurotiales: Trichocomaceae] | Yes (Farr & Rossman 2012; Singh & Chohan 1974) | Yes NSW, Qld (Plant Health Australia 2001) WA (Kelly, Shaw & Clark 1995) Cosmopolitan distribution (Farr & Rossman 2012) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Aureobasidium pullulans</i> (de Bary) G. Arnaud [Dothideales: Dothioraceae] | Yes (Farr & Rossman 2012; Sarbhoy, Lal & Varshney 1975) | Yes ACT, NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Botryosphaeria dothidea</i> (Moug.: Fr.) Ces. & De Not. Synonym: <i>Fusicoccum aesculi</i> Corda [Botryosphaeriales: Botryosphaeriaceae] Macrophoma rot | Yes (CABI 2012) | Yes NSW, Qld, Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|--|---|--|---|--|--------------------------------------|
| <i>Botryosphaeria obtusa</i> (Schwein.) Shoemaker [Botryosphaeriales: Botryosphaeriaceae] Dead arm, Canker | Yes (Farr & Rossman 2012; Rajak & Pandey 1985) | Yes ACT, NSW, Qld, SA, Vic, WA (Cunnington et al. 2007; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Botryosphaeria ribis</i> Grossenb. & Duggar [Botryosphaeriales: Botryosphaeriaceae] | Yes (Farr & Rossman 2012) | Yes ACT, NSW, Qld, Vic, WA (Farr & Rossman 2012; Plant Health Australia 2001; Sakalidis, Hardy & Burgess 2011) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Botryosphaeria stevensii</i> Shoemaker Synonym: <i>Diplodia mutila</i> (Fr.: Fr.) Mont. [Botryosphaeriales: Botryosphaeriaceae] | Yes (Sharma & Bhardwaj 1999) | Yes ACT, NSW, Vic, WA (Farr & Rossman 2012; Plant Health Australia 2001; Taylor et al. 2005) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Botrytis cinerea</i> Pers.: Fr. Synonym: <i>Botryotinia fuckeliana</i> (de Bary) Whetzel [Helotiales: Sclerotiniaceae] Grey mould rot | Yes (CABI 2012; DPP 2007) | Yes ACT, NSW, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Cercospora truncata</i> Ellis & Everh. [Capnodiales: Mycosphaerellaceae] | Yes (Farr & Rossman 2012) | No records found | No <i>Cercospora truncata</i> only infects leaves (Farr & Rossman 2012; Robert, Stegehuis & Stalpers 2009). No records have been found which associate this species with grape bunches. | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|-----------------------------------|--|-----------------------------------|---|--|--------------------------------------|
| <i>Cladosporium cladosporioides</i> (Fresen.) G.A. de Vries 1952 [Capnodiales: Davidiellaceae] | Yes (Chakraborty et al. 2001) | Yes ACT, NSW, NT, Qld, SA, Tas., Vic., WA (Bensch et al. 2010; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Cladosporium herbarum</i> (Pers.: Fr.) Link Synonym: <i>Mycosphaerella tassiana</i> (De Not.) Johanson [Capnodiales: Davidiellaceae] Cladosporium rot | Yes (Farr & Rossman 2012) | Yes NSW, Qld, SA, Tas., Vic., WA (Maxwell & Scott 2008; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Colletotrichum acutatum</i> J.H. Simmonds [Phyllachorales: Phyllachoraceae] Strawberry black spot | Yes (DPP 2007; Kaur & Singh 1990) | Yes NSW, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Cylindrocarpon destructans</i> var. <i>destructans</i> (Zinssm.) Scholten Synonym: <i>Cylindrocarpon destructans</i> (Zinssm.) Scholten [Hypocreales: Nectriaceae] | Yes (Farr & Rossman 2012) | Yes ACT, NSW, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Elsinoë ampelina</i> Shear Synonym: <i>Sphaceloma ampelinum</i> de Bary [Myriangiales: Elsinoaceae] Grape anthracnose, Berry rot, Black spot, Bird's eye spot | Yes (CABI 2012; DPP 2007) | Yes NSW, NT, Qld, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|------------------------------------|---|-----------------------------------|---|--|--------------------------------------|
| <i>Epicoccum nigrum</i> Link [Dothideales: Dothioraceae] Cereal leaf spot | Yes (Farr & Rossman 2012) | Yes ACT, NSW, Qld, Vic., WA (Nair 1985; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Erysiphe necator</i> var. <i>necator</i> Schwein. Synonyms: <i>Erysiphe necator</i> Schwein., <i>Oidium tuckeri</i> Berk. [Erysiphales: Erysiphaceae] Grapevine powdery mildew | Yes (CABI 2012; DPP 2007) | Yes NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Fusarium oxysporum</i> Schltdl. [Hypocreales: Nectriaceae] Fusarium wilt | Yes (CABI 2012; DPP 2007) | Yes ACT, NSW, NT, Qld, SA, Tas., Vic., WA (Elmer et al. 1997; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Fusarium sacchari</i> (E.J. Butler) W. Gams Synonym: <i>Gibberella sacchari</i> Summerell & J.F. Leslie [Hypocreales: Nectriaceae] Fusarium wilt | Yes (DPP 2007; Leslie et al. 2005) | Yes (Summerell et al. 2011) NSW (CABI 2015) NT (Petrovic et al. 2013; Summerell et al. 2011) Qld (Summerell et al. 2011) WA (DAFWA 2015e) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|---|---|-----------------------------------|---|--|--------------------------------------|
| <i>Fusarium solani</i> (Mart.) Sacc. [Hypocreales: Nectriaceae] Dry rot | Yes (Farr & Rossman 2012; Sharma, Kaushik & Kaushik 1997) | Yes NSW, NT, Qld, SA, Tas., Vic., WA (Elmer et al. 1997; Pung & Cox 1999; Sangalang et al. 1995) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Fusarium subglutinans</i> (Wollenw. & Reinking) P.E. Nelson, Toussoun & Marasas [Hypocreales: Nectriaceae] Damping off, Pitch canker | Yes (Farr & Rossman 2012; Rawal 1998) | Yes NSW, NT, Qld, Vic., WA (Elmer et al. 1997; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Fuscoporia gilva</i> (Schwein.) T. Wagner & M. Fisch. Synonym: <i>Phellinus gilvus</i> (Schwein.) Pat. [Hymenochaetales: Hymenochaetaceae] | Yes (Farr & Rossman 2012) | Yes Known as <i>Phellinus gilvus</i> in NSW, NT, Qld, Vic., WA (Plant Health Australia 2001). | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Geotrichum candidum</i> Link [Saccharomycetales: Dipodascaceae] Fruit rot | Yes (Badyal & Sumbali 1990; Farr & Rossman 2012) | Yes NSW, NT, Qld, Tas., Vic., WA (Plant Health Australia 2001; Wade & Morris 1982) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Glomerella cingulata</i> (Stoneman.) Spauld. & H. Schrenk Synonym: <i>Colletotrichum gloeosporioides</i> (Penz.) Penz. & Sacc. [Phyllachorales: Glomerellaceae] Anthracnose | Yes (Farr & Rossman 2012; Kaur & Singh 1990) | Yes ACT, NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001). Tas. (Sampson & Walker 1982) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|--------------------------|---|---|---|---|--------------------------------------|
| <i>Greeneria uvicola</i> (Berkley & M.A. Curtis) Punithalingam [Diaporthales: Gnomoniaceae] Bitter rot | Yes (Reddy & Reddy 1983) | Yes NSW, Qld (Plant Health Australia 2001) Not known to be present in WA. | Yes Bitter rot can affect young shoots, stems of fruit bunches, pedicels and berries. <i>Greeneria uvicola</i> usually attacks berries via the pedicel. Within a few days of infection, berries soften and are bitter to taste; some are easily detached while others shrivel and mummify (McGrew 1988; Momol, Ritchie & Dankers 2007). | Yes In Australia, <i>G. uvicola</i> has been reported from north eastern New South Wales and Queensland (Plant Health Australia 2001; Sergeeva, Nair & Spooner-Hart 2001; Steel, Greer & Savocchia 2007). <i>Greeneria uvicola</i> has also been reported from Brazil, Costa Rica, Greece, India, Japan, New Zealand, South Africa and the USA (Kummuang et al. 1996b; McGrew 1988; Steel 2007; Sutton & Gibson 1977; Ullasa & Rawal 1986). Environments with climates similar to these regions exist in various parts of Western Australia suggesting that <i>G. uvicola</i> has the potential to establish and spread in Western Australia. Hosts are <i>Vitis</i> spp (Farr et al. 2001; Sutton & Gibson 1977), which are widely grown in Australia. | Yes <i>Greeneria uvicola</i> attacks many species of grape, including <i>Vitis vinifera</i> (European grape), <i>V. labrusca</i> (fox grape) and <i>V. rotundifolia</i> (muscadine grape) (Farr et al. 2001; Sutton & Gibson 1977). Affected berries shrivel and rot or become soft, bitter-tasting and are easily detached (McGrew 1988). <i>Greeneria uvicola</i> can also cause girdling of the shoots of <i>V. vinifera</i> cultivars (McGrew 1988). <i>Greeneria uvicola</i> has also been reported to cause rot on mature fruit of apple, cherry, strawberry, peach and banana under experimental conditions (Ridings & Clayton 1970). | Yes (EP, WA) |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|---|---|--|--|--|--------------------------------------|
| <p><i>Guignardia bidwellii</i> (Ellis) Viala & Ravez Synonym: <i>Phyllosticta ampellicida</i> (Engelm.) Van der Aa [Botryosphaerales: Botryosphaeriaceae] Black rot</p> | Yes (CABI 2012; DPP 2007; Farr & Rossman 2012; Singh et al. 1999) | No records found | Yes Affects grape leaf, stem, peduncle and fruit (CABI 2012; NPQS 2007; Ramsdell & Milholland 1988). The pathogen attacks all parts of the vine, particularly the berry clusters (Singh et al. 1999). | Yes <i>Guignardia bidwellii</i> overwinters in mummified berries, either in the vine or on the ground. Ascospores are airborne and disperse moderate distances and conidia are splash dispersed only short distances (Wilcox 2003). <i>Guignardia bidwellii</i> has a range of hosts, including <i>Ampelopsis</i> spp., <i>Cissus</i> spp., <i>Citrus</i> spp., <i>Vitis</i> spp., <i>Arachis hypogaea</i> (peanut) and <i>Asplenium nidus</i> (bird's nest fern), which are widely distributed in home gardens, nurseries and orchards in Australia (Eyres, Wood & Taylor 2006; Farr & Rossman 2012). | Yes Black rot is an important fungal disease of grapes that originated in eastern North America, but now occurs in parts of Europe, South America and Asia (Wilcox 2003). Crop losses can range from 5 to 80 per cent (Ramsdell & Milholland 1988) and are depending on weather, inoculum levels and cultivar susceptibility. | Yes (EP) |
| <p><i>Lasiodiplodia theobromae</i> (Pat.) Griffon & Maubl. Synonyms: <i>Botryodiplodia theobromae</i> Pat., <i>Botryosphaeria rhodina</i> (Berk. & Curtis) Arx [Botryosphaerales: Botryosphaeriaceae] Lasiodiplodia cane dieback</p> | Yes (CABI 2012; DPP 2007; Farr & Rossman 2012) | Yes NSW, NT, Qld, SA, WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|--------------------------------------|--|-----------------------------------|---|--|--------------------------------------|
| <i>Macrophomina phaseolina</i> (Tassi) Goid. [Sphaeropsidales: Sphaeropsidaceae] Charcoal rot | Yes (CABI 2012; Farr & Rossman 2012) | Yes ACT, NSW, NT, Qld, SA, Vic., WA (Plant Health Australia 2001; Sergeeva et al. 2005) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Monilinia fructicola</i> (G. Winter) Honey Synonym: <i>Monilia fructicola</i> L. R. Batra [Helotiales: Sclerotiniaceae] Brown rot | Yes (CABI 2012) | Yes ACT, NSW, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|---|--------------------------|---|---|---|-------------------------------|
| <p><i>Monilinia fructigena</i> (Aderh. & Ruhland) Honey Synonym: <i>Monilia fructigena</i> (Pers.) Pers. [Helotiales: Sclerotiniaceae] Brown rot</p> | Yes (CABI 2012; DPP 2007; Sharma & Kaul 1989) | No records found | <p>Yes</p> <p>Causes raised light brown pustules on the fruit that often expand enclosing the fruit to form a dark, wrinkled, hard mummified fruit (USDA-APHIS 2004). Grape is not a primary host.</p> <p>The original record of <i>M. fructigena</i> on grape was in China (Qi <i>et al.</i> 1966 in Tai (1979)), which provided evidence of the anamorphic stage (<i>Monilia fructigena</i>) being associated with <i>Vitis vinifera</i>. This pathogen has also been reported to cause a soft brown rot of grape berries in both Italy and Japan (Nanni, Zaccaria & Marziano 2003; Ogata, Katsumata & Takaiwa 1999).</p> | <p>Yes</p> <p>Brown rot disease caused by <i>M. fructigena</i> is a common and widespread disease of pome and stone fruit (Mackie, Eyres & Kumar 2005), which are grown widely in Australia.</p> <p>The spores of this fungus can be spread from one orchard to another through wind and water (Jones 1990), as well as potentially being transported by various insects (CABI 2012).</p> | <p>Yes</p> <p><i>Monilinia fructigena</i> produces raised light brown pustules that enclose the fruit to form a wrinkled and mummified fruit (USDA-APHIS 2004).</p> <p><i>Monilinia fructigena</i> causes brown rot disease in pome and stone fruit which is the soft decay of fruit flesh and blighting of spurs and blossoms (Mackie, Eyres & Kumar 2005). This results in significant pre- and post-harvest fruit losses and causes considerable economic losses worldwide (Jones 1990; Mackie, Eyres & Kumar 2005).</p> <p>Brown rot is responsible for great losses to apple after harvest in Himachal Pradesh (Sharma & Kaul 1989).</p> <p>Extensive surveys between 1982–1985 revealed cumulative incidences of the disease varied from 2.0 to 14 per cent (Sharma & Kaul 1989).</p> | Yes (EP) |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|---|---|-----------------------------------|---|--|--------------------------------------|
| <i>Mucor circinelloides</i> Tiegh. [Mucorales: Mucoraceae] | Yes (DPP 2007) | Yes NSW, Qld, Vic., WA (Kew Royal Botanic Gardens 2014; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Mucor racemosus</i> Fresen. Synonym: <i>Mucor varians</i> Povah [Mucorales: Mucoraceae] Spongy storage rot | Yes (Farr & Rossman 2012) | Yes ACT, NSW, Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Neofusicoccum mangiferae</i> (Syd. & P. Syd.) Crous Synonyms: <i>Nattrassia mangiferae</i> (Syd. & P. Syd.) B. Sutton & Dyko; <i>Fusicoccum mangiferae</i> (Syd. & P. Syd.) G.I. Johnson, Slippers & M.J. Wingf. [as 'mangiferum'] [Botryosphaerales: Botryosphaeriaceae] Leaf spot, Stem end rot | Yes (DPP 2007; Farr & Rossman 2012) | Yes Qld (Plant Health Australia 2001) In WA as <i>Nattrassia mangiferae</i> or <i>Fusicoccum mangiferum</i> and in Qld as <i>Fusicoccum mangiferum</i> (Plant Health Australia 2001). | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Neoscytalidium dimidiatum</i> (Penz.) Crous & Slippers Synonyms: <i>Hendersonula toruloidea</i> Natrass; <i>Scytaalidium dimidiatum</i> (Penz.) B. Sutton & Dyko [Botryosphaerales: Botryosphaeriaceae] | Yes (Farr & Rossman 2012; Wangikar, Raut & Gopalkrishna 1969) | Yes NT, WA (Plant Health Australia 2001) Qld as <i>Torula dimidiata</i> (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|-------------------------------------|---|---|---|--|--------------------------------------|
| <i>Passalora dissiliens</i> (Duby) U. Braun & Crous Synonym: <i>Phaeoramularia dissiliens</i> (Duby) Deighton [Capnodiales: Mycosphaerellaceae] | Yes (Farr & Rossman 2012) | Vic. (Washington & Nancarrow 1983) Not known to be present in WA. | No <i>Passalora dissiliens</i> causes variable leaf spot symptoms (Deighton 1976). It has been recorded on leaves of <i>Vitis vinifera</i> in Victoria (Washington & Nancarrow 1983). No report of association with grape bunches was found. | Assessment not required | Assessment not required | No |
| <i>Penicillium aurantiogriseum</i> Dierckx [Eurotiales: Trichocomaceae] Blue mould rot | Yes (Palejwala, Sattur & Modi 1988) | Yes NSW, Qld, Vic., WA (Kew Royal Botanic Gardens 2014; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Penicillium chrysogenum</i> Thom [Eurotiales: Trichocomaceae] | Yes (DPP 2007; Farr & Rossman 2012) | Yes ACT, NSW, Qld, Tas., Vic. (Plant Health Australia 2001) WA (Kew Royal Botanic Gardens 2014) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Penicillium digitatum</i> (Pers.) Sacc. [Eurotiales: Trichocomaceae] Green mould | Yes (CABI 2012) | Yes NSW, Qld, SA, Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|--------------------------------------|--|---|---|--|--------------------------------------|
| <i>Penicillium expansum</i> Link [Eurotiales: Trichocomaceae] Blue mould of stored apple | Yes (CABI 2012; DPP 2007) | Yes NSW, Qld, SA, Vic., WA (CABI 2012; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Penicillium italicum</i> Wehmer [Eurotiales: Trichocomaceae] Blue mould | Yes (CABI 2012) | Yes ACT, NSW, Qld, SA, Vic., WA (CABI 2012; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Pestalotiopsis funerea</i> (Desm.) Steyaert [Xylariales: Amphisphaeriaceae] Leaf spot | Yes (Farr & Rossman 2012) | Yes NSW, Qld, Vic. (Plant Health Australia 2001). Not known to be present in WA | No Affects leaves, stems and roots of its hosts (Mordue 1976). No report of association with grape bunches was found. | Assessment not required | Assessment not required | No |
| <i>Pestalotiopsis mangiferae</i> (Henn.) Steyaert [Xylariales: Amphisphaeriaceae] Grey leaf spot of mango | Yes (DPP 2007; Verma et al. 1991) | Yes NT, WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|------------------------------------|---|--|--|--|--------------------------------------|
| <i>Pestalotiopsis menezesiana</i> (Bres. & Torr.) Bissett [Xylariales: Amphisphaeriaceae] Fruit rot | Yes (Mishra, Prakash & Misra 1974) | Yes NSW (Plant Health Australia 2001; Sergeeva, Priest & Nair 2005) Not known to be present in WA | Yes Infects fruit (Mishra, Prakash & Misra 1974; Xu et al. 1999). | Yes <i>Pestalotiopsis menezesiana</i> infects <i>Cissus rhombifolia</i> (Bissett 1982), grapevine (Mishra, Prakash & Misra 1974; Xu et al. 1999), kiwifruit (Park et al. 1997) and plantain (Huang et al. 2007), which are present in Western Australia. <i>Pestalotiopsis menezesiana</i> is present in Australia (NSW), China, India, Japan and Korea (Huang et al. 2007; Mishra, Prakash & Misra 1974; Park et al. 1997; Plant Health Australia 2001; Xu et al. 1999). Environments with climates similar to these regions exist in various parts of Western Australia. | Yes Grapevine, kiwifruit and plantain are commercially grown in Western Australia. This pathogen causes rot of grape berries (Mishra, Prakash & Misra 1974; Xu et al. 1999), leaf spot of kiwifruit (Park et al. 1997) and leaf spot of plantain (Huang et al. 2007). | Yes (EP, WA) |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|--------------------------|--|--|--|--|--------------------------------------|
| <i>Pestalotiopsis uvicola</i> (Speg.) Bissett [Xylariales: Amphisphaeriaceae] Fruit rot, Berry rot, Leaf spot | Yes (Mohan et al. 2005) | Yes NSW, Qld (Plant Health Australia 2001) NSW (Sergeeva, Priest & Nair 2005) Not regulated for Tas. (DPIPWE 2011). Not known to be present in WA. | Yes Affects grape berries (Guba 1961; Sergeeva, Priest & Nair 2005; Xu et al. 1999). | Yes In Australia, <i>P. uvicola</i> has been reported from NSW and Qld (Plant Health Australia 2001). This pathogen has also been reported from Brazil, France, Italy and the US (Guba 1961). Environments with climates similar to these regions exist in various parts of WA suggesting that <i>P. uvicola</i> has the potential to establish and spread in WA. Hosts include <i>Vitis vinifera</i> , <i>Laurus nobilis</i> and <i>Mangifera indica</i> (Ismail, Cirvilleri & Polizzi 2013; Vitale & Polizzi 2005; Xu et al. 1999), which are grown in WA. | Yes Has been reported to cause post-harvest disease of grapes (Xu et al. 1999), leaf spot and stem blight of bay laurel (<i>Laurus nobilis</i>) (Vitale & Polizzi 2005) and leaf spot of mango (Ismail, Cirvilleri & Polizzi 2013). | Yes (EP, WA) |
| <i>Phakopsora ampelopsidis</i> Diet. & P. Syd. [Pucciniales: Phakopsoraceae] Ampelopsis rust fungus | Yes (Punithalingam 1968) | No records found | No Revised distribution by Ono (2000) did not place this disease in India. This species does not infect <i>Vitis</i> spp. (Ono 2000). | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|---|---|---|---|--|--------------------------------------|
| <i>Phakopsora euvitis</i> Y. Ono [Pucciniales: Phakopsoraceae] Grapevine leaf rust, grapevine rust, grapevine rust fungus | Yes (CABI 2012; DPP 2007; Leu 1988; Persley & Magarey 2009). | No Recorded in NT (Weinert et al. 2003) but has since been eradicated (EPPO 2007; IPPC 2008; Persley & Magarey 2009). | Yes Infects leaves of <i>Vitis vinifera</i> (CABI 2012) and young shoots (Li 2004). Occasionally infects rachises (Leu 1988). | Yes <i>Phakopsora euvitis</i> established in the Northern Territory before eradication (Weinert et al. 2003). Rust fungi spores are wind dispersed (Deacon 2005), and are produced abundantly in warm and humid weather (Persley & Magarey 2009). Hosts are <i>Vitis</i> spp. (Weinert et al. 2003), which are widely grown in Australia. | Yes Rust disease caused by <i>P. euvitis</i> is very destructive (Leu 1988). Heavy infection causes early senescence of the leaves and premature leaf fall. The disease can cause poor shoot growth, reduction of fruit quality and yield loss (CABI 2012). | Yes (EP) |
| <i>Phoma betae</i> A.B. Frank Synonym: <i>Pleospora betae</i> (Berl.) Nevod. F [Pleosporales: Pleosporaceae] | Yes (Farr & Rossmann 2012) | Yes NSW, Qld, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Phoma glomerata</i> (Corda) Wollenw. & Hochapfel [Pleosporales: Pleosporaceae] Phoma blight | Yes (Farr & Rossmann 2012; Pandotra 1976) | Yes NSW, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|---|--|---|---|--|--------------------------------------|
| <p><i>Phomopsis viticola</i> (Sacc.) Sacc. Synonyms: <i>Cryptosporella viticola</i> (Reddick) Shear; <i>Fusicoccum viticola</i> Reddick [Diaporthales: Valsaceae] Phomopsis cane and leaf spot, Excoriose (Europe), Dead arm (USA)</p> | Yes (CABI 2012; DPP 2007; Farr & Rossman 2012; Lal & Arya 1982) | Yes (Merrin, Nair & Tarran 1995) NSW, Qld, SA, Vic. (Plant Health Australia 2001). Tas. (Mostert, Corus & Kang 2001). Plant Health Australia (2001) also shows records for WA, but these have been shown to be <i>Diaporthe australafricana</i> or other species of <i>Phomopsis</i> other than <i>P. viticola</i> by sequencing of the ITS region. | Yes Infects all parts of the grape bunch including rachis, pedicels and berries (Hewitt & Pearson 1988; Persley & Magarey 2009). | Yes <i>Phomopsis viticola</i> is established in temperate climatic regions throughout the viticultural world and has been reported in Africa, Asia, Australia (except WA), Europe and North America (Hewitt & Pearson 1988). Spores of <i>P. viticola</i> are dispersed by rain splash and insects within the vineyard. Long distance dispersal occurs by movement of infected/contaminated propagation material, pruning equipment and agricultural machinery (Burges, Taylor & Kumar 2005). | Yes <i>Phomopsis viticola</i> is a serious pathogen of grapes in several viticultural regions of the world (Hewitt & Pearson 1988). It can cause vine stunting and reduced fruit yield (Burges, Taylor & Kumar 2005), as well as lower the quality of fruit and kill grafted and other nursery stock (Hewitt & Pearson 1988). | Yes (EP, WA) |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|-------------------------|--|---|--|--|--------------------------------------|
| <p><i>Pilidiella castaneicola</i> (Ellis & Everh.) B. Sutton</p> <p>Synonyms: <i>Coniella castaneicola</i> (Ellis & Everh.) B. Sutton, <i>Schizoparme straminea</i> Shear</p> <p>[Diaporthales: Schizoparmaceae]</p> <p>White rot</p> | Yes (Nag Raj 1993) | <p>Yes</p> <p>NSW, NT, Qld, Vic., but not on grape (Plant Health Australia 2001)</p> <p>On leaf of <i>Eucalyptus pellita</i> in Qld (Langrell, Glen & Alfenas 2008)</p> <p>Not regulated for Tas. (DPIPWE 2011).</p> <p>Not known to be present in WA.</p> | <p>Yes</p> <p>Causes white rot of table grapes. It affects rachis, pedicel and berries (Kishi 1998; Yamato 1995).</p> | <p>Yes</p> <p>This fungus has a variety of hosts (Farr & Rossman 2012). Hosts, including grapevine, are widely grown in Western Australia.</p> <p>In Australia, <i>P. castaneicola</i> has been reported from NSW, NT, Qld, Vic., (Plant Health Australia 2001).</p> <p><i>Pilidiella castaneicola</i> has also been reported from Brazil, Canada, China, Cuba, India, Indonesia, Japan, Korea, Pakistan, South Africa, Switzerland, the US and the West Indies (Farr & Rossman 2012).</p> <p>Environments with climates similar to these regions exist in various parts of WA suggesting that <i>P. castaneicola</i> has the potential to establish and spread in WA.</p> | <p>Yes</p> <p>Causes white rot of grapevine berries, (Kishi 1998; Yamato 1995) reducing marketability.</p> <p>Causes fruit rot of strawberries and is found on foliage of broadleaved trees (Farr & Rossman 2012).</p> <p>Is commonly found on leaves of <i>Eucalyptus</i>, but is of minor importance as a leaf pathogen (Van Niekerk et al. 2004).</p> | Yes (EP, WA) |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|---------------------------|--|---|---|--|--------------------------------------|
| <i>Pilidiella diplodiella</i> (Speg.) Crous & Van Niekerk Synonym: <i>Coniella diplodiella</i> (Speg.) Petr. & Syd [Diaporthales: Melanconidaceae] White rot, grapevine white rot | Yes (CABI 2012; DPP 2007) | Yes (Van Niekerk et al. 2004) NSW and WA-as <i>Coniella diplodiella</i> (Plant Health Australia 2001) WA (Shivas 1989). However, the WA isolates were recently re-identified as <i>Coniella fragariae</i> . | Yes Infects young and mature fruit, causing purple-brown spots, yellowing and then browning and drying out of the fruit (Lauber & Schuepp 1968). | Yes Hosts of <i>P. diplodiella</i> , <i>Vitis</i> spp. (Farr & Rossman 2012), are cultivated in Western Australia. | Yes <i>Pilidiella diplodiella</i> causes white rot of grapevine berries, reducing marketability (Bisiach 1988; Van Niekerk et al. 2004). It can also cause cankers in nonlignified shoots of grapevine (Bisiach 1988). | Yes (EP, WA) |
| <i>Pseudocercospora vitis</i> (Lév.) Speg. Synonym: <i>Mycosphaerella personata</i> B.B. Higgins [Capnodiales: Mycosphaerellaceae] Grapevine leaf spot, Leaf blight | Yes (Pons & Sutton 1988) | Yes Qld, NSW, Vic. (Plant Health Australia 2001) Not known to be present in WA. | No Infects leaves (McGrew & Pollack 1988). No report of association with grape bunches was found. | Assessment not required | Assessment not required | No |
| <i>Rhizoctonia solani</i> J.G. Kühn Synonym: <i>Thanatephorus cucumeris</i> (A.B. Frank) Donk [Ceratobasidiales: Ceratobasidiaceae] Damping off | Yes (CABI 2012) | Yes ACT, NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Rhizopus arrhizus</i> A. Fischer [Mucorales: Mucoraceae] Fruit rot | Yes (Farr & Rossman 2012) | Yes NSW, Vic. (Plant Health Australia 2001) WA (Kew Royal Botanic Gardens 2014) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|---|--|-----------------------------------|---|--|--------------------------------------|
| <i>Rhizopus stolonifer</i> (Ehrenb.) Vuill. [Mucorales: Mucoraceae] Rhizopus rot | Yes (DPP 2007) | Yes NSW, NT, Qld, Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Rosellinia necatrix</i> Prill. Synonym: <i>Dematophora necatrix</i> R. Hartig [Xylariales: Xylariaceae] White root rot | Yes (CABI 2012) | Yes NSW, Qld, WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Schizophyllum commune</i> Fr. [Agaricales: Agaricomycetidaeeae] Schizophyllum rot | Yes (Swapna, Syed & Krishnappa 2008) | Yes NSW, NT, Qld, SA, Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Sclerotinia sclerotiorum</i> (Lib.) de Bary Synonym: <i>Sclerotium varium</i> Pers. [Helotiales: Sclerotiniaceae] White mould | Yes (Farr & Rossman 2012) | Yes ACT, NSW, Qld, SA, Tas., Vic., WA (Hall, McMahon & Wicks 2002; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Sclerotium rolfsii</i> Sacc. Synonym: <i>Corticium rolfsii</i> Curzi, <i>Athelia rolfsii</i> (Curzi) C.C. Tu & Kimbr. [Poriales: Atheliaceae] Sclerotium stem rot | Yes (CABI 2012; Farr & Rossman 2012) | Yes ACT, NSW, NT, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001; Vawdrey & Peterson 1990) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|---|--|-----------------------------------|---|--|--------------------------------------|
| <i>Stemphylium botryosum</i> Wallr. Synonym: <i>Pleospora tarda</i> E. G. Simmons [Pleosporales: Pleosporaceae] Stemphylium rot | Yes (Ihsanul Huq & Nowsher Ali Khan 2008) | Yes NSW, SA, Tas., Vic., WA (Barbetti et al. 2006; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Trichoderma harzianum</i> Rifai Synonym: <i>Hypocrea lixii</i> Pat. [Hypocreales: Hypocreaceae] | Yes (DPP 2007) | Yes NSW, NT, Qld, SA, Tas., Vic., WA (Penrose, Nicholls & Koffmann 1984; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Trichothecium roseum</i> (Pers.) Link [Hypocreales: Not Assigned] Pink mould rot | Yes (DPP 2007; Sharma & Agarwal 1997) | Yes ACT, NSW, Qld, SA, Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Verticillium dahliae</i> Kleb. [Not Assigned: Plectosphaerellaceae] Verticillium wilt | Yes (DPP 2007) | Yes ACT, NSW, Qld, SA, Tas., Vic., WA (Harding & Wicks 2007; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| VIRUSES | | | | | | |
| <i>Alfalfa mosaic virus</i> [Bromoviridae: Alfamovirus] | Yes (CABI 2012; Nain, Rishi & Bishnoi 1994) | Yes NSW, Qld, SA, Tas., Vic., WA (CABI 2012; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|---|---|--|--|--|--------------------------------------|
| <i>Broad bean wilt virus 2</i> (BBWV 2) [Secoviridae: Fabavirus] | Yes (CABI 2012; Mali, Vyanjane & Ekbote 1977) | Yes NSW (Schwinghamer et al. 2007). May be present in Qld (Plant Health Australia 2001) but the records could be of <i>Broad bean wilt virus 1</i> . Not known to be present in WA. | Yes Recorded in grapevine (CIHEAM 2006). Probably infects systemically. | No At least one strain is transmitted in seed of <i>Vicia faba</i> , broad bean (Zhou 2002), but no record of seed transmission in <i>Vitis</i> spp. was found. Transmitted in a non-persistent manner by aphids, including <i>Myzus persicae</i> , <i>Aphis craccivora</i> and <i>Acyrtosiphon pisum</i> (Zhou 2002). No records of acquisition of the virus from infected berries. | Assessment not required | No |
| <i>Cucumber mosaic virus</i> [Bromoviridae: Cucumovirus] | Yes (Rishi 2009; Samad et al. 2008) | Yes NSW, Qld, SA, Tas., Vic., WA (Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |
| <i>Peach rosette mosaic virus</i> [Secoviridae: Nepovirus] | No Only one record from India in 1986 exists (CABI 2012). This record is considered unreliable (EPPO 2011). DPP (2012) states that this virus is not present in India. | Assessment not required | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|--|---|--|--|--|--------------------------------------|
| <p><i>Strawberry latent ringspot virus</i> (SLRSV)</p> <p>Synonyms: Aesculus line pattern virus (Schmelzer and Schmidt, 1968); Rhubarb virus 5 [Secoviridae: unassigned]</p> | <p>Yes</p> <p>(CABI 2016; Kulshrestha et al. 2004)</p> | <p>No</p> <p>Recorded in SA on rhubarb (Cook & Dubé 1989) but there are no further records, and the department considers the virus to be absent from Australia.</p> | <p>Yes</p> <p>No specific records for SLRSV in grapevine in India but it is known to occur in <i>Lilium</i> and roses in India (Adekunle et al. 2006; Kulshrestha et al. 2004). However, SLRSV has been associated with grapevines internationally, including from the Czech Republic (Komínek 2008), Italy (Credi et al. 1981), Romania (Eppler, Lesan & Lazar 1989) and Turkey (Akbas & Erdiller 1993).</p> <p>SLRSV infects plants systemically (Murant 1974).</p> <p>Although not reported in Indian viticulture, its presence in India and known association with grapevine internationally suggests that SLRV could potentially be associated with the table grape export pathway.</p> | <p>No</p> <p>This virus has been demonstrated to be seed transmitted in some hosts, including celery, quinoa, raspberry and some weeds (Murant 1983). However, seed transmission has not been recorded in grapevine.</p> <p>Spread occurs via its root-feeding nematode vectors <i>Xiphinema diversicaudatum</i> and <i>X. coxi</i> (CABI 2014). Both nematodes are absent from Australia.</p> <p>Can be transmitted by grafting (Brunt et al. 1996) but rachis material is not suitable for grafting.</p> | <p>Assessment not required</p> | <p>No</p> |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|--|---|--|--|---|--------------------------------------|
| <i>Tobacco necrosis viruses</i> (TNV) [Tombusviridae: Necrovirus] | Yes TNV-D (cb isolate) (CABI 2012; DPP 2012; Ramachandraiah, Venkatarathnam & Sulochana 1979) | Yes TNVs have been recorded in Vic. and Qld (Finlay & Teakle 1969; Teakle 1988), but not on grapevine. It is not known if the species or strain that infects grapevine is present in Australia. Not known to be present in WA. | Yes The strain of <i>Tobacco necrosis virus</i> found in grapevine in South Africa spreads systemically (Cesati & Van Regenmortel 1969); probably present in grape bunches. | Yes TNV strains are established in Australia (Teakle 1988). TNV strains typically have a wide host range (Uyemoto 1981), including grapevine (Zitikaite & Staniulis 2009) and many of these hosts occur in Australia. TNVs are transmitted by <i>Olpidium</i> spp. (Rochon et al. 2004) and at least one of these vectors occurs in Australia (Maccarone et al. 2008). | Yes TNVs cause rusty root disease of carrot, Augusta disease of tulip, stipple streak disease of common bean, necrosis disease of cabbage, cucumber, soybean and zucchini and ABC disease of potato (Smith et al. 1988; Uyemoto 1981; Xi et al. 2008; Zitikaite & Staniulis 2009). | Yes (EP) |
| <i>Tobacco ringspot virus</i> [Secoviridae: Nepovirus] | Yes (CABI 2012; Madhusudan & Govindu 1985) | Yes Qld, SA (Büchen-Osmond et al. 1988). Not considered to be present in WA (DAFWA 2013). However, this virus may be seed-borne in capsicum seed (Stace-Smith 1985) for planting that is permitted entry into Western Australia from Australian states where this virus is present. | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|--|---------------------------------|--|--|---|--------------------------------------|
| <i>Tomato black ring virus</i> (TBRV) [Secoviridae: Nepovirus] | Yes (CABI 2012; DPP 2007; Madhusudan & Govindu 1985) | No records found | Yes <i>Vitis</i> spp. is a principal host (Card, Pearson & Clover 2007). Many plant species infected with TBRV are symptomless, and are difficult to detect (CABI 2012; Murrant 1983). However, some symptoms include necrotic rings and malformation (Harrison 1996). It has also been demonstrated to transmit through seed in at least 25 plant families (Murrant 1983). | Yes TBRV has been recorded throughout Europe, Africa, Asia and the Americas (Harrison 1996). It is known to infect over 76 experimental plant species, including horticultural, agricultural and many weed and endemic species (CABI 2012; Harrison 1996), causing various levels of disease. Many of these plants are present in Australia. It is transmitted in nature by <i>Longidorus</i> spp. of nematodes (Brown, Murrant & Trudgill 1989; Harrison 1996). <i>Longidorus</i> spp. have been reported throughout Australia (Harris 1983; Plant Health Australia 2001). The virus can also be transmitted through sap extracts (Madhusudan & Govindu 1985). It is believed that nearly all of the nematode borne viruses, such as TBRV, can be transmitted and distributed through the seed of their principal hosts (Murrant 1983). | Yes TBRV causes necrotic rings, spots and flecks, mottle stunting and leaf malformation (Harrison 1996). The experimental host range is >9 families susceptible to the virus, many of which include important commodities such as onion, lettuce, tomato, potato, tulip, cucumber, strawberry and grapevines (CABI 2012; Harrison 1996). | Yes |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|--|---|--|--|--|--------------------------------------|
| <i>Tomato ringspot virus</i> [Secoviridae: Nepovirus] | Yes (Rana et al. 2011; Verma, Khan & Handa 2003) | No Recorded in SA (Chu, Francki & Hatta 1983; Cook & Dubé 1989), but there are no further records, the infected plants no longer exist, and the virus is believed to be absent from Australia. | Yes Infects systemically; present in fruit and seed (Gonsalves 1988; Uyemoto 1975). | No Seed transmitted in grapevine occasionally (Uyemoto 1975). Also transmitted by nematodes (<i>Xiphinema</i> spp.) and by grafting (Stace-Smith 1984). Transmission via nematode from fruit for human consumption is unlikely. Infected grapevine seedlings are very unlikely to establish, as demonstrated by the full assessment of the likelihood of establishment for <i>Tomato black ring virus</i> in this report. The chance that infected grape seeds from fruit waste will germinate is small. If germination does occur, seedlings are unlikely to survive. | Assessment not required | No |
| <i>Tomato spotted wilt virus</i> [Bunyaviridae: Tospovirus] | Yes (CABI 2012) | Yes NSW, NT, Qld, SA, Tas., Vic., WA (Persley, Thomas & Sharman 2006; Plant Health Australia 2001) | Assessment not required | Assessment not required | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|---|--------------------------------|--|--|---|--|--------------------------------------|
| VIROIDS | | | | | | |
| <i>Citrus exocortis viroid</i> (CEVd) [Pospiviroidae: Pospiviroid] | Yes (Ramachandran et al. 1993) | Yes Only known to be present in NSW, Qld and SA (Barkley & Büchen-Osmond 1988). Not known to be present in WA. | Yes Grapevine is a host of CEVd (Garcia-Arenal, Pallas & Flores 1987) and transmission of the viroid via grape seed has been observed (Wan Chow Wah & Symons 1997). | No The viroid may be transmitted by grafting, abrasion and through seed (Little & Rezaian 2003; Singh, Ready & Nie 2003; Wan Chow Wah & Symons 1997). Mechanical transmission from fruit for human consumption is unlikely. Infected grapevine seedlings are very unlikely to establish, as demonstrated by the full assessment of the likelihood of establishment for <i>Tomato black ring virus</i> in this report. The chance that infected grape seeds from fruit waste will germinate is small. If germination does occur, seedlings are unlikely to survive. | Assessment not required | No |

| Pest | Present in India | Present within Australia | Potential to be on pathway | Potential for establishment and spread | Potential for economic consequences | Pest risk assessment required |
|--|--------------------------------|--|--|--|--|--------------------------------------|
| <i>Hop stunt viroid</i> (HSVd) [Pospiviroidae: Hostuviroid] | Yes (Ramachandran et al. 2005) | Yes Only known to be present in SA and Vic. (Koltunow, Krake & Rezaian 1988). Not known to be present in WA. | Yes HSVd has been demonstrated to be seed transmitted in grapevines (Wan Chow Wah & Symons 1999), but not in any other species. Wan Chow Wah and Symons (1999) confirmed that, in grapevines, HSVd can be transmitted by seed to seedlings. (This authority is cited in Little and Rezaian (2003) which is then cited in Albrechtsen (2006)). HSVd infects systemically and is present in all parts of the plant (Li et al. 2006; Yaguchi & Takahashi 1984). | No The viroid may be transmitted via mechanical means (Sano 2003), through cuttings and grafting (European Food Safety Authority 2008) or via grape seed (Wan Chow Wah & Symons 1999). Mechanical transmission from fruit for human consumption is unlikely. Infected grapevine seedlings are very unlikely to establish, as demonstrated by the full assessment of the likelihood of establishment for <i>Tomato black ring virus</i> in this report. The chance that infected grape seeds from fruit waste will germinate is small. If germination does occur, seedlings are unlikely to survive. | Assessment not required | No |

Appendix B Issues raised in stakeholder comments

A summary of key stakeholder comments and how they were considered in the final report is given below.

Comment 1: One stakeholder asked for the opportunity to review and provide comments on any changes prior to the release of the final policy review.

Response: The Australian Government Department of Agriculture and Water Resources has conducted this non-regulated analysis of existing policy consistent with the *Import Risk Analysis Handbook 2011*. Please note that from 16 June 2016, the *Import Risk Analysis Handbook 2011* will be replaced by the *Biosecurity Import Risk Analysis Guidelines 2015*. For further information on the new guidelines please refer to the [department's website](#). The department is willing to consider new information and modify import conditions if warranted.

Comment 2: Two stakeholders raised concerns over the methodology used in the pest categorisation process that led to several organisms not requiring a pest risk assessment.

Response: The Australian Government Department of Agriculture and Water Resources has conducted the pest categorisation process in accordance with ISPM 11. The evidence considered does not support a strong pathway association for all pests considered. For some of the pests, for example for pathogens which are seed transmitted in grapes and have no other means of establishment, the evidence considered (including full assessments for similar pathogens that are seed transmitted) does not support a potential for the pests to establish and spread. The department considers that the available evidence does not justify a full pest risk assessment for these organisms.

Comment 3: One stakeholder raised concerns that unlisted organisms (organisms which require further assessment by Western Australia before they can be declared as permitted (section 11) or prohibited (section 12)), which may or may not be present in Western Australia, have not been adequately considered in the report.

Response: The Australian Government Department of Agriculture and Water Resources considered all available scientific literature and information in conducting the pest categorisation. Pests that are not known to be present in Western Australia have been assessed further in the pest categorisation process, except for the pests for which Western Australia does not require mitigation measures for other hosts which pose a similar risk from Australian states/territories where this pest is present. The potential of the pests being on the pathway and their potentials for establishment and spread as well as economic consequences have been assessed where appropriate. The Australian Department of Agriculture and Water Resources would review any evidence that stakeholders could provide on pests or pathogens they believe should be considered further.

*Comment 4: One stakeholder considered that some of the pests for which a pest risk assessment was conducted, for example *Planococcus ficus*, *Planococcus lilacinus*, *Rastroccus iceryoides* and *Bactrocera correcta*, are not associated with table grapes in India.*

Response: As the stakeholder did not provide any additional evidence to support its claims, the Australian Government Department of Agriculture and Water Resources further reviewed the existing evidence and concluded that the report does not need to be amended.

Comment 5: One stakeholder considered that pest risk assessments should be conducted for additional pests.

Response: The Australian Government Department of Agriculture and Water Resources further reviewed the evidence for all of the pests the stakeholder considers a pest risk assessment should be conducted for. Following this review, the department conducted a pest risk assessment for one of the pests (*Cryptoblabes gnidiella*). However, the unrestricted risk for this pest for table grapes from India achieves Australia's ALOP and risk management measures are not required for this pest.

Comment 6: One stakeholder considered that some of the likelihood ratings and/or consequence estimates in the pest risk assessments for some pathogens are too low.

Response: The Australian Government Department of Agriculture and Water Resources re-examined the evidence supporting the assessments for these pathogens and concluded that a change of likelihood ratings and/or consequence estimates for these pathogens is not justified. The stakeholder did not provide any new/additional evidence to support the claims.

Comment 7: One stakeholder expressed concerns over the implementation of risk management options.

Response: The Australian Government Department of Agriculture and Water Resources will verify the specific export measures prior to commencement of trade. The risk management measures recommended in this report are considered appropriate to reduce the risk for all identified quarantine pests to achieve Australia's appropriate level of protection (ALOP).

Comment 8: One stakeholder raised concerns over the proposed risk management measure (visual inspection and remedial action) for mealybugs, spider mite and thrips.

Response: The Australian Government Department of Agriculture and Water Resources has successfully used the same measure for these or similar arthropod pests for existing policy. Phytosanitary inspection specifically targets the identified pests and optical enhancement is used where necessary. The efficacy of this measure is supported by considerable trade of table grapes from California to Australia since 2002.

Comment 9: One stakeholder considered that the draft report does not provide enough information about the inspection processes.

Response: Inspection and verification processes are outlined in Chapter 5 of the report. Any consignment of table grapes will be inspected by both India and Australia. As a signatory to the IPPC, India is obligated to fulfil its responsibilities for pre-export inspection and certification required by the importing countries, including Australia. DAC as India's NPPO is responsible for verifying and certifying that the table grapes have been produced in accordance with Australian import requirements. This includes any pest and disease risk mitigation measures in place in the production areas.

The Australian Government Department of Agriculture and Water Resources verifies that imported table grapes comply with Australian import requirements. The department completes a verification inspection of all consignments of table grapes, either in Australia or offshore if requested. The inspection will be conducted in accordance with the department's standard inspection protocol for table grapes, using optical enhancement where necessary.

Glossary

| Term or abbreviation | Definition |
|--|---|
| Additional declaration | A statement that is required by an importing country to be entered on a phytosanitary certificate and which provides specific additional information on a consignment in relation to regulated pests (FAO 2015a). |
| Anamorph | An asexual stage in the life cycle of a fungus, also known as the imperfect state of a fungus. |
| Appropriate level of protection (ALOP) | The level of protection deemed appropriate by the Member establishing a sanitary or phytosanitary measure to protect human, animal or plant life or health within its territory (WTO 1995). |
| Area | An officially defined country, part of a country or all or parts of several countries (FAO 2015a). |
| Area of low pest prevalence | An area, whether all of a country, part of a country, or all parts of several countries, as identified by the competent authorities, in which a specific pest occurs at low levels and which is subject to effective surveillance, control or eradication measures (FAO 2015a). |
| Arthropod | The largest phylum of animals, including the insects, arachnids and crustaceans. |
| Asexual reproduction | The development of new individual from a single cell or group of cells in the absence of meiosis. |
| Biosecurity Australia | The previous name for the unit, within the Australian Government Department of Agriculture and Water Resources, responsible for recommendations for the development of Australia's biosecurity policy. These functions are undertaken within the Plant Division of the Department. |
| Cane (grapevine) | A cane is a ripened shoot of a grapevine that has grown from a new bud located on the cordon. A shoot is called a cane when it changes colour from green to brown during veraison. Shoots give rise to leaves, tendrils and grape clusters. |
| Consignment | A quantity of plants, plant products and/or other articles being moved from one country to another and covered, when required, by a single phytosanitary certificate (a consignment may be composed of one or more commodities or lots) (FAO 2015a). |
| Control (of a pest) | Suppression, containment or eradication of a pest population (FAO 2015a). |
| Diapause | Period of suspended development/growth occurring in some insects, in which metabolism is decreased. |
| Disease | A condition of part or all of an organism that may result from various causes such as infection, genetic defect or environmental stress. |
| Endangered area | An area where ecological factors favour the establishment of a pest whose presence in the area will result in economically important loss (FAO 2015a). |
| Endemic | Belonging to, native to, or prevalent in a particular geography, area or environment. |
| Entry (of a pest) | Movement of a pest into an area where it is not yet present, or present but not widely distributed and being officially controlled (FAO 2015a). |
| Establishment (of a pest) | Perpetuation, for the foreseeable future, of a pest within an area after entry (FAO 2015a). |
| Fecundity | The fertility of an organism. |
| Fresh | Living; not dried, deep-frozen or otherwise conserved (FAO 2015a). |
| Fumigation | A method of pest control that completely fills an area with gaseous pesticides to suffocate or poison the pests within. |
| Genus | A taxonomic category ranking below a family and above a species and generally consisting of a group of species exhibiting similar characteristics. In taxonomic nomenclature the genus name is used, either alone or followed by a Latin adjective or epithet, to form the name of a species. |
| Host | An organism that harbours a parasite, mutual partner, or commensal partner, typically providing nourishment and shelter. |

| Term or abbreviation | Definition |
|--|---|
| Host range | Species capable, under natural conditions, of sustaining a specific pest or other organism (FAO 2015a). |
| Import permit | Official document authorising importation of a commodity in accordance with specified phytosanitary import requirements (FAO 2015a). |
| Import risk analysis | An administrative process through which quarantine policy is developed or reviewed, incorporating risk assessment, risk management and risk communication. |
| Infection | The internal 'endophytic' colonisation of a plant, or plant organ, and is generally associated with the development of disease symptoms as the integrity of cells and/or biological processes are disrupted. |
| Infestation (of a commodity) | Presence in a commodity of a living pest of the plant or plant product concerned. Infestation includes infection (FAO 2015a). |
| Inspection | Official visual examination of plants, plant products or other regulated articles to determine if pests are present and/or to determine compliance with phytosanitary regulations (FAO 2015a). |
| Intended use | Declared purpose for which plants, plant products, or other regulated articles are imported, produced, or used (FAO 2015a). |
| Interception (of a pest) | The detection of a pest during inspection or testing of an imported consignment (FAO 2015a). |
| International Standard for Phytosanitary Measures (ISPM) | An international standard adopted by the Conference of the Food and Agriculture Organization, the Interim Commission on phytosanitary measures or the Commission on phytosanitary measures, established under the IPPC (FAO 2015a). |
| Introduction (of a pest) | The entry of a pest resulting in its establishment (FAO 2015a). |
| Lot | A number of units of a single commodity, identifiable by its homogeneity of composition, origin etc., forming part of a consignment (FAO 2015a). Within this report a 'lot' refers to a quantity of fruit of a single variety, harvested from a single production site during a single pick and packed at one time. |
| Mature fruit | Commercial maturity is the start of the ripening process. The ripening process will then continue and provide a product that is consumer-acceptable. Maturity assessments include colour, starch, index, soluble solids content, flesh firmness, acidity, and ethylene production rate. |
| Mortality | The total number of organisms killed by a particular disease. |
| National Plant Protection Organization (NPPO) | Official service established by a government to discharge the functions specified by the IPPC (FAO 2015a). |
| Official control | The active enforcement of mandatory phytosanitary regulations and the application of mandatory phytosanitary procedures with the objective of eradication or containment of quarantine pests or for the management of regulated non-quarantine pests (FAO 2015a). |
| Parthenogenesis | Production of an embryo from unfertilised egg. |
| Pathogen | A biological agent that can cause disease to its host. |
| Pathway | Any means that allows the entry or spread of a pest (FAO 2015a). |
| Pest | Any species, strain or biotype of plant, animal, or pathogenic agent injurious to plants or plant products (FAO 2015a). |
| Pest categorisation | The process for determining whether a pest has or has not the characteristics of a quarantine pest or those of a regulated non-quarantine pest (FAO 2015a). |
| Pest free area (PFA) | An area in which a specific pest does not occur as demonstrated by scientific evidence and in which, where appropriate, this condition is being officially maintained (FAO 2015a). |
| Pest free place of production | Place of production in which a specific pest does not occur as demonstrated by scientific evidence and in which, where appropriate, this condition is being officially maintained for a defined period (FAO 2015a). |

| Term or abbreviation | Definition |
|---|--|
| Pest free production site | A defined portion of a place of production in which a specific pest does not occur as demonstrated by scientific evidence and in which, where appropriate, this condition is being officially maintained for a defined period and that is managed as a separate unit in the same way as a pest free place of production (FAO 2015a). |
| Pest risk analysis (PRA) | The process of evaluating biological or other scientific and economic evidence to determine whether an organism is a pest, whether it should be regulated, and the strength of any phytosanitary measures to be taken against it (FAO 2015a). |
| Pest risk assessment (for quarantine pests) | Evaluation of the probability of the introduction and spread of a pest and of the associated potential economic consequences (FAO 2015a). |
| Pest risk management (for quarantine pests) | Evaluation and selection of options to reduce the risk of introduction and spread of a pest (FAO 2015a). |
| Pest status (in an area) | Presence or absence, at the present time, of a pest in an area, including where appropriate its distribution, as officially determined using expert judgement on the basis of current and historical pest records and other information (FAO 2015a). |
| Phloem | In vascular plants, the tissue that carries organic nutrients to all parts of the plant where needed. |
| Phytosanitary certificate | An official paper document or its official electronic equivalent, consistent with the model of certificates of the IPPC, attesting that a consignment meets phytosanitary import requirements (FAO 2015a). |
| Phytosanitary certification | Use of phytosanitary procedures leading to the issue of a phytosanitary certificate (FAO 2015a). |
| Phytosanitary measure | Any legislation, regulation or official procedure having the purpose to prevent the introduction and/or spread of quarantine pests, or to limit the economic impact of regulated non-quarantine pests (FAO 2015a). |
| Phytosanitary procedure | An official method for implementing phytosanitary measures including the performance of inspections, tests, surveillance or treatments in connection with regulated pests (FAO 2015a). |
| Phytosanitary regulation | Official rule to prevent the introduction and/or spread of quarantine pests, or to limit the economic impact of regulated non-quarantine pests, including establishment of procedures for phytosanitary certification (FAO 2015a). |
| Polyphagous | Feeding on a relatively large number of hosts from different plant family and/or genera. |
| PRA area | Area in relation to which a pest risk analysis is conducted (FAO 2015a). |
| Practically free | Of a consignment, field or place of production, without pests (or a specific pest) in numbers or quantities in excess of those that can be expected to result from, and be consistent with good cultural and handling practices employed in the production and marketing of the commodity (FAO 2015a). |
| Production site | In this report, a production site is a continuous planting of table grape trees treated as a single unit for pest management purposes. If an orchard is subdivided into one or more units for pest management purposes, then each unit is a production site. If the orchard is not subdivided, then the orchard is also the production site. |
| Quarantine | Official confinement of regulated articles for observation and research or for further inspection, testing or treatment (FAO 2015a). |
| Quarantine pest | A pest of potential economic importance to the area endangered thereby and not yet present there, or present but not widely distributed and being officially controlled (FAO 2015a). |
| Regulated article | Any plant, plant product, storage place, packing, conveyance, container, soil and any other organism, object or material capable of harbouring or spreading pests, deemed to require phytosanitary measures, particularly where international transportation is involved (FAO 2015a). |
| Regulated pest | A quarantine pest or a regulated non-quarantine pest (FAO 2015a). |
| Restricted risk | Risk estimate with phytosanitary measure(s) applied. |
| Saprophyte | An organism deriving its nourishment from dead organic matter. |

| Term or abbreviation | Definition |
|-----------------------------|--|
| Spread (of a pest) | Expansion of the geographical distribution of a pest within an area (FAO 2015a). |
| SPS Agreement | WTO Agreement on the Application of Sanitary and Phytosanitary Measures. |
| Stakeholders | Government agencies, individuals, community or industry groups or organizations, whether in Australia or overseas, including the proponent/applicant for a specific proposal, who have an interest in the policy issues. |
| Systems approach(es) | The integration of different risk management measures, at least two of which act independently, and which cumulatively achieve the appropriate level of protection against regulated pests. |
| Teleomorph | The sexual stage of the life cycle of a fungus. Also called the perfect stage. |
| Trash | Soil, splinters, twigs, leaves, and other plant material, other than fruit stalks. |
| Treatment | Official procedure for the killing, inactivation or removal of pests, or for rendering pests infertile or for devitalisation (FAO 2015a). |
| Unrestricted risk | Unrestricted risk estimates apply in the absence of risk mitigation measures. |
| Vector | An organism that does not cause disease itself, but which causes infection by conveying pathogens from one host to another. |
| Viable | Alive, able to germinate or capable of growth. |

References

- Abdullagatov, A & Abdullagatov, K 1986, 'The gamma moth as a vineyard pest' (in Russian), *Zashchita Rastenii*, vol. 11, pp. 40-2.
- ABRS 2009, 'Australian Faunal Directory', available at <http://www.environment.gov.au/biodiversity/abrs/online-resources/fauna/afd/index.html>, accessed 2015.
- ABS 2009a, *Agricultural commodities, Australia 2007-08*, No. 7121.0, Australian Bureau of Statistics, Canberra.
- -- 2009b, *Australian wine and grape industry 2008 (re-issue)*, No. 1329.0, Australian Bureau of Statistics, Canberra, available at www.abs.gov.au.
- -- 2012a, *Agricultural commodities, Australia 2010-11*, No. 7121.0, Australian Bureau of Statistics, Canberra.
- -- 2012b, *Data for the value of agricultural commodities produced, Australia, 2010-11 report*, Australian Bureau of Statistics, available at http://www.abs.gov.au/AUSSTATS/SUBSCRIBER.NSF/log?openagent&75030do002_201011.xls&7503.0&DataCubes&D5C7C3AFA0BD9D16CA257A2B0017ABB7&0&2010-11&29.06.2012&Previous.
- Adekunle, OK, Kulshrestha, S, Prasad, R, Hallan, V, Raikhy, G, Verma, N, Ram, R, Kumar, S & Zaidi, AA 2006, 'Plant parasitic and vector nematodes associated with Asiatic and Oriental hybrid lilies', *Bioresource Technology*, vol. 97, pp. 364-71.
- Ahmed, MA, Kumar, ST & Dharmaraju, E 1977, 'Cockchafer beetle: *Adoretus bicolor* Brenske damaging grape vine', *Indian Journal of Entomology*, vol. 39, pp. 389-90.
- Akbas, B & Erdiller, G 1993, 'Researches on grapevine virus diseases and determination of their incidences in Ankara, Türkiye', *Journal of Turkish Phytopathology*, vol. 22, no. 2-3, pp. 55-63.
- ALA 2013, 'Atlas of Living Australia', available at www.ala.org.au, accessed 2013.
- Albrechtsen, SE 2006, 'Appendix 1: list of seed-transmitted viruses and viroids', in *Testing methods for seed-transmitted viruses: principles and protocols*, CAB International, Wallingford.
- Alcock, J 1971, 'The behavior of a stinkbug, *Euschistus conspersus* Uhler (Hemiptera: Pentatomidae)', *Psyche*, vol. 78, no. 4, pp. 215-28.
- Alford, DV 2007, *Pest of fruit crops: a colour handbook*, Manson Publishing, London, UK.
- Alma, A, Davis, RE, Vibio, M, Danielli, A, Bosco, D, Arzone, A & Bertaccini, A 1996, 'Mixed infection of grapevines in northern Italy by phytoplasmas including 16S rRNA RFLP subgroup 16SrI-B strains previously unreported in this host', *Plant Disease*, vol. 80, no. 4, pp. 418-21.
- APEDA 2013, *Procedures for export of fresh table grapes to the European Union through control of residues of agro chemicals. Trade notice no: QMC/GEN/056/2013-14*, Agricultural and Processed Food Products Export Development Authority, available at <http://www.apeda.gov.in/apedawebsite/Grapenet/procedureforexportofgrapes2013-14.pdf> (pdf 557kb).

-- -- 2015, *Product profile: fresh grapes*, Agricultural and Processed Food Products Export Development Authority Agri-exchange, available at <http://agriexchange.apeda.gov.in>.

AQIS 1999, *Draft import risk analysis for the importation of fresh table grapes [Vitis vinifera L.] from California (USA)*, Australian Quarantine and Inspection Service, Canberra.

-- -- 2000, *Final import risk analysis for the importation of fresh table grapes (Vitis vinifera L.) from the state of California in the United States of America*, Australian Quarantine and Inspection Service, Canberra.

Armstrong, JW, Silva, ST & Shishido, VM 1995, 'Quarantine cold treatment for Hawaiian Carambola fruit infested with Mediterranean fruit fly, melon fly, or Oriental fruit fly (Diptera: Tephritidae) eggs and larvae', *Journal of Economic Entomology*, vol. 88, no. 3, pp. 683-7.

Arrigo, N & Arnold, C 2007, 'Naturalised *Vitis* rootstocks in Europe and consequences to native wild grapevine' *PloS ONE*, vol. 2, no. 6, pp. e521, available at DOI 10.1371/journal.pone.0000521.

Ashihara, W 1996, 'Effect of grape cultivars and insecticide application on reproduction and occurrence of *Tetranychus kanzawai* Kishida (Acarina: Tetranychidae)' (in Japanese), *Japanese Journal of Applied Entomology and Zoology*, vol. 40, no. 1, pp. 55-9.

Asjes, CJ & Blom-Barnhoorn, GJ 2002, 'Control of spread of Augusta disease caused by *Tobacco necrosis virus* in tulip by composting residual waste of small bulbs, tunics, roots and soil debris', *Acta Horticulturae (ISHS)*, vol. 570, pp. 283-6.

Astridge, D & Elder, R 2005, *Oriental scale in papayas*, Department of Employment, Economic Development and Innovation, Primary Industries and Fisheries, Queensland, available at <http://www2.dpi.qld.gov.au/>.

Atallah, J, Teixeira, L, Salazar, R, Zaragoza, G & Kopp, A 2014, 'The making of a pest: the evolution of a fruit-penetrating ovipositor in *Drosophila suzukii* and related species' *Proceedings of the Royal Society Biological Sciences*, vol. 281, no. 1781, pp. e20132840, available at <http://dx.doi.org/10.1098/rspb.2013.2840>.

ATGA 2013, *Varieties*, Australian Table Grape Association, available at <http://www.australiangrapes.com.au/about-atga2/varieties>.

Australian Museum 2009, *Animal species: hawk moths*, available at <http://australianmuseum.net.au/Hawk-Moths>.

Australian Table Grape Association 2013, *Australian table grapes: the pick of the bunch*, Australian Table Grape Association, available at http://www.australiangrapes.com.au/data/assets/pdf_file/0017/17225/HAL0462_Grapes_Industry_Booklet.pdf (pdf 770 kb).

Ausveg 2013a, *Vegetables - where current data is available*, Ausveg, Camberwell.

-- -- 2013b, *Vegetables - where current data is not available*, Ausveg, Camberwell.

Badyal, K & Sumbali, G 1990, 'Market diseases of mango', *Indian Journal of Mycology and Plant Pathology*, vol. 20, p. 281.

Bagnoli, B & Lucchi, A 2001, 'Bionomics of *Cryptoblabes gnidiella* (Millière) (Pyralidae Phycitinae) in Tuscan vineyards', *IOBC wprs Bulletin*, vol. 24, no. 7, pp. 79-83.

Barbetti, MJ, Riley, IT, You, MP, Li, H & Sivasithamparam, K 2006, 'The association of necrotrophic fungal pathogens and plant parasitic nematodes with the loss of productivity of

annual medic-based pastures in Australia and options for their management', *Australasian Plant Pathology*, vol. 35, pp. 691-706.

Barkley, P & Büchen-Osmond, C 1988, 'Distribution, hosts and vectors of viruses in Australia', in *Viruses of plants in Australia: descriptions and lists from the VIDE database*, Büchen-Osmond, C, Crabtree, K, Gibbs, K & McLean, G (eds), The Australian National University Printing Press, Canberra.

Batra, RC, Brar, SS, Khangura, JS & Dhillon, WS 1987, 'A new record of *Planococcus pacificus* Cox (Pseudococcidae: Hemiptera) as a pest of grapevine in India', *Punjab Horticultural Journal*, vol. 27, pp. 250-1.

Batra, RC, Brar, SS, Singh, R & Kapoor, S 1980, 'Carbaryl for the control of *Rhipiphorothrips cruentatus* Hood and thinning of grape berries', *Science and Culture*, vol. 46, pp. 75-6.

Belli, G, Bianco, PA & Conti, M 2010, 'Grapevine yellows in Italy: past, present and future', *Journal of Plant Pathology*, vol. 92, no. 2, pp. 303-26.

Ben-Dov, Y 2012a, 'Coccidae', available at <http://www.sel.barc.usda.gov/scalenet/scalenet.htm>, accessed 2012.

-- -- 2012b, 'Margarodidae', available at <http://www.sel.barc.usda.gov/scalenet/scalenet.htm>, accessed 2012.

-- -- 2012c, 'Pseudococcidae', available at <http://www.sel.barc.usda.gov/scalenet/scalenet.htm>, accessed 2012.

-- -- 2013, 'Coccidae', available at <http://www.sel.barc.usda.gov/scalenet/scalenet.htm>, accessed 2013.

Bensch, K, Groenewald, JZ, Dijksterhuis, J, Starink-Willemse, M, Andersen, B, Summerell, BA, Shin, H-D, Dugan, FM, Schroers, H-J, Braun, U & Crous, PW 2010, 'Species and ecological diversity within the *Cladosporium cladosporioides* complex (*Davidiellaceae*, *Capnodiales*)', *Studies in Mycology*, vol. 67, pp. 1-94.

Bentley, WJ, Varela, LG, Zalom, FG, Smith, RJ, Purcell, AH, Phillips, PA, Haviland, DR, Daane, KM & Battany, MC 2008, *Omnivorous leafroller*, UC IPM Pest Management Guidelines: Grape, available at <http://www.ipm.ucdavis.edu/PMG/r302300311.html>.

Bhagat, RC, Ramzan, A & Farhan, N 1991, 'New records of scale insects (Homoptera: Coccoidea) and host-plants from Kashmir Valley, India', *Entomon.*, vol. 16, no. 1, pp. 91-4.

Bharathie, KP 1975, 'Tortricid moth [*Archips micaceana*] damage to Pinus [*P. patula* and *P. insularis*] seedlings in Sri Lanka', *Sri Lanka Forester*, vol. 12, no. 1, pp. 37-9.

Bianco, PA, Davis, RE, Prince, JP, Fortusini, A, Casati, P & Belli, G 1994, 'Elm yellows and aster yellows MLOs associated with a grapevine yellows disease very similar to Flavescence doree in northern Italy', *International Organization of Mycoplasmaology (IOM) Letters*, vol. 3, pp. 251-2.

Bindra, OS & Varma, GC 1979, 'Insect pests of grape-vines in North India', *Food Farming and Agriculture*, vol. 10, pp. 277-82.

Biosecurity Australia 2002, *Risk analysis glassy-winged sharpshooter (GWSS), Homalodisca coagulata on Californian table grapes*, Biosecurity Australia, Canberra.

-- -- 2003, *Netherlands truss tomatoes import policy*, Department of Agriculture, Fisheries and Forestry, available at <http://www.daff.gov.au/>.

- -- 2005a, *Final report: extension of existing policy for sweet oranges from Italy*, Department of Agriculture, Fisheries and Forestry, Canberra.
- -- 2005b, *Final report: import risk analysis for table grapes from Chile*, Department of Agriculture, Fisheries and Forestry, Canberra.
- -- 2006a, *Biosecurity Australia policy memorandum 2006/32: revised conditions for the importation of Californian table grapes*, Department of Agriculture, Fisheries and Forestry, Canberra.
- -- 2006b, *Policy for the importation of fresh mangoes (Mangifera indica L.) from Taiwan*, Biosecurity Australia, Department of Agriculture, Fisheries and Forestry, Canberra, available at http://www.daff.gov.au/_data/assets/word_doc/0017/27062/2006_24a.doc.
- -- 2008, *Final import risk analysis report for fresh mango fruit from India*, Biosecurity Australia, Department of Agriculture, Fisheries and Forestry, Canberra, available at http://www.daff.gov.au/_data/assets/pdf_file/0003/771906/Mangoes_from_India_Final_Report.pdf.
- -- 2009, *Final import risk analysis report for fresh unshu mandarin fruit from Shizuoka prefecture in Japan*, Department of Agriculture, Fisheries and Forestry, Canberra.
- -- 2010, *Final import risk analysis report for fresh apple fruit from the People's Republic of China*, Biosecurity Australia, Department of Agriculture, Fisheries and Forestry, Canberra, available at [http://www.daff.gov.au/_data/assets/pdf_file/0006/1686471/Final IRA report for apples from the Peoples Republic of China.pdf](http://www.daff.gov.au/_data/assets/pdf_file/0006/1686471/Final_IRA_report_for_apples_from_the_Peoples_Republic_of_China.pdf).
- -- 2011a, *Final import risk analysis report for table grapes from the People's Republic of China*, Department of Agriculture, Fisheries and Forestry, Canberra, available at http://www.daff.gov.au/_data/assets/pdf_file/0010/1943938/FinalIRA_Tablegrapes_China.pdf (pdf 4.28 mb).
- -- 2011b, *Final non-regulated risk analysis report for table grapes from the Republic of Korea*, Department of Agriculture, Fisheries and Forestry, Canberra, available at http://www.daff.gov.au/_data/assets/pdf_file/0003/1990560/Final-report-Korean-table-grapes.pdf (pdf 2.55 mb).
- Bisiach, M 1988, 'White rot', in *Compendium of grape diseases*, Pearson, RC & Goheen, AC (eds), The American Phytopathological Society Press, St. Paul, Minnesota.
- Bisiach, M & Viterbo, AB 1973, 'Further researches on grapevine cluster drying-off caused by *Coniothyrium diplodiella*', *Mededelingen Fakulteit Landbouwwetenschappen, Rijksuniversiteit Gent*, vol. 38, no. 8, pp. 1561-71.
- Bisotto-de-Oliveira, R, Redaelli, LR, Sant'ana, J, Cover, C & Botton, M 2007, 'Ocorrência de *Cryptoblabes gnidiella* (Millière) (Lepidoptera: Pyralidae) relacionada à fenologia da videira em Bento Gonçalves, RS' (Occurrence of *Cryptoblabes gnidiella* (Milliere) (Lepidoptera: Pyralidae) associated with grape phenology in Bento Gonçalves, RS), *Neotropical Entomology*, vol. 36, no. 4, pp. 555-9.
- Bissett, J 1982, '*Pestalotiopsis menezesiana* on greenhouse plantings of *Cissus rhombifolia* with notes on related fungi occurring on Vitaceae', *Canadian Journal of Botany*, vol. 60, pp. 2570-4.
- Bos, L 1999, 'Symptoms of virus diseases', in *Plant viruses, unique and intriguing pathogens: a textbook of plant virology*, Backhuys Publishers, Leiden.

- Botha, J & Learmonth, S 2005, *Management of European red mite in Western Australia*, Farmnote no. 40/2005, Department of Agriculture, Western Australia.
- Bournier, A 1977, 'Grape insects', *Annual Review of Entomology*, vol. 22, pp. 355-76.
- Bradbury, JF 1986, *Guide to plant pathogenic bacteria*, CAB International, Slough, UK.
- Brewer, LJ, Walton, V, Dreves, A, Shearer, P, Zalom, F & Walsh, D 2012, 'Biology and management of spotted wing drosophilla on small and stone fruits: Year 2 reporting cycle', Department of Horticulture, Oregon State University, Oregon, available at http://spottedwing.org/system/files/SWD_ResearchReviewYear%202012_7.16.12.pdf (pdf 3.21 mb).
- Brookes, HM 1964, 'The Coccoidea (Homoptera) naturalized in South Australia: a second annotated list', *Transactions of the Royal Society of South Australia*, vol. 88, pp. 15-20.
- Brown, DJF, Murant, AF & Trudgill, DL 1989, 'Differences between isolates of the English serotype of *Tomato black ring virus* in their transmissibility by an English population of *Longidorus attenuatus* (Nematoda: Dorylaimoidea)', *Revue de Nematologie*, vol. 12, no. 1, pp. 51-6.
- Brunt, AA, Crabtree, K, Dallwitz, MJ, Gibbs, AJ, Watson, L & Zurcher, EJ 1996, 'Plant viruses online: descriptions and lists from the VIDE database', available at <http://pvo.bio-mirror.cn/refs.htm>, accessed 2015.
- Brunt, AA & Teakle, DS 1996, '*Tobacco necrosis necrovirus*', in *Viruses of plants: descriptions and lists from the VIDE database*, Brunt, AA, Crabtree, K, Dallwitz, MJ, Gibbs, A & Watson, L (eds), CAB International, Wallingford, UK.
- Buchanan, GA & Whiting, JR 1991, 'Phylloxera management: prevention is better than cure', *The Australian and New Zealand Wine Industry Journal*, vol. 6, no. 3, pp. 223-30.
- Büchen-Osmond, C, Crabtree, K, Gibbs, A & McLean, G, (eds) 1988, *Viruses of plants in Australia: descriptions and lists from the VIDE database*, The Australian National University Research School of Biological Sciences, Canberra.
- Buck, M, Marshall, SA & Cheung, DKB 2008, 'Identification atlas of the Vespidae (Hymenoptera, Aculeata) of the northeastern Nearctic region', *Canadian Journal of Arthropod Identification*, vol. 5, pp. 1-492.
- Bureau of Meteorology 2010, 'Average annual and monthly maximum, minimum, and mean temperature', available at <http://www.bom.gov.au/>, accessed 2010.
- Burges, N, Taylor, A & Kumar, S 2005, 'Phomopsis cane and leaf spot: *Phomopsis viticola* (synonym: *P. viticola* taxon 2) an exotic threat to Western Australia', *Department of Agriculture Factsheet*, vol. 9, pp. 1-2.
- Butani, DK 1993, *Mango: pest problems*, Periodical Expert Book Agency, Dehli.
- CABI 2012, 'Crop Protection Compendium', available at <http://www.cabi.org/cpc>, accessed 2012.
- -- 2013, 'Crop Protection Compendium', available at <http://www.cabi.org/cpc>, accessed 2013.
- -- 2014, 'Crop Protection Compendium', available at <http://www.cabi.org/cpc>, accessed 2014.
- -- 2015, 'Crop Protection Compendium', available at <http://www.cabi.org/cpc>, accessed 2015.

- -- 2016, 'Crop Protection Compendium', available at <http://www.cabi.org/cpc>, accessed 2016.
- CABI & EPPO 1979, *Commonwealth Institute of Entomology, distribution maps of plant pests, map no. 45: Myzus persicae (Sulzer)*, available at <http://www.cabi.org/DMPP/>.
- -- 1986, *Commonwealth Institute of Entomology, distribution maps of plant pests, map no. 76: Eupoecilia ambiguella*, CAB Abstracts plus, available at <http://www.cababstractsplus.org/DMPP/Index.asp>.
- -- 1997a, *Data sheets on quarantine pests: Aleurocanthus spiniferus*, European and Mediterranean Plant Protection Organization, available at http://www.eppo.int/QUARANTINE/insects/Aleurocanthus_spiniferus/ALECSN_ds.pdf.
- -- 1997b, *Datasheets on quarantine pests: Aleurocanthus woglumi*, European and Mediterranean Plant Protection Organization, available at <http://www.eppo.org/>.
- -- 1997c, *Datasheets on quarantine pests: Scirtothrips citri*, European and Mediterranean Plant Protection Organization, available at <http://www.eppo.org>.
- -- 1997d, 'Datasheets on quarantine pests: Tomato black-ring nepovirus', in *Quarantine pests for Europe*, 2nd edn, Smith, IM, McNamara, DG, Scott, PR & Holderness, M (eds), CAB International, Wallingford.
- Card, SD, Pearson, MN & Clover, GRG 2007, 'Plant pathogens transmitted by pollen', *Australasian Plant Pathology*, vol. 36, no. 5, pp. 455-61.
- Cardoso, JM, Felix, MR, Clara, MI & Oliveira, S 2005, 'The complete genome sequence of a new necrovirus isolated from *Olea europaea* L', *Archives of Virology*, vol. 150, pp. 815-23.
- Carrieri, R, Carotenuto, G & Lahoz, E 2013, 'Characterization and pathogenicity of *Pestalotiopsis uvicola* causing black leaf spot on carob (*Ceratonia siliqua* L.) in Italy', *European Journal of Plant Pathology*, vol. 137, pp. 655-61.
- Carter, DJ 1984, *Pest Lepidoptera of Europe: with special reference to the British Isles*, Dr W. Junk Publishers, Dordrecht, The Netherlands.
- Castillo-Pando, M, Somers, A, Green, CD, Priest, M & Sriskanthades, M 2001, 'Fungi associated with dieback of Semillon grapevines in the Hunter Valley of New South Wales', *Australasian Plant Pathology*, vol. 30, pp. 59-63.
- Castillo-Pando, M, Steel, C & Somers, T 1999, 'Fungal bunch rots of grapes in the Hunter and Hastings Valleys', *The Australian Grapegrower and Winemaker*, vol. September, pp. 33-6.
- Cesati, RR & Van Regenmortel, MHV 1969, 'Serological detection of a strain of *Tobacco necrosis virus* in grapevine leaves', *Journal of Phytopathology*, vol. 64, no. 4, pp. 362-6.
- CFIA 2010, *Plant health risk assessment: Drosophila suzukii (Matsumura) Spotted wing drosophila*, Canadian Food Inspection Agency, Ottawa.
- Chakraborty, P, Gupta-Bhattacharya, S, Chowdhury, I, Ranjan Majumdar, M & Chanda, S 2001, 'Differences in concentrations of allergenic pollens and spores at different heights on an agricultural farm in West Bengal, India', *Annals of Agricultural and Environmental Medicine*, vol. 8, pp. 123-30.
- Chand, R, Chaurasia, S, Singh, SK & Sokhi, SS 1999, 'Grapevine bacterial canker disease: profile of problem and its management', in *Diseases of horticultural crops: fruits*, Verma, LR & Sharma, RC (eds), Indus Publishing Company, New Delhi.

- Chand, R & Kishun, R 1990, 'Outbreak of grapevine bacterial canker disease in India', *Vitis*, vol. 29, pp. 183-8.
- Chand, R, Patil, BP & Kishun, R 1991, 'Management of bacterial canker disease (*Xanthomonas campestris* pv *viticola*) of grape vine (*Vitis vinifera*) by pruning', *Indian Journal of Agricultural Sciences*, vol. 61, no. 3, pp. 220-2.
- -- 1992, 'Efficacy of different chemicals against grapevine bacterial canker disease (*Xanthomonas campestris* pv. *viticola*)', *Indian Journal of Plant Protection*, vol. 20, pp. 108-10.
- Chand, R, Singh, PN, Singh, D & Singh, R 1994, 'Copper and streptomycin resistance in *Xanthomonas campestris* pv. *viticola*', *Journal of Plant Diseases and Protection*, vol. 101, no. 5, pp. 487-91.
- Chowfla, SC, Sharma, PK & Thakur, PD 1999, 'Viruses infecting tomato and their management', in *Diseases of horticultural crops: vegetables, ornamentals and mushrooms*, Verma, LR & Sharma, RC (eds), Indus Publishing Company, New Delhi.
- Christenson, LD & Foote, RH 1960, 'Biology of fruit flies', *Annual Review of Entomology*, vol. 5, pp. 171-92.
- Chu, PWG, Francki, RIB & Hatta, T 1983, 'Some properties of *Tomato ringspot virus* isolated from *Pentas lanceolata* in South Australia', *Plant Pathology*, vol. 32, pp. 353-6.
- Chu, Y & Tung, CH 1996, 'The laboratory observations on the attack of Oriental fruit fly, *Bactrocera dorsalis* (Hendel) on grapes' (in Chinese), *Plant Protection Bulletin (Taichung)*, vol. 38, pp. 49-57.
- CIA 2013, *Page size base map: India transportation*, available at <https://www.cia.gov/library/publications/resources/cia-maps-publications/India.html> (PDF 2.65 MB).
- CIHEAM 2006, *Directory of infectious diseases of grapevines and viruses and virus-like diseases of the grapevine: bibliographic report 1998-2004*, Options Méditerranéennes série B, No. 55, Centre International de Hautes études Agronomiques Méditerranéennes-Istituto Agronomico Mediterraneo di Bari, Bari.
- Cline, H 1986, 'Berry scarring problem tied to early thrips', *California Grape Grower*, vol. 17, no. 10, pp. 36-7.
- Collyer, E 1976, 'Integrated control of apple pests in New Zealand: vi. incidence of European red mite, *Panonychus ulmi* (Koch), and its predators', *New Zealand Journal of Zoology*, vol. 3, no. 1, pp. 39-50.
- Common, IFB 1990, *Moths of Australia*, Melbourne University Press, Carlton.
- Conn, BJ 2010, *Vitis vinifera L.*, New South Wales Flora Online: PlantNET, available at <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Vitis~vinifera>.
- Cook, RP & Dubé, AJ 1989, *Host-pathogen index of plant diseases in South Australia*, Field Crops Pathology Group, South Australian Department of Agriculture, Adelaide.
- Coutts, RHA, Rigden, JE, Slabas, AR, Lomonosoff, GP & Wise, PJ 1991, 'The complete nucleotide sequence of *Tobacco necrosis virus* strain D', *Journal of General Virology*, vol. 72, pp. 1521-9.
- Cranshaw, W 2008, *European paper wasp*, no. 5.611, Colorado State University Extension, Fort Collins, available at <http://www.ext.colostate.edu/pubs/insect/05611.html>.

- Cranshaw, WS, Larsen, HJ, Jnr & Zimmerman, RJ 2011, 'Notes on fruit damage by the European paper wasp, *Polistes dominula* (Christ) (Hymenoptera: Vespidae)', *Southwestern Entomologist*, vol. 36, no. 1, pp. 103-5.
- Credi, R, Babini, AR, Betti, L, Bertaccini, A & Gelli, C 1981, 'A distinctive isolate of *Strawberry latent ringspot virus* from grapevines in Italy', *Phytopathologia Mediterranea*, vol. 20, pp. 56-63.
- CSIRO 2005, 'Australian insect common names version 1.53', available at http://www.ces.csiro.au/aicn/name_s/b_1.htm, accessed 2015.
- Cunnington, JH, Priest, MJ, Powney, RA & Cother, NJ 2007, 'Diversity of *Botryosphaeria* species on horticultural plants in Victoria and New South Wales', *Australasian Plant Pathology*, vol. 36, no. 2, pp. 157-9.
- DAC 2004, *Fruits and vegetables grading and marking rules, 2004*, Gazette of India, Department of Agriculture and Cooperation, available at <http://agmarknet.nic.in/fveggmrules04.htm>.
- -- 2007, *Fruits and vegetables grading and marking (amendment) rules, 2007*, Gazette of India, Department of Agriculture and Cooperation, available at <http://agmarknet.nic.in/FruitVegGmRules.pdf> (pdf 104 kb).
- Dadmal, SM, Pawar, NP, Ghawde, SM & Kale, KB 2001, 'Efficacy of plant products and chemical insecticides against the citrus thrips, *Scritothrips citri* Moulton', *Pest Management and Economic Zoology*, vol. 9, no. 1, pp. 93-5.
- DAFF 2004a, *Longan and lychee fruit from the People's Republic of China and Thailand: Final import risk analysis report - Part A and Part B*, Department of Agriculture, Fisheries and Forestry, Canberra, available at http://www.daff.gov.au/_data/assets/word_doc/0013/164002/ll_finala.doc.
- -- 2004b, *Persimmon fruit (*Diospyros kaki* L.) from Japan, Korea and Israel: final import policy*, Department of Agriculture, Fisheries and Forestry, Canberra, available at http://www.daff.gov.au/_data/assets/pdf_file/0013/25141/persimmon_final.pdf.
- -- 2008, *Horticulture-Thailand: phytosanitary requirements update*, Department of Agriculture, Fisheries and Forestry, available at <http://www.daff.gov.au/>.
- -- 2010, *Report on the verification visit to the table grape production area of Nasik in India, 10-16 April 2010*, Biosecurity Australia, Canberra.
- -- 2012, *Final import risk analysis report for the importation of fresh decrowned pineapple (*Ananas comosus* (L.) Merr.) fruit from Malaysia*, Department of Agriculture, Fisheries and Forestry, Canberra.
- -- 2013, *Final non-regulated analysis of existing policy for Californian table grapes to Western Australia*, Department of Agriculture, Fisheries and Forestry, Canberra, available at http://daff.gov.au/_data/assets/pdf_file/0005/2325380/Final-non-regulated-CA-tgrapes-to-WA.pdf.
- DAFWA 2006, *Wine at a glance from Western Australia*, Bulletin 4684, Department of Agriculture and Food, Western Australia, available at http://www.agric.wa.gov.au/objtwr/imported_assets/content/hort/vit/bulletin46854.pdf (pdf 378 kb).
- -- 2008, *Department of Agriculture and Food, Western Australia's submission to the draft import risk analysis for the importation of fresh Unshu mandarin fruit from Japan*, Department of Agriculture and Food, Western Australia.

- -- 2009a, *Western Australian certified seed potato scheme production rules*, AGWEST Plant Laboratories, Perth.
- -- 2009b, *Western Australian registered seed potato scheme production rules*, AGWEST Plant Laboratories, Perth.
- -- 2013, *Comments on draft pest categorisation for table grapes from Japan*, Government of Western Australia, Department of Agriculture and Food, South Perth.
- -- 2014, 'Quarantine WA import requirements', available at <http://www.agric.wa.gov.au/qtine/default.asp>, accessed 2014.
- -- 2015a, *Draft policy review: a categorisation of invertebrate and pathogen organisms associated with fresh table grape bunches (Vitis spp.) imported from other Australian states and territories*, Department of Agriculture and Food, Western Australia, South Perth, available at <https://www.agric.wa.gov.au/plant-biosecurity/table-grapes-stakeholder-consultation-draft-pest-risk-analysis> (pdf 5312 KB).
- -- 2015b, *Draft policy review: fresh table grape bunches (Vitis spp.) imported into Western Australia from other states and territories*, Department of Agriculture and Food, Western Australia, South Perth, available at <https://www.agric.wa.gov.au/plant-biosecurity/table-grapes-stakeholder-consultation-draft-pest-risk-analysis> (pdf 6560 kb).
- -- 2015c, *Final policy review: A categorisation of invertebrate and pathogen organisms associated with fresh table grape bunches (Vitis spp.) imported from other Australian states and territories*, Department of Agriculture and Food, Western Australia, South Perth, available at <https://www.agric.wa.gov.au/plant-biosecurity/table-grapes-final-policy-review> (4553 kb pdf).
- -- 2015d, *Final policy review: Fresh table grape bunches (Vitis spp.) imported into Western Australia from other states and territories*, Department of Agriculture and Food, Western Australia, South Perth, available at <https://www.agric.wa.gov.au/plant-biosecurity/table-grapes-final-policy-review> (5883 kb pdf).
- -- 2015e, *Submission on the Draft report for the non-regulated analysis of existing policy for table grapes from India*, Government of Western Australia, Department of Agriculture and Food, Perth.
- Dahiya, KK & Lakra, RK 2001, 'Seasonal occurrence and succession of thrips, *Rhipiphorothrips cruentatus* Hood in important horticultural crops of Haryana', *Crop Research (Hisar)*, vol. 21, no. 1, pp. 112-4.
- Damm, U, Fourie, PH & Crous, PW 2007, '*Aplosporella prunicola*, a novel species of anamorphic *Botryosphaeriaceae*', *Fungal Diversity*, vol. 27, pp. 35-43.
- Das, BP & Gupta, VK 1989, *The social wasps of India and the adjacent countries (Hymenoptera: Vespidae)*, Association for the Study of Oriental Insects, Gainesville, Florida.
- Davey, PM 1954, 'Quantities of food eaten by the desert locust, *Schistocerca gregaria* (Forsk), in relation to growth', *Bulletin of Entomological Research*, vol. 45, pp. 539-51.
- David'yan, GE 2008, *Parthenolecanium corni* Bouche: *European fruit scale*, Interactive agricultural ecological atlas of Russia and neighbouring countries: economic plants and their diseases, pests and weeds, available at <http://www.agroatlas.ru>.
- David, Z & Rafaila, C 1966, 'Cercetari asupra putregaiului alb la struguri produs de ciuperca *Coniothyrium diplodiella* (Speg.) Sacc' (Study on white rot of Grapevine caused by *Coniella diplodiella*), *Analele Institutului de Cercetari pentru Protectia Plantelor*, vol. 4, pp. 143-53.

Davis, RE, Jomantiene, R, Dally, EL & Wolf, TK 1998, 'Phytoplasmas associated with grapevine yellows in Virginia belong to group 16SrI, subgroup A (tomato big bud phytoplasma subgroup), and group 16SrIII, new subgroup I', *Vitis*, vol. 37, no. 3, pp. 131-7.

DAWA 2006a, *Draft plant pathology lists of pathogen species: table one: pathogens (fungi, bacteria, viruses, viroids and nematodes) recorded from Western Australia*, Department of Agriculture, Perth.

-- -- 2006b, *Table grapes from Western Australia at a glance*, Bulletin 4626, Department of Agriculture, Western Australia, available at http://www.agric.wa.gov.au/objtwr/imported_assets/content/hort/vit/cp/tg/bulletin4626.pdf.

Deacon, J 2005, *Profiles of fungi: rust fungi (Basidiomycota)*, Fungal Biology, available at <http://www.biology.ed.ac.uk/research/groups/jdeacon/FungalBiology/rust.htm>.

Deighton, FC 1976, 'Three fungi on leaves of *Vitis*', *Transactions of the British Mycological Society*, vol. 67, no. 2, pp. 223-32.

Department of Agriculture 2013a, 'Import Conditions Database', available at http://apps.daff.gov.au/icon32/asp/ex_querycontent.asp, accessed 2013.

-- -- 2014, *Final report for the non-regulated analysis of existing policy for fresh table grapes from Japan*, Department of Agriculture, Canberra, available at <http://www.agriculture.gov.au/SiteCollectionDocuments/ba/reviews/final-plant/table-grapes-japan/final-report-non-regulated-analysis-table-grapes-japan.pdf>.

Department of Agriculture, Fisheries and Forestry Biosecurity 2013b, *Final pest risk analysis report for *Drosophila suzukii**, Department of Agriculture, Fisheries and Forestry, Canberra, available at http://daff.gov.au/_data/assets/pdf_file/0004/2281738/Final-PRA-report-Drosophila-suzukii.pdf (pdf 1.97 mb).

Department of Primary Industry, Parks, Water and Environment 2011, *European and English wasps*, available at <http://www.dpiw.tas.gov.au/inter.nsf/webpages/cpas-5y9793?open>.

Devasahayam, S & Nair, CPR 1986, 'The tea mosquito bug *Helopeltis antonii* Signoret on cashew in India', *Journal of Plantation Crops*, vol. 14, no. 1, pp. 1-10.

Dhooria, MS 2003, 'Economic status of phytophagous/predaceous mites from different plants in Punjab', *Journal of Insect Science*, vol. 16, no. 1-2, pp. 12-7.

Dhooria, MS, Bhullar, MB & Mallik, B 2005, *Mite pests of citrus and their management in India*, AINP Agricultural Acarology, Bangalore.

Di Vecchi-Staraz, M, Laucou, V, Bruno, G, Lacombe, T, Gerber, S, Bourse, T, Boselli, M & This, P 2009, 'Low level of pollen-mediated gene flow from cultivated to wild grapevine: consequences for the evolution of the endangered subspecies *Vitis vinifera* L. subsp. *silvestris*', *Journal of Heredity*, vol. 100, no. 1, pp. 66-75.

Doijode, SD 2001, 'Grape: *Vitis vinifera* L.', in *Seed storage in fruit crops*, Haworth Press, New York.

DPIF 2013, *Northern Territory Plant Health Manual - Version 3.0*, Department of Primary Industry and Fisheries, Northern Territory, available at http://www.nt.gov.au/d/Primary_Industry/Content/File/quarantine/Plant%20Health%20Manual%20Version%203_0%2001%20July%202013.pdf.

- DPIPWE 2011, *Plant Quarantine Manual Tasmania - 2012 Edition*, Department of Primary Industries, Parks, Water and Environment, Hobart, Tasmania, available at www.dpipwe.tas.gov.au.
- DPIPWE Tasmania 2015, *Plant Biosecurity Manual Tasmania - 2016 Edition*, Hobart, Australia, available at <http://dpiipwe.tas.gov.au/biosecurity/plant-biosecurity/plant-biosecurity-manual> (PDF 1.385kb).
- DPP 2007, *Export of grapes from India to Australia*, Directorate of Plant Protection, Ministry of Agriculture, India.
- -- 2009, 'Information provided by the Directorate of Plant Protection for conducting a pest risk analysis for export of table grapes from India', Directorate of Plant Protection, Ministry of Agriculture, India.
- -- 2012, *Additional information: pest and pathogens and commercial production practices of table grapes in India: export to Australia*, Directorate of Plant Protection, Quarantine and Storage, Department of Agriculture and Cooperation, India, Faridabad.
- Drew, RAI & Hancock, DL 1994, 'The *Bactrocera dorsalis* complex of fruit flies (Diptera: Tephritidae: Dacinae) in Asia', *Bulletin of Entomological Research*, vol. Suppl. 2, pp. 1-68.
- Dubey, AK & Ko, CC 2009, 'A review of the genus *Aleurolobus* Quaintance and Baker (Hemiptera: Aleyrodidae) from Taiwan, based mainly on pupal morphology with a description of a new species', *Entomological Science*, vol. 12, pp. 51-66.
- Dubey, AK, Ko, CC & David, BV 2008, 'The genus *Singhius* Takahashi (Hemiptera: Aleyrodidae): with notes on pupal dimorphism and intraspecific variation, and a key to the species', *Zoological Studies*, vol. 47, no. 4, pp. 507-18.
- Dye, DW 1978, 'Genus IX: *Xanthomonas* Dowson 1939. In: Young JM, Dye DW, Bradbury JF, Panagopoulos CG, Robbs CF (1978) A proposed nomenclature and classification for plant pathogenic bacteria', *New Zealand Journal of Agricultural Research*, vol. 21, pp. 153-77.
- Ellis, MA 2008, *Bitter rot of grape*, The Ohio State University Extension Factsheet, available at http://ohioline.osu.edu/hyg-fact/3000/pdf/HYG_3032_08.pdf (pdf 267 kb).
- Ellis, RH, Hong, TD & Roberts, EH 1985, 'Vitaceae', in *Handbook of seed technology for genebanks: vol. 2*, International Board for Plant Genetic Resources, Rome.
- Elmer, WH, Summerell, BA, Burgess, LW, Backhouse, D & Abubaker, AA 1997, 'Fusarium species associated with asparagus crowns and soil in Australia and New Zealand', *Australasian Plant Pathology*, vol. 26, pp. 255-61.
- Emmett, BJ 2006, *Preliminary table grape pathogens of concern for China*, Department of Primary Industries, Mildura.
- Eppler, A, Lesan, V & Lazar, A 1989, 'Virus and virus diseases in some vineyards in Romania', *Mededelingen van de Faculteit Landbouwkundige, Rijksuniversiteit Gent*, vol. 54, pp. 491-7.
- EPPO 2007, *Phakopsora euvitis eradicated from Australia. Num. article: 2007/153*, EPPO Global Database, available at <http://gd3.eppo.int/reporting.php/article1172>.
- -- 2011, 'PQR-EPPO database on quarantine pests (available online)', available at <http://www.eppo.org/DATABASES/pqr/pqr.htm>, accessed 2011.

European Food Safety Authority 2008, 'Pest risk assessment made by France on *Hop stunt viroid* (HSVd) considered by France as harmful in the French Overseas Department of Réunion', *The EFSA Journal*, vol. 687, pp. 1-17.

Eyres, N, Wood, C & Taylor, A 2006, *Black rot Guignardia bidwellii*, Factsheet note 167, Department of Agriculture, Western Australia.

FAO 1995, *International Standards for Phytosanitary Measures (ISPM) no. 4: Requirements for the establishment of pest free areas*, Food and Agriculture Organization of the United Nations, Rome, available at <https://www.ippc.int/en/core-activities/standards-setting/ispms/>.

-- -- 1999, *International Standards for Phytosanitary Measures (ISPM) no. 10: Requirements for the establishment of pest free places of production and pest free production sites*, Food and Agriculture Organization of the United Nations, Rome, available at <https://www.ippc.int/en/core-activities/standards-setting/ispms/>.

-- -- 2002, *International Standards for Phytosanitary Measures (ISPM) no. 14: The use of integrated measures in a systems approach for pest risk management*, Food and Agriculture Organization of the United Nations, Rome, available at <https://www.ippc.int/en/core-activities/standards-setting/ispms/>.

-- -- 2007a, *Hyphantria cunea (insect)*, available at <http://www.issg.org/>.

-- -- 2007b, *International Standards for Phytosanitary Measures (ISPM) no. 2: Framework for pest risk analysis*, Food and Agriculture Organization of the United Nations, Rome, available at <https://www.ippc.int/en/core-activities/standards-setting/ispms/>.

-- -- 2009, *International Standards for Phytosanitary Measures (ISPM) no. 28: phytosanitary treatments pt 7: irradiation treatment for fruit flies of the family Tephritidae (generic)*, Food and Agriculture Organization of the United Nations, Rome, available at <https://www.ippc.int/en/core-activities/standards-setting/ispms/>.

-- -- 2013, *International Standards for Phytosanitary Measures (ISPM) no. 11: Pest risk analysis for quarantine pests*, Food and Agriculture Organization of the United Nations, Rome, available at <https://www.ippc.int/en/core-activities/standards-setting/ispms/>.

-- -- 2015a, *International Standards for Phytosanitary Measures (ISPM) no. 5: Glossary of phytosanitary terms*, Food and Agriculture Organization of the United Nations, Rome, available at <https://www.ippc.int/en/core-activities/standards-setting/ispms/>.

-- -- 2015b, *International Standards for Phytosanitary Measures (ISPM) no. 26: Establishment of pest free areas for fruit flies (Tephritidae)*, Food and Agriculture Organization of the United Nations, Rome, available at <https://www.ippc.int/en/core-activities/standards-setting/ispms/>.

Farr, DF, Castlebury, LA, Rossman, AY & Erincik, O 2001, '*Greeneria uvicola*, cause of bitter rot of grapes, belongs in the Diaporthales', *Sydowia*, vol. 53, no. 2, pp. 185-99.

Farr, DF & Rossman, AY 2012, 'Fungal Databases, Systematic Mycology and Microbiology Laboratory, ARS, USDA', available at <http://nt.ars-grin.gov/fungalDATABASES/index.cfm>, accessed 2012.

Farr, DF & Rossman, AY 2013a, '*Coniella diplodiella*', available at http://nt.ars-grin.gov/fungalDATABASES/new_allView.cfm?whichone=all&thisName=Coniella_diplodiella&organismtype=Fungus&fromAllCount=yes, accessed 2013.

- Farr, DF & Rossman, AY 2013b, 'Fungal Databases, Systematic Mycology and Microbiology Laboratory, ARS, USDA', available at <http://nt.ars-grin.gov/fungalDATABASES/index.cfm>, accessed 2013.
- -- 2015, 'Fungal Databases, Systematic Mycology and Microbiology Laboratory, ARS, USDA', available at <http://nt.ars-grin.gov/fungalDATABASES/index.cfm>, accessed 2015.
- Finlay, JR & Teakle, DS 1969, 'The effect of pH on the particle stability of a phosphotungstate-stained *Tobacco necrosis virus*', *Journal of General Virology*, vol. 5, pp. 93-6.
- Firrao, G, Gibbs, K & Stretten, C 2005, 'Short taxonomic guide to the genus '*Candidatus* Phytoplasma'', *Journal of Plant Pathology*, vol. 87, pp. 249-63.
- Flaherty, DL, Christensen, LP, Lanini, WT, Marois, JJ, Phillips, PA & Wilson, LT, (eds) 1992, *Grape pest management*, 2nd edn, University of California, Oakland.
- Follett, PA, Swedman, A & Price, DK 2014, 'Postharvest irradiation treatment for quarantine control of *Drosophila suzukii* (Diptera: Drosophilidae) in fresh commodities', *Journal of Economic Entomology*, vol. 107, no. 3, pp. 964-9.
- Forlani, M & Coppola, V 1977, 'Effetti della frigoconservazione, dell'acido gibberellico e della temperatura sulla germinazione dei vinaccioli di *Vitis vinifera* L. Premessa' (The effects of cold storage, gibberellic acid, and temperature on the germination of *Vitis vinifera* seeds), *Revista di Viticoltura di Enologia*, vol. 30, pp. 445-51.
- Garcia-Arenal, F, Pallas, V & Flores, R 1987, 'The sequence of a viroid from grapevine closely related to severe isolates of *Citrus exocortis viroid*', *Nucleic Acids Research*, vol. 15, pp. 4203-10.
- Gibbs, A & Harrison, B 1976, 'Effects of viruses in plants', in *Plant virology: the principles*, Edward Arnold, London.
- Gildow, F, Damsteegt, VD, Stone, AL, Schneider, W, Luster, D & Levy, L 2004, 'Plum pox in North America: identification of aphid vectors and a potential role for fruit in virus spread', *Phytopathology*, vol. 94, no. 8, pp. 868-74.
- Gonsalves, D 1988, '*Tomato ringspot virus* decline', in *Compendium of grape diseases*, Pearson, RC & Goheen, AC (eds), The American Phytopathological Society Press, St. Paul, Minnesota.
- Gonzalez, RH 1983, *Manejo de plagas de la vid. Publicacions en Ciencias Agrícolas No. 13*, Facultad de Ciencias Agrarias, Veterinarias y Forestales, Universidad de Chile, Santiago.
- Government of Western Australia 2013, 'Western Australia Organism List (WAOL)', available at <http://www.biosecurity.wa.gov.au/western-australian-organism-list-waol>, accessed 2013.
- -- 2014, 'Western Australia Organism List (WAOL)', available at <http://www.biosecurity.wa.gov.au/western-australian-organism-list-waol>, accessed 2015.
- Grasso, EM & Granata, G 2008, 'Disease Note: First report of stem blight on *Metrosideros kermadecensis* induced by *Pestalotiopsis uvicola* in Italy', *Journal of Plant Pathology*, vol. 90, p. 587.
- Grichanov, IY 2009, *Cossus cossus* L.: *goat moth*, Interactive Agricultural Ecological Atlas of Russia and Neighbouring Countries: Economic Plants and their Diseases, Pests and Weeds, available at http://www.agroatlas.ru/en/content/pests/Cossus_cossus/.

- Grichanov, IY & Ovsyannikova, EI 2009, *Hyphantria cunea* Drury, Interactive Agricultural Ecological Atlas of Russia and Neighbouring Countries: Economic Plants and their Diseases, Pests and Weeds, available at <http://www.agroatlas.ru/en/>.
- Guba, EF 1961, *Monograph of Monochaetia and Pestalotia*, Harvard University Press, Cambridge, Massachusetts.
- Gupta, SK & Dhooria, MS 1972, 'Some new records of grape vine mite pests and their predators in India', *Current Science*, vol. 41, no. 22, pp. 824-5.
- Guruprasad, BR, Hegde, SN & Krishna, MS 2010, 'Seasonal and altitudinal changes in population density of 20 species of *Drosophila* in Chamundi hill', *Journal of Insect Science*, vol. 10, no. 123, pp. 1-12.
- Gutierrez, J & Schicha, E 1983, 'The spider mite family Tetranychidae (Acari) in New South Wales', *International Journal of Acarology*, vol. 9, no. 3, pp. 99-116.
- Gyeltshen, J, Hodges, A & Hodges, GS 2008, *Orange spiny whitefly, Aleurocanthus spiniferus (Quaintance) (Insecta: Hemiptera: Aleyrodidae)*, Institute of Food and Agricultural Sciences Extension, University of Florida, available at <http://edis.ifas.ufl.edu/IN618>.
- HAL 2012, *The Australian horticulture statistics handbook 2012*, available at <http://horticulture.com.au/> (pdf 9035 kb).
- Halfeld-Vieira, BA & De Lima Nechet, K 2006, 'Bacterial canker of grapevine in Roraima, Brazil', *Fitopatologia Brasileira*, vol. 31, no. 6, p. 604.
- Hall, BH, McMahon, RL & Wicks, TJ 2002, 'First report of *Sclerotinia sclerotiorum* on grape (*Vitis vinifera*) in South Australia', *Australasian Plant Pathology*, vol. 31, pp. 417-8.
- Harari, AR, Zahavi, T, Gordon, D, Anshelevich, L, Harel, M, Ovadia, S & Dunkelblum, E 2007, 'Pest management programmes in vineyards using male mating disruption', *Pest Management Science*, vol. 63, pp. 769-75.
- Harding, RB & Wicks, TJ 2007, '*Verticillium dahliae* and *Pratylenchus* spp: populations in potato soils and plants in Australia', *Australasian Plant Pathology*, vol. 36, pp. 62-7.
- Harper, SJ, Delmiglio, C, Ward, LI & Clover, GRG 2011, 'Detection of *Tomato black ring virus* by real-time one-step RT-PCR', *Journal of Virological Methods*, vol. 171, pp. 190-4.
- Harris, A, Gibbs, AJ & Gibbs, MJ 2002, *Nepoviruses and their diagnosis in plants: a novel polymerase chain reaction diagnostic test for nepoviruses in nursery stock*, Biosecurity Australia, Department of Agriculture, Fisheries and Forestry, Canberra.
- Harris, AR 1983, 'Distribution of plant parasitic nematodes in vineyards in Central and North-Eastern Victoria', *Australian Journal of Experimental Agriculture and Animal Husbandry*, vol. 23, pp. 221-4.
- Harrison, BD 1958, 'Relationship between Beet Ringspot, Potato Bouquet and Tomato Black Ring Viruses', *Journal of General Microbiology*, vol. 18, pp. 450-60.
- -- 1996, '*Tomato black ring nepovirus*', in *Viruses of plants: descriptions and lists from the VIDE database*, Brunt, AA, Crabtree, K, Dallwitz, MJ, Gibbs, A & Watson, L (eds), CAB International, Wallingford.
- Hattori, I 1969, 'Fruit-piercing moths in Japan', *Japan Agricultural Research Quarterly*, vol. 4, no. 4, pp. 32-6.

- Hauser, M, Gaimari, S & Damus, M 2009, 'Drosophila suzukii new to North America', *Fly Times*, vol. 43, pp. 12-5.
- Hewitt, WB & Pearson, RC 1988, 'Phomopsis cane and leaf spot', in *Compendium of grape diseases*, Pearson, RC & Goheen, AC (eds), The American Phytopathological Society Press, St. Paul, Minnesota.
- Hibben, CR, Bozarth, RF & Reese, J 1979, 'Identification of Tobacco necrosis virus in deteriorating clones of aspen', *Forest Science*, vol. 25, pp. 557-67.
- Hill, DS 1987, *Agricultural insect pests of temperate regions and their control*, University of Cambridge, Cambridge.
- Ho, CC & Chen, WH 1994, 'Infestation of grape berry by spider mites', *Plant Protection Bulletin*, vol. 36, no. 3, pp. 251-6.
- Hossain, MS & Williams, DG 2003, 'Phenology of carpophilus beetle populations (Coleoptera: Nitidulidae, *Carpophilus* spp.) in a fruit dump in northern Victoria', *Australian Journal of Experimental Agriculture*, vol. 43, pp. 1275-9.
- Hren, M, Nikolic, P, Rotter, A, Blejec, A, Terrier, N, Ravnikar, M, Dermastia, M & Gruden, K 2009, 'Bois noir' phytoplasma induces significant reprogramming of the leaf transcriptome in the field grown grapevine', *BMC Genomics*, vol. 10, pp. 460-76.
- Huang, SL, Yan, B, Wei, JG, Yan, WH, Cen, ZL & Yang, T 2007, 'First report of plantain zonate leaf spot caused by *Pestalotiopsis menezesiana* in China', *Australian Plant Disease Notes*, vol. 2, pp. 61-2.
- ICRISAT 1985, 'Proceeding of the international sorghum entomology workshop', ICRISAT Patancheru. Andhra Pradesh, India.
- Ihsanul Huq, M & Nowsher Ali Khan, AZM 2008, 'Epidemiology of *Stemphylium* blight of lentil', *Bangladesh Journal of Scientific and Industrial Research*, vol. 43, no. 4, pp. 513-20.
- India Meteorological Department 2000, *Monthly mean maximum & minimum temperature and total rainfall based upon 1901-2000 data*, available at www.imd.gov.in/.
- Ingels, CA, Bugg, RL, McGourty, GT & Christensen, IP 1998, *Cover cropping in vineyards: a grower's handbook*, Publication 3338 edn, University of California, Oakland.
- INRA 1997, 'HYPPZ (Hypermédia pour la protection des plantes)', available at <http://www.inra.fr/hyppz/pa.htm>, accessed 2012.
- -- 1998, 'HYPPZ (Hypermédia pour la protection des plantes)', available at <http://www.inra.fr/hyppz/pa.htm>, accessed 2015.
- -- 2012, *Vespula germanica*, available at <http://www.inra.fr/hyppz/RAVAGEUR/6vesger.htm>.
- Integrated Pest Management Center 2007, *Potato leafhopper (Empoasca fabae)*, Michigan State University, Michigan.
- International Trade Centre 2015, *Trade map: trade statistics for international business development*, Trade Map, available at <http://www.trademap.org/Index.aspx>.
- Ioriatti, C, Lucchi, A & Varela, LG 2012, 'Grape berry moths in Western European vineyards and their recent movement into the New World', in *Arthropod management in vineyards: pests, approaches, and future directions*, Bostonian, NJ, Vincent, C & Isaacs, R (eds), Springer, New York.

IPPC 2008, *Eradication of grapevine leaf rust (GVLR) from Australia*, International Plant Protection Convention, available at <https://www.ippc.int/en/countries/australia/pestreports/>.

Isaacs, R 2007, *Potato leafhopper*, Michigan State University, available at <http://www.grapes.msu.edu/potatoleafhopper.htm>.

Isaacs, R & van Timmeren, S 2009, *Potato leafhopper control in winegrapes*, Michigan State University, available at <http://dnndev.anr.msu.edu/>.

Ismail, AM, Cirvilleri, G & Polizzi, G 2013, 'Characterisation and pathogenicity of *Pestalotiopsis uvicola* and *Pestalotiopsis clavispora* causing grey leaf spot of mango (*Mangifera indica* L.) in Italy', *European Journal of Plant Pathology*, vol. 135, pp. 619-25.

Iwaizumi, R, Kumagai, M & Katsumata, S 1994, 'Research of infestation to several kinds of fruits by the melon fly, *Bactrocera cucurbitae* (Coquillett) and the oriental fruit fly, *B. dorsalis* (Hendel) (Diptera: Tephritidae)', *Research Bulletin of Plant Protection of Japan*, vol. 30, pp. 93-7.

Jambenal, S 2008, 'Studies on bacterial leaf spot of grape caused by *Xanthomonas campestris* pv. *viticola* (Nayudu) Dye in Northern Karnataka', Master of Science (Agriculture) in Plant Pathology, University of Agricultural Sciences.

Jaramillo, J, Chapman, EG, Vega, FE & Harwood, JD 2010, 'Molecular diagnosis of a previously unreported predator-prey association in coffee: *Karnyothrips flavipes* Jones (Thysanoptera: Phlaeothripidae) predation on the coffee berry borer', *Naturwissenschaften*, vol. 97, no. 3, pp. 291-8.

Jeppson, LR, Keifer, HH & Baker, EW 1975, *Mites injurious to economic plants*, University of California, Berkeley.

Jindal, KK & Sharma, RC 1987, 'Almond leaf scorch: a new disease from India', *FAO Plant Protection Bulletin*, vol. 35, no. 2, pp. 64-5.

Jones, AL 1990, 'Brown rot diseases', in *Compendium of apple and pear diseases*, Jones, AL & Aldwinckle, HS (eds), The American Phytopathological Society Press, St. Paul, Minnesota.

Jones, AT & McGavin, WJ 1996, 'A damaging outbreak of arabis mosaic nepovirus in blackcurrant, the occurrence of other nepoviruses in *Ribes* species, and the demonstration that alfalfa mosaic virus is the cause of interveinal white mosaic in blackcurrant', *Annals of Applied Biology*, vol. 129, no. 1, pp. 47-55.

JSAEZ 1987, *Major insect and other pests of economic plants in Japan*, The Japanese Society of Applied Entomology and Zoology (ed), Japan Plant Protection Association, Japan.

Kaiser, WJ 1980, 'Use of thermotherapy to free potato tubers of Alfalfa mosaic, Potato leaf roll, and Tomato black ring viruses', *Phytopathology*, vol. 70, pp. 1119-22.

Kanzawa, T 1935, *Research into the fruit-fly *Drosophila suzukii* Matsumura (preliminary report)*, Yamanashi Agricultural Experimental Station, Kofu.

-- -- 1939, *Studies on *Drosophila suzukii* Mats*, Yamanashi Agricultural Experimental Station, Japan.

Kapoor, VC 2002, 'Fruit-fly pests and their present status in India', *Proceedings of 6th International Fruit Fly Symposium, Stellenbosch, South Africa*, CRP, South Africa, pp. 23-33.

Kapur, AP 1949, 'On the Indian species of *Rodolia* Mulsiant (Coleoptera-Coccinellidae)', *Bulletin of Entomological Research*, vol. 39, no. 4, pp. 531-8.

- Kassanis, B 1970, *Tobacco necrosis virus*, Association of Applied Biologists: Descriptions of Plant Viruses, available at <http://www.dpvweb.net>.
- Kasy, F 1973, 'Beitrag zur Kenntnis der Familie Stathmopodidae Meyrick, 1913 (Lepidoptera, Gelechioidea)' (Contribution to the knowledge of the family Stathmopodidae Meyrick, 1913 (Lepidoptera, Gelechioidea)), *Tijdschrift voor Entomologie*, vol. 116, no. 13, pp. 227-99.
- Kaur, S & Singh, J 1990, '*Colletotrichum acutatum*: a new threat to chilli crop in Punjab', *Indian Phytopathology*, vol. 43, pp. 108-10.
- Kelly, SE, Shaw, SE & Clark, WT 1995, 'Long-term survival of four dogs with disseminated *Aspergillus terreus* infection treated with itraconazole', *Australian Veterinary Journal*, vol. 72, pp. 311-3.
- Keshavareddy, G, Verghese, A & Reddy, BSL 2007, 'Faunistic survey of scolytids in fruit ecosystems in Karnataka, South India', *Pest Management in Horticultural Ecosystems*, vol. 13, no. 2, pp. 122-7. (Abstract only)
- Kew Royal Botanic Gardens 2014, 'Herb IMI', available at <http://www.herbimi.info/herbimi/home.htm>, accessed 2015.
- Kiri-ganai Research Pty Ltd 2006, *The Australian wine grape industry taking stock and setting directions final report, December 2006*, available at http://www.wgga.com.au/industry/tssd/TSSD%20final%20report_27Dec2006.pdf (2.35 mb).
- Kishi, K, (ed) 1998, *Plant diseases in Japan*, Zenkoku Noson Kyoiku Kyokai, Tokyo.
- Kobayashi, T 2007, *Index of fungi inhabiting woody plants in Japan: host, distribution and literature*, Zenkoku-Noson-Kyoiku Kyokai Publishing Company Ltd, Tokyo.
- Koltunow, AM, Krake, LR & Rezaian, MA 1988, '*Hop stunt viroid* in Australian grapevine cultivars: potential for hop infection', *Australasian Plant Pathology*, vol. 17, no. 1, pp. 7-10.
- Komínek, P 2008, 'Distribution of grapevine viruses in vineyards of the Czech Republic', *Journal of Plant Pathology*, vol. 90, pp. 357-8.
- Krizek, GO 1991, 'Sphingids in photography', *Tropical Lepidoptera*, vol. 2, no. 1, pp. 43-51.
- Krüger, K, De Klerk, A, Douglas-Smit, N, Joubert, J, Pittersen, G & Stiller, M 2011, 'Aster yellows phytoplasma in grapevines: identification of vectors in South Africa', *Bulletin of Insectology*, vol. 64, no. Supplement, pp. S137-8.
- Kulkarni, KA 1971, 'Bionomics of the grape flea beetle *Scelodonta strigicollis* (Motschulsky) (Coleoptera: Chrysomelidae)', *Mysore Journal of Agricultural Sciences*, vol. 5, pp. 308-16.
- Kulkarni, NS, Mani, M & Banerjee, K 2007, *Management of thrips on grapes*, Extension folder no. 13, National Research Centre for Grapes, Pune.
- Kulshrestha, S, Hallan, V, Raikhy, G, Ram, R & Zaidi, AA 2004, '*Strawberry latent ringspot virus* infecting roses in India', *Plant Disease*, vol. 88, no. 1, p. 86.
- Kummuang, N, Diehl, SV, Smith, BJ & Graves, CH, Jnr 1996a, 'Muscadine grape berry rot diseases in Mississippi: disease epidemiology and crop reduction', *Plant Disease*, vol. 80, pp. 244-7.
- Kummuang, N, Smith, BJ, Diehl, SV & Graves, CH, Jnr 1996b, 'Muscadine grape berry rot diseases in Mississippi: disease identification and incidence', *Plant Disease*, vol. 80, no. 3, pp. 238-43.

- Lakra, RK & Dahiya, KK 2000, 'Chemical control of thrips *Rhipiphorothrips cruentatus* Hood (Thripidae: Thysanoptera) infesting grapevines (*Vitis vinifera* L.)', *Haryana Journal of Horticultural Science*, vol. 29, no. 1-2, pp. 52-4.
- Lal, B & Arya, A 1982, 'A soft rot of grapes caused by *Phomopsis viticola*', *Indian Phytopathology*, vol. 35, no. 2, pp. 261-4.
- Lal, SS & Pillai, KS 1981, 'Cassava pests and their control in Southern India', *Tropical Pest Management*, vol. 27, no. 4, pp. 480-91.
- Langrell, SRH, Glen, M & Alfenas, AC 2008, 'Molecular diagnosis of *Puccinia psidii* (guava rust) - a quarantine threat to Australian eucalypt and Myrtaceae biodiversity', *Plant Pathology*, vol. 57, pp. 687-701.
- Lauber, HP & Schuepp, H 1968, 'Der praktische rebbauer' (Grape diseases), *Schweizerische Zeitschrift für Obst*, vol. 104, pp. 550-5.
- Laveau, C, van Helden, M, Darrietort, G, Esmenjaud, D & Demangeat, G 2013, 'First detection of *Tomato black ring virus* (TBRV) in a French vineyard', *Journal International des Sciences de la Vigne et du Vin*, vol. 47, no. 3, pp. 191-4.
- Lee, IM, Davis, RE & Gundersen-Rindal, DE 2000, 'Phytoplasma: phytopathogenic mollicutes', *Annual Review of Microbiology*, vol. 54, pp. 221-55.
- Lee, IM, Gundersen, DE, Davis, RE, Bottner, KD, Marcone, C & Seemüller, E 2004, '*Candidatus Phytoplasma asteris*', a novel phytoplasma taxon associated with aster yellows and related diseases', *International Journal of Systematic and Evolutionary Microbiology*, vol. 54, pp. 1037-48.
- Lee, JC, Bruck, DJ, Curry, H, Edwards, D, Haviland, DR, Van Steenwyk, RA & Yorgey, BM 2011, 'The susceptibility of small fruits and cherries to the spotted-wing drosophila, *Drosophila suzukii*', *Pest Management Science*, vol. 67, pp. 1358-67.
- Leslie, JF, Summerell, BA, Bullock, S & Doe, FJ 2005, 'Description of *Gibberella sacchari* and neotypification of its anamorph *Fusarium sacchari*', *Mycologia*, vol. 97, no. 3, pp. 718-24.
- Leu, LS 1988, 'Rust', in *Compendium of grape diseases*, Pearson, RC & Goheen, AC (eds), The American Phytopathological Society Press, St. Paul, Minnesota.
- Li, SF, Guo, R, Tsuji, M & Sano, T 2006, 'Two grapevine viroids in China and the possible detection of a third', *Plant Pathology*, vol. 55, no. 4, p. 564.
- Li, ZX 2004, *Control of pests and diseases of table grape*, 2nd edn, Jindun, Beijing.
- Lima, MF, Ferreira, MASV, Moreira, WA & Dianese, JC 1999, 'Bacterial canker of grapevine in Brazil', *Fitopatologia Brasileira*, vol. 24, no. 3, pp. 440-3.
- Liotta, G & Mineo, G 1964, 'La *Cryptoblabes gnidiella* Mill. o tignola rigata degli agrumi (Lep. Pyralidae) osservazioni biologiche in Sicilia' (The *Cryptoblabes gnidiella* Mill. or citrus striped moth (Lep. Pyralidae) biological observations in Sicily), *Entomologia Agraria e dell'Osservatorio di Fitopatologia di Palermo*, vol. 5, pp. 155-72.
- Lister, RM & Murant, AF 1967, 'Seed-transmission of nematode-borne viruses', *Annals of Applied Biology*, vol. 59, pp. 49-62.
- Lit, IL, Jnr, Caasi-Lit, M & Calilung, VJ 1998, 'The mealybugs (Pseudococcidae, Coccoidea, Hemiptera) of eggplant (*Solanum melongena* Linn.) in the Philippines', *Philippine Entomologist*, vol. 12, no. 1, pp. 29-41.

- Little, A & Rezaian, MA 2003, 'Grapevine viroids', in *Viroids*, Hadidi, A, Flores, R, Randles, JW & Semancik, JS (eds), Commonwealth Scientific and Industrial Research Organisation, Collingwood.
- Liu, AR, Xu, T & Guo, LD 2007, 'Molecular and morphological description of *Pestalotiopsis hainanensis* sp. nov., a new endophyte from a tropical region of China', *Fungal Diversity*, vol. 24, pp. 23-36.
- Liu, H, Wang, H, Zhao, JY, Yuan, C, Li, WY & Liu, YG 2013, 'Endophytic *Pestalotiopsis* from the main pine species in Sichuan', *Mycosystema*, vol. 32, pp. 15-24.
- Liu, X & Ye, H 2009, 'Effect of temperature on development and survival of *Bactrocera correcta* (Diptera: Tephritidae)', *Scientific Research and Essay*, vol. 4, no. 5, pp. 467-72.
- Locci, R & Quaroni, S 1972, 'Studies on *Coniothyrium diplodiella* I. isolation, cultivation and identification of the fungus', *Rivista de Patologia Vegetale*, vol. 8, no. 1, pp. 59-82.
- Loch, A 2007, *Grapevine pests and their management: Primefact 511*, NSW Department of Primary Industries, available at http://www.dpi.nsw.gov.au/data/assets/pdf_file/0010/110998/Grapevine-pests-and-their-management.pdf (pdf 221 kb).
- Longland, JM & Sutton, TB 2008, 'Factors affecting the infection of fruit of *Vitis vinifera* by the bitter rot pathogen *Greeneria uvicola*', *Phytopathology*, vol. 98, pp. 580-4.
- Maccarone, LD, Sivasithamparam, K, Barbetti, MJ & Jones, RAC 2008, 'Phylogenetic analysis of *Olpidium virulentus*, *Lettuce big-vein associated virus* and *Mirafiori lettuce big-vein virus* isolates from Western Australia', *Plant Health Research Symposium 2008, 24 October 2008*, Australian Plant Pathology Society, Western Australia, pp. 1-6.
- Mackie, AE, Eyres, N & Kumar, S 2005, *Brown rot *Monilinia fructigena**, Factsheet note 181, Department of Agriculture and Food, Western Australia.
- MacLeod, A 2006, *CSL pest risk analysis for *Planococcus lilacinus**, Central Science Laboratory, Sand Hutton.
- Madhusudan, T & Govindu, HC 1985, 'Identification of virus diseases of tobacco in South India', *Indian Journal of Virology*, vol. 1, no. 2, pp. 163-73.
- MAF Biosecurity New Zealand 2009, *Import risk analysis: table grapes (*Vitis vinifera*) from China final*, Ministry of Agriculture and Forestry, Wellington, available at <http://www.biosecurity.govt.nz/files/regs/imports/risk/table-grapes-china-ra.pdf> (pdf 1.68 mb).
- -- 2011, *Rubus (raspberries, blackberries & hybridberries) post-entry quarantine testing manual. January 2011*, Biosecurity New Zealand, Ministry of Agriculture and Forestry, Auckland.
- MAFF 2008, *Information of pests and their risk management: major pests on grape in Japan*, Ministry of Agriculture, Forestry and Fisheries, Japan, Tokyo.
- Maharachchikumbura, SSN, Guo, LD, Chukeatirote, E, Bahkali, AH & Hyde, KD 2011, '*Pestalotiopsis* - morphology, phylogeny, biochemistry and diversity', *Fungal Diversity*, vol. 50, pp. 167-87.
- Mauguashca, F, Ferguson, H, Bahder, B, Brooks, T, O'Neal, S & Walsh, D 2010, *Spotted wing drosophila, 4 October 2010 grape update: injured and ripening fruit may become more attractive: monitoring strongly recommended*, Washington State University Extension, available at http://extension.wsu.edu/swd/Documents/SWDGrapeUpdate10_4_10.pdf (pdf 241 kb).

- Malavolta, VA, Jnr, Almeida, IMG, Sugimori, MH, Ribeiro, IJA, Neto, JR, Pires, EJP & Nogueira, EMC 1999, 'Ocorrência de *Xanthomonas campestris* pv. *viticola* en videira no Brasil' (Occurrence of *Xanthomonas campestris* pv. *viticola* in grape in Brazil), *Summa Phytopathologica*, vol. 25, pp. 262-4.
- Mali, VR, Vyanjane, NT & Ekbote, AU 1977, 'Petunia ringspot virus disease - a new record for India', *Current Science*, vol. 46, pp. 567-8.
- Mani, M 1992, '*Bactrocera correcta* on grapevine in India', *FAO Plant Protection Bulletin*, vol. 40, no. 4, pp. 162-3.
- Mani, M & Thontadarya, TS 1987, 'Record of mealybug species on grapevine in Karnataka', *Current Science*, vol. 56, no. 22, p. 1192.
- Martelli, GP 1978, 'Nematode-borne viruses of grapevine, their epidemiology and control', *Nematologia Mediterranea*, vol. 6, pp. 1-27.
- -- 1999, 'Infectious diseases and certification of grapevines', in *Proceedings of the Mediterranean network on grapevine closteroviruses 1992-1997 and the viruses and virus-like diseases of the grapevine a bibliographic report, 1985-1997*, Centre International de Hautes études Agronomiques Méditerranéennes-Istituto Agronomico Mediterraneo di Bari, Bari.
- Martin, RR & Tzanetakis, IE 2006, 'Characterization and recent advances in detection of strawberry viruses', *Plant Disease*, vol. 90, no. 4, pp. 384-96.
- Mathew, G 1987, 'Insect borers of commercially important stored timber in the state of Kerala, India', *Journal of Stored Products Research*, vol. 23, no. 4, pp. 185-90.
- Mau, RFL & Martin Kessing, JL 1992, *Cyptoblabs gnidiella* (Milliere), Crop Knowledge Master, available at <http://www.extento.hawaii.edu/Kbase/crop/Type/cryptobl.htm>.
- -- 2007, *Bactrocera dorsalis* (Hendel), Crop Knowledge Master, available at http://www.extento.hawaii.edu/kbase/crop/Type/bactro_d.htm.
- Maxwell, A & Scott, JK 2008, 'Pathogens on wild radish, *Raphanus raphanistrum* (Brassicaceae), in south-western Australia - implications for biological control', *Australasian Plant Pathology*, vol. 37, pp. 523-33.
- McCormick, KM & Hollaway, GJ 1999, 'First report of bacterial blight in fenugreek (*Trigonella foenum-graecum*) caused by *Pseudomonas syringae* pv. *syringae*', *Australasian Plant Pathology*, vol. 28, p. 338.
- McGrew, JR 1988, 'Bitter rot', in *Compendium of grape diseases*, Pearson, RC & Goheen, AC (eds), The American Phytopathological Society Press, St. Paul, Minnesota.
- McGrew, JR & Pollack, FG 1988, 'Leaf blight', in *Compendium of grape diseases*, Pearson, RC & Goheen, AC (eds), The American Phytopathological Society Press, St. Paul.
- McKeown, KC 1947, 'Catalogue of the Cerambycidae (Coleoptera) of Australia', *Memoirs of the Australian Museum*, vol. 10, pp. 1-190.
- McLeod, R, Reay, F & Smyth, J 1994, *Plant nematodes of Australia listed by plant and by genus*, New South Wales Agriculture, Rural Industries Research and Development Corporation, Australia.

- Medina-Gaud, S & Franqui, RA 2001, '*Retithrips syriacus* (Mayet), the black vine thrips (Insecta: Thysanoptera: Thripidae) new to Puerto Rico', *Journal of Agriculture of the University of Puerto Rico*, vol. 85, no. 1-2, pp. 85-9.
- Meijerman, L & Ulenberg, SA 2000, *Eurasian Tortricidae: Archips podana*, Arthropods of Economic Importance, available at <http://nlbif.eti.uva.nl/bis/projects.php>.
- -- 2011, *Archips machlopiis* (Meyrick 1912), Arthropods of Economic Importance, available at <http://wbd.etibioinformatics.nl/bis/tortricidae.php?menuentry=soorten&id=127>.
- Mencarelli, F, Bellincontro, A & DiRenzo, G 2005, "*Grape postharvest operations*" with emphasis in developing countries, available at <http://www.fao.org/inpho/content/compend/text/ch34/Ae615e02.htm#TopOfPage>.
- Menzies, R & Brien, J 2002, *AGFACTS: raspberry growing in NSW*, Agfact H3.1.46 (2nd edition), NSW Agriculture, New South Wales, available at http://www.dpi.nsw.gov.au/_data/assets/pdf_file/0020/140249/raspberry-growing-in-nsw.pdf (289 kb pdf).
- Merrin, SJ, Nair, NG & Tarran, J 1995, 'Variation in *Phomopsis* recorded on grapevine in Australia and its taxonomic and biological implications', *Australasian Plant Pathology*, vol. 24, pp. 44-56.
- Meulewaeter, F, Seurinck, J & Van Emmelo, J 1990, 'Genome structure of *Tobacco necrosis virus* strain A', *Virology*, vol. 177, pp. 699-709.
- Meyrick, E 1897, 'Descriptions of Australian Micro-Lepidoptera. XVII. Elachistidae', *Proceedings of the Linnean Society of New South Wales*, vol. 22, pp. 297-435.
- Mickelbart, MV 1996, 'Sapodilla: a potential crop for subtropical climates', in *Progress in new crops*, Janick, J (ed), ASHS Press, Alexandria.
- Migeon, A & Dorkeld, F 2006, 'Spider mites web: a comprehensive database for the Tetranychidae', available at <http://www.montpellier.inra.fr/CBGP/spmweb>, accessed 2010.
- -- 2012, 'Spider mites web: a comprehensive database for the Tetranychidae', available at <http://www1.montpellier.inra.fr/CBGP/spmweb/>, accessed 2012.
- Mijuskovic, M & Badulovic, S 1960, 'Masovna pojava *Celerio lineata* Fabr. var. *livornica* esp. na vinogradima u okolini titograda 1958. Godine' (Mass appearance of *Celerio lineata* Fabr. var. *livornica* Esp. on vines near Titograd in 1958), *Zaštita Bilja*, vol. 61, pp. 21-30.
- Mikusova, P, Santini, A, Ritieni, A, Pavlokin, J & Srobarova, A 2012, 'Berries contamination by microfungi in Slovakia vineyard regions: Impact of climate conditions on microfungi biodiversity', *Revista Iberoamericana de Micología*, vol. 29, no. 3, pp. 126-31.
- Miles, PW 1987, 'Feeding process of Aphidoidea in relation to effects on their food plants.', in *Aphids: their biology, natural enemies and control Volume A*, Minks, AK & Harrewijn, P (eds), Elsevier Science Publishers, Amsterdam.
- Milholland, RD 1991, 'Muscadine grapes: some important diseases and their control', *Plant Disease*, vol. 75, pp. 113-7.
- Millar, JG, Daane, KM, McElfresh, JS, Moreira, JA, Malakar-Kuenen, R, Guillén, M & Bentley, WJ 2002, 'Development and optimization of methods for using sex pheromone for monitoring the mealybug *Planococcus ficus* (Homoptera: Pseudococcidae) in California vineyards', *Journal of Economic Entomology*, vol. 95, no. 4, pp. 706-14.

- Miller, DR & Davidson, JA 2005, *Armored scale insect pests of trees and shrubs (Hemiptera: Diaspididae)*, Cornell University Press, Ithaca & London.
- Miller, DR, Denno, BD & Gimpel, ME 2012, 'Diaspididae: Diaspidinae & Leucaspinae', available at <http://www.sel.barc.usda.gov/scalenet/scalenet.htm>, accessed 2012.
- Miller, DR, Miller, GL & Watson, GW 2002, 'Invasive species of mealybugs (Hemiptera: Pseudococcidae) and their threat to US agriculture', *Proceedings of the Entomological Society of Washington*, vol. 104, no. 4, pp. 825-36.
- Miller, DR, Rung, A, Venable, GL & Gill, RJ 2007, 'Scale insects: identification tools for species of quarantine significance', available at <http://www.sel.barc.usda.gov/ScaleKeys/ScaleInsectsHome/ScaleInsectsHome.html>, accessed 2012.
- Mink, GI 1993, 'Pollen-and seed-transmitted viruses and viroids', *Annual Review of Phytopathology*, vol. 31, pp. 375-402.
- Mishra, B, Prakash, O & Misra, AP 1974, '*Pestalotia menezesiana* on grape berries from India', *Indian Phytopathology*, vol. 27, pp. 257-8.
- Mohanan, C, Ratheesh, N, Nair, LP & Kumar, KCR 2005, 'Disease problems in root trainer forest nurseries in Kerala State and their management', *Proceedings of the 5th Meeting of IUFRO Working Party, Peechi, Kerala, India*, Finnish Forest Research Institute, Helsinki, pp. 7-12.
- Momol, TM, Ritchie, L & Dankers, H 2007, *2007 Florida plant disease management guide: grape (Vitis spp.)*, Institute of Food and Agricultural Sciences Extension, University of Florida, available at <http://edis.ifas.ufl.edu/pdffiles/PG/PG01100.pdf> (pdf 343 kb).
- Moon, H, Woong, K, Choi, M, Kwon, S, Shin, Y, Kim, D & Hwang, C 2008, 'Seasonal occurrences of insect pests in watermelon under greenhouses as affected by cropping season' (in Korean), *Korean Journal of Applied Entomology*, vol. 47, no. 4, pp. 345-52.
- Mordue, JEM 1976, 'Commonwealth Mycological Institute descriptions of fungi and bacteria no. 514: *Pestalotiopsis funerea*', CAB International. Wallingford.
- Mostert, L, Corus, PW & Kang, JC 2001, 'Species of *Phomopsis* and a *Libertella* sp. occurring on grapevines with specific reference to South Africa: morphological, cultural, molecular and pathological characterization', *Mycologia*, vol. 93, no. 1, pp. 146-67.
- Mound, L 2005, *Black vine thrips (Retithrips syriacus)*, PaDIL: Species Factsheet, available at <http://www.padil.gov.au:80/pests-and-diseases/Pest/Main/136401>.
- -- 2009, *Aeolothripidae*, World Thysanoptera, available at http://anic.ento.csiro.au/thrips/identifying_thrips/Aeolothripidae.htm.
- Mundkur, BB & Thirumalachar, MJ 1946, *Revisions of and additions to Indian fungi I*, Mycological Papers No. 16, The Imperial Mycological Institute, Kew.
- Murant, AF 1970, *Tomato black ring virus*, Descriptions of Plant Viruses, available at <http://www.dpvweb.net/dpv/showdpv.php?dpvno=38>.
- -- 1974, *Strawberry latent ringspot virus*, Descriptions of Plant Viruses, available at <http://www.dpvweb.net/dpv/showdpv.php?dpvno=126>.
- -- 1983, 'Seed and pollen transmission of nematode-borne viruses', *Seed Science and Technology*, vol. 11, pp. 973-87.

- -- 1987, 'Raspberry ringspot and associated diseases of *Rubus* caused by raspberry ringspot and tomato black ring viruses', *Agriculture handbook: United States Department of Agriculture, Combine Forest Pest Research and Development Program*, vol. 631, pp. 211-20.
- Nag Raj, TR 1993, *Coelomycetous anamorphs with appendage-bearing conidia*, Mycologue Publications, Ontario.
- Nain, PS, Rishi, N & Bishnoi, SS 1994, 'Profile of viral diseases of cowpea (*Vigna unguiculata*) in Northern India', *Indian Journal of Virology*, vol. 10, no. 2, pp. 128-36.
- Nair, NG 1985, 'Fungi associated with bunch rot of grapes in the Hunter Valley', *Australian Journal of Agricultural Research*, vol. 36, pp. 435-42.
- Nanni, B, Zaccaria, F & Marziano, F 2003, 'First record of brown rot', *Petria*, vol. 13, no. 1-2, pp. 105-14.
- Nascimento, ARP & Mariano, RLR 2004, 'Bacterial canker of grapevine: etiology, epidemiology and control strategies', *Ciência Rural*, vol. 34, no. 1, pp. 301-7.
- Nascimento, ARP, Michereff, SJ, Mariano, RLR & Viana, IO 2006, 'Reaction of grapevine clones to *Xanthomonas campestris* pv. *viticola*, based on epidemiological components of bacterial canker', *Ciência Rural*, vol. 36, no. 1, pp. 1-7.
- Nassar, OA & Ghai, S 1981, 'Taxonomic studies on Tetranychoid mites infesting vegetable and fruit crops in Delhi and surrounding areas.', *Oriental Insects*, vol. 15, no. 4, pp. 333-96.
- Navajas, M, Gutierrez, J, Williams, M & Gotoh, T 2001, 'Synonymy between two spider mite species, *Tetranychus kanzawai* and *T. hydrangeae* (Acari: Tetranychidae), shown by ribosomal ITS2 sequences and cross-breeding experiments', *Bulletin of Entomological Research*, vol. 91, pp. 117-23.
- Nayudu, MV 1972, '*Pseudomonas viticola* sp. nov., incitant of a new bacterial disease of grape vine', *Phytopathology*, vol. 73, pp. 183-6.
- NCBI 2009, '*Necrovirus*', available at <http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=12052>, accessed 2009.
- Nemeth, M 1986, '*Tobacco necrosis virus*', in *Virus, mycoplasma and rickettsia diseases of fruit trees*, Martinus Nijhoff Publishers, Dordrecht.
- NHB 2009, *Grape pests, National Horticulture Board, India*, available at http://nhb.gov.in/bulletin_files/fruits/grape/gra008.pdf (pdf 303 kb).
- -- 2011, *Grape: Diseases*, National Horticulture Board, available at <http://nhb.gov.in/fruits/grape/gra002.pdf> (pdf 451kb).
- Nielsen, AL, Hamilton, GC & Matadha, D 2008, 'Developmental rate estimation and life table analysis for *Halyomorpha halys* (Hemiptera: Pentatomidae)', *Environmental Entomology*, vol. 37, no. 2, pp. 348-55.
- Nielsen, ES & Common, IFB 1991, 'Lepidoptera', in *The insects of Australia: a textbook for students and research workers*, Naumann, ID, Carne, PB, Lawrence, JF, Nielsen, ES, Spradbery, JP, Taylor, RW, Whitten, MJ & Littlejohn, MJ (eds), Melbourne University Press, Carlton.
- Nielsen, ES, Edwards, ED & Rangsi, TV 1996, *Checklist of the Lepidoptera of Australia*, Commonwealth Scientific and Industrial Research Organisation, Collingwood.
- NPQS 2007, *PRA materials of grapes from Korea*, National Plant Quarantine Service, Korea.

- NRC 2013, *Good agricultural practices for production of quality table grapes*, National Research Centre for Grapes, Indian Council of Agricultural Research, Pune.
- NVHSC 2005, *Pest risk analysis for the establishment of a new infestation of phylloxera in Australia as a result of harvest processes*, National Vine Health Steering Committee, South Australia, available at <http://www.gwrdc.com.au/downloads/0708docs/PRAgenericOct05.pdf> (pdf 180 kb).
- -- 2008, *National phylloxera management protocol: definitions of phylloxera management zones*, National Vine Health Steering Committee, South Australia.
- Ocete, R, López, MÁ, Gallardo, A & Arnold, C 2008, 'Comparative analysis of wild and cultivated grapevine (*Vitis vinifera*) in the Basque region of Spain and France', *Agriculture, Ecosystems and Environment*, vol. 123, pp. 95-8.
- Odak, SC & Dhamdhare, SV 1970, 'New record of Hymenopterous parasites of *Sylepta lunalis* Guen (Lepidoptera: Pyralidae), a pest of grapevines', *The Indian Journal of Entomology*, vol. 32, p. 395.
- Office of the Gene Technology Regulator 2003, *Application for licence for dealings involving an intentional release into the environment-field trial of GM grapevines: evaluation of berry colour, sugar composition, flower and fruit development and pollen flow study*, DIR 031/2002 Risk assessment and risk management plan, Department of Health and Ageing, Canberra.
- Ogata, T, Katsumata, H & Takaiwa, K 1999, 'Brown rot of grape berry caused by *Monilia fructigena*' (in Japanese), *Annual Report of the Society of Plant Protection of North Japan*, vol. 50, pp. 97-9.
- Olmo, HP 1976, 'Grapes', in *Evolution of crop plants*, Simmonds, NW (ed), Longmans, London.
- Ono, Y 2000, 'Taxonomy of the *Phakopsora ampelopsidis* species complex on vitaceous hosts in Asia including a new species, *P. euvitis*', *Mycologia*, vol. 92, no. 1, pp. 154-73.
- Ovsyannikova, EI & Grichanov, IY 2009a, *Archips podana Scop.: large fruit-tree tortrix*, Interactive Agricultural Ecological Atlas of Russia and Neighbouring Countries: Economic Plants and their Diseases, Pests and Weeds, available at http://www.agroatlas.ru/en/content/pests/Archips_podana/.
- -- 2009b, *Mamestra brassicae L.*, Interactive Agricultural Ecological Atlas of Russia and Neighbouring Countries: Economic Plants and their Diseases, Pests and Weeds, available at <http://www.agroatlas.ru/en/>.
- Palejwala, VA, Sattur, AP & Modi, VV 1988, 'Strategies for the control of post-harvest spoilage of mangoes by *Penicillium cyclopium*', *Acta Horticulturae (ISHS)*, vol. 231, pp. 688-96.
- Pandotra, VR 1976, 'Occurrence of fruit spot disease of grapes in Jammu', *Indian Journal of Mycology and Plant Pathology*, vol. 6, no. 2, p. 182.
- Park, JY, Lee, W, Song, DU, Seong, KY, Cho, BH & Kim, KC 1997, 'Leaf blight of kiwifruit caused by *Pestalotiopsis menezesiana* (Bress. & Torr.) Bissett and its ecopathology' (in Korean), *Korean Journal of Plant Pathology*, vol. 13, no. 1, pp. 22-9.
- Parkinson, N, Cowie, C, Heeney, J & Stead, D 2009, 'Phylogenetic structure of *Xanthomonas* determined by comparison of *gyrB* sequences', *International Journal of Systematic Bacteriology*, vol. 59, pp. 264-74.

- Pathak, AK & Verma, KS 2009, 'Aero-bacteriological study of vegetables market at Jabalpur', *Iranian Journal of Environmental Health Science & Engineering*, vol. 6, no. 3, pp. 187-94.
- Pavan, F, Stefanelli, G, Villani, A, Gasparinetti, P, Cofussi, G, Mucignat, D, Bernard, D & Mutton, P 1998, 'Danni da *Empoasca vitis* (Gothe) (Homoptera: Cicadellidae) in vigne ti dell'Italia nord-orientale e soglie d'intervento' (Damage caused by *Empoasca vitis* (Gothe) (Homoptera: Cicadellidae) in vineyards of North-Eastern Italy and intervention thresholds), *Frutsula Entomologica*, vol. 21, no. 34, pp. 109-24.
- Pearson, RC & Goheen, AC 1988, *Compendium of grape diseases*, Pearson, RC & Goheen, AC (eds), The American Phytopathological Society Press, St. Paul, Minnesota.
- Peixoto, AR, Mariano, RLR, Moreira, JOT & Viana, IO 2007, 'Alternative hosts of *Xanthomonas campestris* pv. *viticola*', *Fitopatologia Brasileira*, vol. 32, no. 2, pp. 161-4.
- Penrose, LJ, Nicholls, MR & Koffmann, W 1984, 'Apple fruit rot caused by *Trichoderma harzianum*', *Australasian Plant Pathology*, vol. 13, no. 3, pp. 46-7.
- Persley, D & Magarey, P 2009, 'Grape', in *Diseases of fruit crops in Australia*, Cooke, T, Persley, D & House, S (eds), CSIRO Publishing, Collingwood.
- Persley, DM, Thomas, JE & Sharman, M 2006, '*Tospoviruses*: an Australian perspective', *Australasian Plant Pathology*, vol. 35, pp. 161-80.
- Pethybridge, SJ, Hay, FS & Groom, T 2003, 'Seasonal fluctuations in fungi associated with *Pyrethrum* foliage in Tasmania', *Australasian Plant Pathology*, vol. 32, pp. 223-30.
- Petrovic, T, Burgess, LW, Cowle, I, Warren, RA & Harvey, PR 2013, 'Diversity and fertility of *Fusarium sachari* from wild rice (*Oryza australiensis*) in Northern Australia, and pathogenicity tests with wild rice, rice, sorghum and maize', *European Journal of Plant Pathology*, vol. 136, no. 4, pp. 773-88.
- Pfeiffer, DG 2009, *Climbing cutworms in grapes*, Mid-Atlantic Regional Fruit Loop, available at www.virginiafruit.ento.vt.edu/cutwormsgrape.html.
- PGIBSA 2009, *Phylloxera management zones of Australia*, Phylloxera and Grape Industry Board of South Australia, South Australia.
- Phadke, CH, Rao, VG & Pawar, SK 1978, 'Natural outbreak of the muscardine fungus *Nomuraea rileyi* (Farlow) Samson on leaf-eating caterpillar, *Spodoptera exigua* Hb. in Maharashtra', *Current Science*, vol. 47, no. 13, p. 476.
- Pitkin, BR 1976, 'A revision of the Indian species of Haplothrips and related genera (Thysanoptera: Phlaeothripidae).', *Bulletin of the British Museum of Natural History, Entomology*, vol. 34, pp. 223-80.
- Pittaway, AR & Kitching, IJ 2006, 'Sphingidae of the eastern Palaearctic', available at <http://tpittaway.tripod.com/china/china.htm>, accessed 2006.
- -- 2012, 'Sphingidae of the Eastern Palaearctic', available at <http://tpittaway.tripod.com/china/china.htm>, accessed 2012.
- Plant Health Australia 2001, 'Australian plant pest database, online database', available at www.planthealthaustralia.com.au/appd, accessed 2013.
- -- 2009a, *Industry biosecurity plan for the strawberry industry*, Plant Health Australia Limited, Canberra.

- -- 2009b, *Industry biosecurity plan for the viticulture industry*, Version 2.0, Plant Health Australia, Canberra.
- -- 2010, *National Plant Biosecurity Status Report 2010*, Plant Health Australia, Canberra.
- -- 2011, *Industry biosecurity plan for the vegetable industry*, Plant Health Australia Limited, Canberra.
- -- 2013a, *Industry biosecurity plan for the potato industry*, Plant Health Australia Limited, Canberra.
- -- 2013b, *Industry biosecurity plan for the Rubus industry*, Version 1.0 October 2013, Plant Health Australia Limited, Canberra.
- -- 2014, *Strawberries*, available at <http://www.planthealthaustralia.com.au/industries/strawberries/>.
- Pons, N & Sutton, BC 1988, *Cercospora and similar fungi on yams (Dioscorea species)*, Mycological Papers No. 160, CAB International, Wallingford.
- Poole, MC 2010, *An annotated catalogue of insect and allied species associated with Western Australian agriculture and related industries: perennial draft, July 2010*, Department of Agriculture and Food, Government of Western Australia.
- Poole, MC, Wood, CE, Lanoiselet, V, Tuten, SJ & Hammond, NE 2011, *Final pest risk analysis report for the importation of fresh summerfruit from South Australia, Tasmania, New South Wales and Victoria into Western Australia*, Department of Agriculture and Food, Western Australia.
- Prakash, O & Raoof, MA 1985, 'New records of fungi on leaves and twigs of mango (*Mangifera indica*)', *Indian Journal of Plant Pathology*, vol. 3, no. 2, pp. 243-4.
- Prasad, SK 1960, 'Effect of insecticides of Potato leaf hopper, *Empoasca fabae* (Harris): abundance and yield of potatoes', *Indian Journal of Entomology*, vol. 22, no. 4, pp. 283-6.
- Pratt, C 1988, 'Grapevine structure and growth stages', in *Compendium of grape diseases*, Pearson, RC & Goheen, AC (eds), The American Phytopathological Society Press, St. Paul, Minnesota.
- Pung, H & Cox, P 1999, 'The development of crown rot disease on carrots', *12th Biennial Australasian Plant Pathology Conference 27-30 September 1999, Devonport, Tasmania, Serve-Ag Research, Tasmania*.
- Punithalingam, E 1968, *Commonwealth Mycological Institute descriptions of fungi and bacteria no. 173: Physopella ampelopsidis*, CAB Abstracts plus, available at <http://cababstractplus.org>.
- Puttarudriah, M & Appanna, GB 1955, 'Two new hosts of *Helopeltis antonii* Signoret in Mysore', *Indian Journal of Entomology*, vol. 17, pp. 391-2.
- Puttarudriah, M, Kataoihallimath, SS & Chandrasekhar, B 1961, 'A note on the occurrence of *Cacoecia micaceana* Walker as a pest on grape vine', *Mysore Agricultural Journal*, vol. 35, no. 4, pp. 228-32.
- Queensland Government 2014, *Introduction into Queensland of grape phylloxera, grape plants, grape plant products, vineyard appliances or soil which has been in contact with a grape plant*, Queensland Government, Brisbane.
- Rajak, RC & Pandey, AK 1985, 'Fungi from Jabalpur', *Indian Journal of Mycology and Plant Pathology*, vol. 15, no. 2, pp. 186-94.

- Ramachandraiah, M, Venkatarathnam, P & Sulochana, CB 1979, 'Tobacco necrosis virus: occurrence in India', *Plant Disease Reporter*, vol. 63, no. 11, pp. 949-51.
- Ramachandran, P, Agrawal, J, Roy, A, Ghosh, DK, Das, DR & Ahlawat, YS 2005, *First record of a Hop stunt viroid variant on Nagpur mandarin and Mosambi sweet orange trees on rough lemon and Rangpur lime rootstocks*, New Disease Reports, available at <http://www.bspp.org.uk/publications/new-disease-reports/jan2005/2004-85.asp>.
- Ramachandran, P, Pandey, PK, Ahlawat, YS, Varma, A & Kapur, SP 1993, 'Viroid diseases of citrus in India', *Proceedings of the 12th Conference for the International Organisation of Citrus Virologists, Dehli, India, 23-27 November, 1992, Dehli*, University of California, Riverside, pp. 438-40.
- Ramsdell, DC & Milholland, RD 1988, 'Black rot', in *Compendium of grape diseases*, Pearson, RC & Goheen, AC (eds), The American Phytopathological Society Press, St. Paul, Minnesota.
- Rana, T, Negi, A, Dhir, S, Thockchom, T, Chandel, V, Walia, Y, Singh, RM, Ram, R, Hallan, V & Zaidi, AA 2011, 'Molecular diagnosis of apple virus and viroid pathogens from India', *Archives of Phytopathology and Plant Protection*, vol. 44, no. 6, pp. 505-12.
- Rather, AQ 1999, 'Mites associated with viticulture in India with a key for their identification', *Journal of Acarology*, vol. 15, no. 1-2, pp. 18-24.
- Rawal, RD 1998, 'Management of fungal diseases in tropical fruits', in *Tropical fruits in Asia: diversity, maintenance, conservation and use*, Arora, RK & Rao, VR (eds), International Plant Genetic Resources Institute, Bangalore.
- Reddy, DJ 2006, 'Estimation of avoidable losses due to pests of grapevine', *Indian Journal of Agricultural Research*, vol. 40, no. 4, pp. 282-5.
- Reddy, K 2011, *Division of Plant Pathology: research achievements*, Indian Institute of Horticultural Research, available at <http://www.iihr.res.in/frmDivisionDetails.aspx?id=31>.
- Reddy, MS & Reddy, KRC 1983, 'Greeneria fruit rot: an endemic disease of grape in India', *Indian Phytopathology*, vol. 36, no. 1, pp. 110-4.
- Richardson, FJ, Richardson, RG & Shepherd, RCH 2006, *Weeds of the South-East: an identification guide for Australia*, R.G. and F.J. Richardson, Victoria.
- Ridings, WH & Clayton, CN 1970, '*Melanconium fuliginum* and the bitter rot disease of grape', *Phytopathology*, vol. 60, no. 8, pp. 1203-11.
- Rishi, N 2009, 'Significant plant virus diseases in India and a glimpse of modern disease management technology', *Journal of General Plant Pathology*, vol. 75, pp. 1-18.
- Robert, V, Stegehuis, G & Stalpers, J 2009, 'The mycobank engine and related databases', available at www.mycobank.org, accessed 2009.
- Roberts, SJ, Hiltunen, LH, Hunter, PJ & Brough, J 1999, 'Transmission from seed to seedling and secondary spread of *Xanthomonas campestris* pv. *campestris* in brassica transplants: effects of dose and watering regime', *European Journal of Plant Pathology*, vol. 105, no. 9, pp. 879-89.
- Robinson, GS, Ackery, PR, Kitching, IJ, Beccaloni, GW & Hernández, LM 2010, 'HOSTS: a database of the world's lepidopteran hostplants', available at <http://www.nhm.ac.uk/research-curation/research/projects/hostplants/>, accessed 2012.

- Robinson, GS, Tuck, KR & Shaffer, M 1994, *A field guide to the smaller moths of South-East Asia*, Malasian Nature Society, Kuala Lumpur.
- Rochon, D, Kakani, K, Robbins, M & Reade, R 2004, 'Molecular aspects of plant virus transmission by Olpidium and Plasmodiophorid vectors', *Annual Review of Phytopathology*, vol. 42, pp. 211-41.
- Rodrigues Neto, J, Destéfano, SAL, Rodrigues, LMR, Souza Peloso, D & da Cruz Oliveira, L, Jnr 2011, 'Grapevine bacterial canker in the State of São Paulo, Brazil: detection and eradication', *Tropical Plant Pathology*, vol. 36, no. 1, pp. 42-4.
- Rose, HS & Pooni, HS 2004, 'Taxonomic studies on the family Tortricidae (Tortricoidae: Lepidoptera) from North-West India: I. Tribe Archipini (Tortricinae)', *Entomon*, vol. 29, no. 2, pp. 137-81.
- Roychoudhury, N & Joshi, KC 2011, 'Effect of ivermectin on larval mortality and reproductive potential of *Albizia* foliage feeder, *Spirama retorta* Cramer (Lepidoptera: Noctuidae)', *Journal of Pest Science*, vol. 84, pp. 117-22.
- Rüdel, M 1985, 'Grapevine damage induced by particular virus-vector combinations', *Phytopathologia Mediterranea*, vol. 24, pp. 183-5.
- Ryan, RP, Vorhölter, FJ, Potnis, N, Jones, JB, Van Sluys, MA, Bogdanove, AJ & Dow, JM 2011, 'Pathogenomics of *Xanthomonas*: understanding bacterium-plant interactions', *Nature Reviews Microbiology*, vol. 9, pp. 344-55.
- Saeki, K, Takahashi, Y, Oh-oka, H, Umeoka, T, Oda, Y & Fukuyama, K 2001, 'Primary structure and phylogenetic analysis of the coat protein of a toyama isolate of *Tobacco necrosis virus*', *Bioscience, Biotechnology and Biochemistry*, vol. 65, no. 3, pp. 719-24.
- Saguez, J, Lasnier, J & Vincent, C 2013, 'First record of *Drosophila suzukii* in Quebec vineyards', *Journal International des Sciences de la Vigne et du Vin*, vol. 47, no. 1, pp. 69-72.
- Sakalidis, ML, Hardy, GES & Burgess, TI 2011, 'Endophytes as potential pathogens of the baobab species *Adansonia gregorii*: a focus on the Botryosphaeriaceae', *Fungal Ecology*, vol. 4, pp. 1-14.
- Samad, A, Ajayakumar, PV, Gupta, MK, Shukla, AK, Darokar, MP, Somkuwar, B & Alam, M 2008, 'Natural infection of periwinkle (*Catharanthus roseus*) with *Cucumber mosaic virus*, subgroup 1b', *Australasian Plant Disease Notes*, vol. 3, pp. 30-4.
- Sampson, PJ & Walker, J 1982, *An annotated list of plant diseases in Tasmania*, Department of Agriculture, Tasmania.
- Sánchez, EC, Garcia, CFO, de la Cruz, MT, Muñoz, CDB, Cruz, MDR & Espinoza, LDL 2011, 'Etiología de la Mancha Acuosa de la Jamaica (*Hibiscus sabdariffa*) en Tabasco, México' (Etiology of Roselle (*Hibiscus sabdariffa*) watery spot in Tabasco, Mexico), *Revista Mexicana de Fitopatología*, vol. 29, no. 2, pp. 165-7.
- Sangalang, AE, Burgess, LW, Backhouse, D, Duff, J & Wurst, M 1995, 'Mycogeography of *Fusarium* species in soils from tropical, arid and mediterranean regions of Australia', *Mycological Research*, vol. 99, no. 5, pp. 523-8.
- Sano, T 2003, '*Hop stunt viroid*', in *Viroids*, Hadidi, A, Flores, R, Randles, JW & Semancik, JS (eds), Commonwealth Scientific and Industrial Research Organisation, Collingwood.
- Sarbhoy, AK, Lal, G & Varshney, JL 1975, *Fungi of India, 1967-71*, Navyug Traders, New Delhi.

- Sasmal, S & Kulshreshtha, JP 1985, '*Cryptoblabes gnidiella*, an azolla pest on rice in India', in *International Rice Research Newsletter*, International Rice Institute, Manila, Philippines.
- Satarkar, VR, Krishnamurthy, SV, Faleiro, JR & Verghese, A 2009, 'Spatial distribution of major *Bactrocera* fruit flies attracted to methyl eugenol in different ecological zones of Goa, India', *International Journal of Tropical Insect Science*, vol. 29, no. 4, pp. 195-201.
- Schwinghamer, MW, Schilg, M, Moore, K, Knights, T, Bambach, R, Southwell, R, Kumari, S, Srivastava, M & Wratten, K 2007, *Virus situation in chickpea, faba beans and canola*, Grains Research and Development Corporation, available at http://www.grdc.com.au/director/events/onfarmtrials?item_id=4829BB0CA9AC5A109CF7A53B484E5905&pageNumber=2.
- Scott, DH & Ink, DP 1950, 'Grape seed germination experiments', *Proceedings of the American Society for Horticultural Science*, vol. 56, pp. 134-9.
- Sergeeva, V 2010, 'Grapevine and olive diseases in Australia', *Petria*, vol. 20, no. 2, pp. 99-100.
- Sergeeva, V, Nair, T & Spooner-Hart, RN 2001, 'Fungi recorded on grapevines during the course of an industry service on *Botrytis* monitoring and fungicide resistance', *Australian and New Zealand Grapegrower and Winemaker*, vol. Annual Technical Issue, pp. 7-11.
- Sergeeva, V, Priest, M & Nair, NG 2005, 'Species of *Pestalotiopsis* and related genera occurring on grapevines in Australia', *Australasian Plant Pathology*, vol. 34, pp. 255-8.
- Sergeeva, V, Tesoriero, L, Spooner-Hart, R & Nair, N 2005, 'First report of *Macrophomina phaseolina* on olives (*Olea europaea*) in Australia', *Australasian Plant Pathology*, vol. 34, pp. 273-4.
- Sforza, R, Boudon-Padieu, E & Greif, C 2003, 'New mealybug species vectoring *Grapevine leafroll associated viruses-1 and -3 (GLRaV-1 and -3)*', *European Journal of Plant Pathology*, vol. 109, pp. 975-81.
- Sharma, HC & Agarwal, DK 1997, 'A new anthracnose disease of grape vine (*Vitis vinifera*) in India', *Indian Phytopathology*, vol. 50, no. 1, p. 150.
- Sharma, HC & Bhardwaj, SS 1999, 'Canker and foliage diseases of apple', in *Diseases of horticultural crops: fruit*, Verma, LR & Sharma, RC (eds), Indus Publishing Company, New Dehli.
- Sharma, HC, Singh, F & Nwanze, KF 1997, *Plant resistance to insects in sorghum*, International Crops Research Institute for the Semi-Arid Tropics, India.
- Sharma, RL & Kaul, JL 1989, 'Incidence of brown rot (*Monilinia* spp.) of apple in Himachal Pradesh', *Indian Journal of Mycology and Plant Pathology*, vol. 19, no. 2, pp. 208-11.
- Sharma, S, Kaushik, N & Kaushik, JC 1997, 'Root rot of *Prosopis cineraria* caused by *Fusarium solani*: a new host record', *Indian Phytopathology*, vol. 50, no. 2, p. 300.
- Shekhawat, GS, Gadewar, AV & Chakrabarti, SK 1999, 'Bacterial diseases of potato in India', in *Diseases of horticultural crops: fruit*, Verma, LR & Sharma, RC (eds), Indus Publishing Company, New Dehli.
- Shepherd, CJ & Pratt, BH 1973, 'Separation of two ecotypes of *Phytophthora drechsleri* Tucker occurring in Australian native forests', *Australian Journal of Biological Sciences*, vol. 26, pp. 1095-107.

- Shikhamany, SD 2001, 'Grape production in India', in *Grape production in the Asia-Pacific region*, Papademetriou, MK & Dent, FJ (eds), Food and Agriculture Organization for the United Nations, Bangkok.
- Shivas, RG 1989, 'Fungal and bacterial diseases of plants in Western Australia', *Journal of the Royal Society of Western Australia*, vol. 72, no. 1-2, pp. 1-62.
- Shreemali, PG 1973, 'A new disease of *Artabotrys hexapetalous*', *Indian Phytopathology*, vol. 26, no. 3, pp. 605-7.
- Shukla, RP & Tandon, PL 1984, 'Insect pests on custard apple', *FAO Plant Protection Bulletin*, vol. 32, no. 1, p. 31.
- Sidhu, AK, Chandra, K & Pathania, PC 2010, *A check-list of microlepidoptera of India (Part I: Family Pterophoridae)*, Zoological Survey of India online version, available at <http://zsi.gov.in/checklist/Family%20Pterophoridae.pdf> (pdf 143 kb).
- Silva, AMF, de Menezes, EF, de Souza, EB, de Melo, NF & Mariano, RDLR 2012, 'Survival of *Xanthomonas campestris* pv. *viticola* in infected tissue of grapevine', *Revista Brasileira de Fruticultura*, vol. 34, no. 3, pp. 757-65.
- Silva, EB & Mexia, A 1999, 'The pest complex *Cryptoblabes gnidiella* (Millière) (Lepidoptera: *Pyalidae*) and *Planococcus citri* (Risso) (Homoptera: *Pseudococcidae*) on sweet orange groves (*Citrus sinensis* (L.) Osbeck) in Portugal: interspecific association', *Boletim de Sanidad Vegetal, Plagas*, vol. 25, pp. 89-98.
- Simmonds, JH 1966, *Host index of plant diseases in Queensland*, Department of Primary Industries, Brisbane.
- Singh, BK & Bhatt, M 1988, 'A preliminary report on the Drosophilidae of Kumaun region with description of two new species and three new records', *Oriental Insects*, vol. 22, pp. 147-61.
- Singh, BK & Negi, NS 1989, 'Drosophilidae of Garhwal region with the description of one new species', *Proceedings of the Zoological Society Calcutta*, vol. 40, pp. 19-26.
- Singh, DV & Sinch, RS 1966, 'Occurrence of *Coniella diplodiella* (Speg.) Petrak and Sydow on *Geranium* sp.', *Science and Culture*, vol. 32, no. 10, p. 504.
- Singh, I & Chohan, JS 1974, 'Seed-borne fungi on cowpea (*Vigna sinensis*)', *Phytopathological Notes*, vol. 27, pp. 239-40.
- Singh, OT, Chakravorty, J & Varatharajan, R 2010, 'Entomofauna of Kane Wildlife Sanctuary, Arunachal Pradesh, northeastern India', *Journal of Threatened Taxa*, vol. 2, no. 13, pp. 1392-400.
- Singh, RP, Ready, KFM & Nie, X 2003, 'Biology', in *Viroids*, Hadidi, A, Flores, R, Randles, JW & Semancik, JS (eds), Commonwealth Scientific and Industrial Research Organisation, Collingwood.
- Singh, S, Nagaich, BB, Arai, K & Khurana, SMP 1983, 'Clover phyllody in Simla hills', *Phytopathological Notes*, vol. 36, pp. 771-5.
- Singh, SK, Motilal, SC, Chand, R & Ojha, KL 1999, 'Grapevine diseases and their management', in *Diseases of horticultural crops: fruits*, Verma, LR & Sharma, RC (eds), M.L. Gidwani, Indus Publishing Company, New Delhi.
- Singh, SN 1961, 'Germination of grape (*Vitis vinifera* L.) hybrid seeds by chilling', *Current Science*, vol. 30, p. 62.

- Singh, YP & Singh, DK 1995, 'Bionomics of *Cryptoblabes gnidiella* Miller A, a pest of sorghum', *Advances in Agricultural Research in India*, vol. 3, pp. 119-29.
- Smith, D 2012, *Bitter rot of grapes*, eXtension, available at <http://www.extension.org/pages/31135/bitter-rot-of-grapes/print/>.
- Smith, IM, Dunez, J, Lelliot, RA, Phillips, DH & Archer, SA 1988, *European handbook of plant diseases*, Smith, IM, Dunez, J, Lelliot, RA, Phillips, DH & Archer, SA (eds), Blackwell Scientific Publications, Oxford.
- Snare, L 2006, *Pest and disease analysis in hazelnuts*, Project no: NT05002, Department of Primary Industries, New South Wales.
- Sohi, BS, Bindra, OS & Sohi, AS 1975, 'Effectiveness of insecticidal sprays against grape-vine Cicadellid, *Arboridia (Arborifera) viniferata* Sohi and Sandhu in the Punjab, India', *The Indian Journal of Horticulture*, vol. 32, no. 1-2, pp. 68-70.
- Srivastava, RP 1997, *Mango insect pest management*, International Book Distributing Co., Lucknow, India.
- Srivastava, RP & Verghese, A 1985, 'Record of a new mealy bug, *Perissopneumon ferox* Newstead (Margarodidae: Homoptera) on mango from Uttar Pradesh, India', *Entomologica*, vol. 10, no. 2, pp. 184-5.
- Stace-Smith, R 1984, *Tomato ringspot virus*, Descriptions of Plant Viruses, available at <http://www.dpvweb.net/dpv/showdpv.php?dpvno=290>.
- -- 1985, *Tobacco ringspot virus*, Descriptions of Plant Viruses, available at <http://www.dpvweb.net/dpv/showdpv.php?dpvno=309>.
- Steel, CC 2007, *Management of non-Botrytis bunch rots: final report to Grape and Wine Research and Development Corporation, Hunter Valley Vineyard Association, Mid-North Coast Winegrape Growers Associations*, National Wine and Grape Industry Centre, Wagga Wagga.
- Steel, CC, Greer, LA & Savocchia, S 2007, 'Studies on *Colletotrichum acutatum* and *Greenaria uvicola*: two fungi associated with bunch grapes in sub-tropical Australia', *Australian Journal of Grape and Wine Research*, vol. 13, pp. 23-9.
- -- 2012, 'Grapevine inflorescences are susceptible to the bunch rot pathogens, *Greeneria uvicola* (bitter rot) and *Colletotrichum acutatum* (ripe rot)', *European Journal of Plant Pathology*, vol. 133, pp. 773-8.
- Stellmach, G 1970, 'Tomato black ring in *Vitis*', in *Virus diseases of small fruits and grapevines*, Frazier, NW, Fulton, JP, Thresh, JM, Converse, RH, Varney, EH & Hewitt, WB (eds), University of California, Berkeley.
- Stobbs, LW & van Schagen, JG 1984, 'Occurrence of *Tomato black ring virus* on grapevine in Southern Ontario', *Canadian Plant Disease Survey*, vol. 64, no. 1, pp. 3-4.
- Streten, C & Gibb, KS 2006, 'Phytoplasma diseases in sub-tropical and tropical Australia', *Australasian Plant Pathology*, vol. 35, pp. 129-46.
- Sugimoto, S 1994, 'Scale insects intercepted on banana fruits from Mindanao Island, The Philippines (Coccoidea: Homoptera)' (in Japanese), *Research Bulletin of the Plant Protection Service, Japan*, vol. 30, no. 1, pp. 115-21.

- Suhag, LS, Kaushik, JC & Duhan, JC 1982, 'Etiology and epidemiology of fungal foliar diseases on grapevine', *Indian Journal of Mycology and Plant Pathology*, vol. 12, no. 2, pp. 191-7.
- Summerell, BA, Leslie, JF, Liew, ECY, Laurence, MH, Bullock, S, Petrovic, T, Bentley, AR, Howard, CG, Peterson, SA, Walsh, JL & Burgess, LW 2011, 'Fusarium species associated with plants in Australia', *Fungal Diversity*, vol. 46, pp. 1-27.
- Suresh, S & Mohanasundaram, M 1996, 'Coccoid (Coccoidea: Homoptera) fauna of Tamil Nadu, India', *Journal of Entomological Research*, vol. 20, no. 3, pp. 233-74.
- Sutton, BC 1980, *The Coelomycetes: fungi imperfecti with pycnidia, acervuli, and stromata*, Commonwealth Mycological Institute, Kew.
- Sutton, BC & Gibson, IAS 1977, *Commonwealth Mycological Institute descriptions of fungi and bacteria no. 538: Greeneria uvicola*, I.M.I. Descriptions of Pathogenic Fungi and Bacteria, available at <http://cababstractsplus.org/DMPD/Index.asp>.
- Sutton, BC & Waterston, JM 1964, *Commonwealth Mycological Institute descriptions of pathogenic fungi and bacteria no. 82: Coniella diplodiella*, CABI Bioscience, available at <http://www.cabi.org/dfb/>.
- Swapna, S, Syed, A & Krishnappa, M 2008, 'Diversity of macrofungi in semi-evergreen and moist deciduous forest of Shimoga District-Karnataka, India', *Journal of Mycology and Plant Pathology*, vol. 38, no. 1, pp. 21-6.
- Swirski, E, Izhar, Y, Wysoki, M, Gurevitz, E & Greenberg, S 1980, 'Integrated control of the long-tailed mealybug, *Pseudococcus longispinus* (Homoptera: Pseudococcidae), in avocado plantations in Israel', *Entomophaga*, vol. 25, no. 4, pp. 415-26.
- Tai, FL 1979, *Sylloge fungorum sinicorum*, Science Press, Academia Sinica, Peking.
- Tandon, PL & Verghese, A 1987, 'New insect pests of certain fruit crops', *Indian Journal of Horticulture*, vol. 44, no. 1-2, pp. 121-2.
- Tashiro, N 1992, 'Girdling of grapevine (cv. "Kyoho") canes by *Greeneria uvicola* (Syn. *Melanconium fuligineum*) in a plastic green house' (in Japanese), *Kyushu Agricultural Research*, vol. 54, p. 88.
- Taylor, A 2012, *Know your grapevine bunch rots*, Farmnote no. 528, Department of Agriculture and Food, Western Australia, available at http://www.agric.wa.gov.au/PC_95052.html?s=1001.
- Taylor, A, St.J.Hardy, GE, Wood, P & Burgess, T 2005, 'Identification and pathogenicity of *Botryosphaeria* species associated with grapevine decline in Western Australia', *Australasian Plant Pathology*, vol. 34, pp. 187-95.
- Taylor, CE & Brown, DJF 1997, *Nematode vectors of plant viruses*, CAB International, Wallingford.
- Teakle, DS 1988, '*Tobacco necrosis necrovirus*', in *Viruses of plants in Australia*, Büchen-Osmond, C, Crabtree, K, Gibbs, A & McLean, G (eds), Australian National University Printing Service, Canberra.
- Toda, MJ 1991, 'Drosophilidae (Diptera) in Myanmar (Burma) vii. the *Drosophila melanogaster* species-group, excepting the *D. montium* species sub-group', *Oriental Insects*, vol. 25, pp. 69-94.
- Tomlinson, JA, Faithful, EM, Webb, MJW, Fraser, RSS & Seeley, ND 1983, 'Chenopodium necrosis: a distinctive strain of *Tobacco necrosis virus* isolated from river water', *Annals of Applied Biology*, vol. 102, pp. 135-47.

- Tostes, GO, Araujo, JSP, Farias, ARG, Frade, DAR & Olivares, FL 2014, 'Detection and cellular localization of *Xanthomonas campestris* pv. *viticola* in seeds of commercial 'Red Globe' grapes', *Tropical Plant Pathology*, vol. 39, no. 2, pp. 134-40.
- Trindade, LC, Lima, MF & Ferreira, MASV 2005, 'Molecular characterization of Brazilian strains of *Xanthomonas campestris* pv. *viticola* by rep-PCR fingerprinting', *Fitopatologia Brasileira*, vol. 30, no. 1, pp. 46-54.
- Trindade, LC, Marques, E, Lopes, DB & Ferreira, MASV 2007, 'Development of a molecular method for detection and identification of *Xanthomonas campestris* pv. *viticola*', *Summa Phytopathologica*, vol. 33, no. 1, pp. 16-23.
- Tuck, KR 1990, 'A taxonomic revision of the Malaysian and Indonesian species of *Archips* Hübner (Lepidoptera: Tortricidae)', *Entomologica Scandinavica*, vol. 21, no. 2, pp. 179-96.
- Ullasa, BA & Rawal, RD 1986, 'Studies on American rot of grapes due to *Greenaria uvicola* from Bangalore, India', *Indian Journal of Plant Pathology*, vol. 4, no. 2, pp. 154-61.
- University of Queensland 2011, *Rosella - Hibiscus sabdariffa*, The University of Queensland. Special edition of Environmental Weeds of Australia for Biosecurity Queensland, available at http://keyserver.lucidcentral.org/weeds/data/03030800-0b07-490a-8d04-0605030c0f01/media/Html/Hibiscus_sabdariffa.htm.
- Úrbez-Torres, JR, Adams, P, Kamas, J & Gubler, WD 2009, 'Identification, incidence, and pathogenicity of fungal species associated with grapevine dieback in Texas', *American Journal of Enology and Viticulture*, vol. 60, no. 4, pp. 497-507.
- Úrbez-Torres, JR, Peduto, F, Striegler, RK, Urrea-Romero, KE, Rupe, JC, Cartwright, RD & Gubler, WD 2012, 'Characterization of fungal pathogens associated with grapevine trunk diseases in Arkansas and Missouri', *Fungal Diversity*, vol. 52, pp. 169-89.
- USDA-APHIS 2002, *Importation of grapes (Vitis spp.) from Korea into the United States: a qualitative, pathway-initiated pest risk assessment*, Animal and Plant Health Inspection Service, United States Department of Agriculture, Riverdale, Maryland, available at [https://web01.aphis.usda.gov/oxygen fod/fb_md_ppq.nsf/0/6a4c6eb951006d8685256e2a005eccdc0/\\$FILE/0058.pdf](https://web01.aphis.usda.gov/oxygen fod/fb_md_ppq.nsf/0/6a4c6eb951006d8685256e2a005eccdc0/$FILE/0058.pdf) (pdf 199 kb).
- -- 2004, '7 CFR part 319: importation of fruits and vegetables. Docket number 02-106-2', *Federal Register*, vol. 69 no. 217, pp. 65053-67.
- USDA 2015, *USDA treatment manual*, United States Department of Agriculture, USA, available at http://www.aphis.usda.gov/import_export/plants/manuals/ports/downloads/treatment.pdf (PDF 4.99MB).
- Uyemoto, JK 1975, 'A severe outbreak of virus-induced grapevine decline in cascade grapes in New York', *Plant Disease Reporter*, vol. 59, pp. 98-101.
- -- 1981, 'Tobacco necrosis and satellite viruses', in *Handbook of plant virus infections and comparative diagnosis*, Kurstak, E (ed), Elsevier North-Holland Biomedical Press, Amsterdam.
- Uyemoto, JK & Gilmer, RM 1972, 'Properties of *Tobacco necrosis virus* strains isolated from apple', *Phytopathology*, vol. 62, pp. 478-81.
- Van Niekerk, JM, Groenewald, JZE, Verkley, GJM, Fourie, PH, Wingfield, MJ & Crous, PW 2004, 'Systematic reappraisal of *Coniella* and *Pilidiella*, with specific reference to species occurring on eucalyptus and *Vitis* in South Africa', *Mycological Research*, vol. 108, no. 3, pp. 283-303.

- Vanitha, K, Karuppuchamy, P & Sivasubramanian, P 2011, 'Pests of Vanilla (*Vanilla planifolia* Andrews) and their natural enemies in Tamil Nadu, India', *International Journal of Biodiversity and Conservation*, vol. 3, no. 4, pp. 116-20.
- Varma, RV 1984, 'New records of insects damaging *Eucalyptus* in the nursery in Kerala, India', *Indian Journal of Forestry*, vol. 7, no. 2, pp. 160-1.
- Vawdrey, LL & Peterson, RA 1990, 'Diseases of kenaf (*Hibiscus cannabinus*) in the Burdekin River irrigation area', *Australasian Plant Pathology*, vol. 19, no. 2, pp. 34-5.
- Venette, RC & Davis, EE 2004, *Mini risk assessment: passionvine mealybug: Planococcus minor (Maskell) [Pseudococcidae: Hemiptera]*, University of Minnesota, St Paul, Minnesota, available at http://www.aphis.usda.gov/plant_health/plant_pest_info/pest_detection/downloads/prp/pminorpra.pdf (pdf 7.31 mb).
- Venette, RC, Davis, EE, Heisler, H & Larson, M 2003, *Mini risk assessment: silver Y moth, Autographa gamma (L.) [Lepidoptera: Noctuidae]*, University of Minnesota, St Paul, Minnesota, available at http://www.aphis.usda.gov/plant_health/plant_pest_info/pest_detection/downloads/prp/agammapra.pdf (pdf 4.76 mb).
- Vergheese, A & Harish, R 2010, 'Studies on thrips of grapes in South India, with special reference to *Scirtothrips dorsalis* Hood', *Journal of Insect Science*, vol. 10, no. 166, pp. 52-3.
- Vergheese, A, Madhura, HS, Kamala Jayanthi, PD & Stonehouse, JM 2002, 'Fruit flies of economic significance in India, with special reference to *Bactrocera dorsalis* (Hendel)', *Proceedings of 6th International Fruit Fly Symposium, Stellenbosch, South Africa*, Isteg Scientific Publications, South Africa, pp. 317-24.
- Vergheese, A, Tandon, PL & Stonehouse, JM 2004, 'Economic evaluation of the integrated management of the Oriental fruit fly *Bactrocera dorsalis* (Diptera: Tephritidae) in mango in India', *Crop Protection*, vol. 23, pp. 61-3.
- Verma, AK, Khan, ML & Handa, A 2003, 'Transmission of tomato ringspot virus by *Xiphinema inaequale* (Khan and Ahmad 1975) Bajaj and Jairajpuri 1979, associated with gladiolus in Himachal Pradesh', *Pest Management and Economic Zoology*, vol. 11, no. 2, pp. 189-92.
- Verma, KS, Cheema, SS, Kang, MS & Sharma, AK 1991, 'Hitherto unrecorded disease problems of mango from Punjab', *Plant Disease Research*, vol. 6, no. 2, pp. 141-2.
- Verma, SK, Chand, G & Kumar, P 2012, 'New record of *Icerya purchasi* in one year old seedlings of *Albizia procera* (Roxb.) Benth', *Indian Forester*, vol. 138, no. 2, p. 200.
- ViCSPA 2009, *Certified seed potatoes from Victoria*, available at <http://www.vicspa.org.au/>.
- Villate, L, Fievet, V, Hanse, B, Delemarre, F, Plantard, O, Esmenjaud, D & van Helden, M 2008, 'Spatial distribution of the dagger nematode *Xiphinema index* and its associated *Grapevine fanleaf virus* in french vineyard', *Phytopathology*, vol. 98, no. 8, pp. 942-8.
- VINA 2008, *Vine Industry Nursery Association: accreditation scheme*, available at <http://www.vina.net.au/index.php?pageid=11>.
- Vitale, A & Polizzi, G 2005, 'Disease Notes: Occurrence of *Pestalotiopsis uvicola* causing leaf spots and stem blight on bay laurel (*Laurel nobilis*) in Sicily', *Plant Disease*, vol. 89, p. 1362.
- Wade, NL & Morris, SC 1982, 'Causes and control of cantaloupe postharvest wastage in Australia', *Plant Disease*, vol. 66, pp. 549-52.

- Waite, GK 2009, *Diagnostic methods for Oriental fruit fly: Bactrocera dorsalis*, PaDIL: Pests and Diseases Image Library, available at <http://www.padil.gov.au/pbt>.
- Waldecks 2013, *Fresh fruit trees: how to squeeze fruit trees into home gardens*, Waldecks Garden Centres Care Sheets, available at www.waldecks.com.au/caresheets.htm.
- Walker, K 2008, *Bamboo longhorn beetle: Chlorophorus annularis (Fabricius, 1787) (Coleoptera: Cerambycidae)*, PaDIL: Pests and Diseases Image Library, available at <http://www.padil.gov.au/viewPest.aspx?id=122>.
- -- 2011, *Auger beetle: Xylothrips flavipes*, PaDIL: Pests and Diseases Image Library, available at <http://www.padil.gov.au/pests-and-diseases/Pest/Main/135586>.
- Walter, DE, Halliday, RB & Smith, D 1995, 'The Oriental red mite, *Eutetranychus orientalis* (Klein) (Acarina: Tetranychidae), in Australia', *Journal of the Australian Entomological Society*, vol. 34, no. 4, pp. 307-8.
- Walton, VM, Daane, KM, Bentley, WJ, Millar, JG & Larsen, TE 2006, 'Pheromone-based mating disruption of *Planococcus ficus* (Hemiptera: Pseudococcidae) in California vineyards', *Journal of Economic Entomology*, vol. 99, no. 4, pp. 1280-90.
- Walton, VM & Pringle, KL 2004a, 'Seasonal population studies of vine mealybug, *Planococcus ficus* (Signoret), and its natural enemies in vineyards in the Western Cape Province, South Africa', *California Conference on Biological Control IV, Berkeley, California*, University of California, Berkeley, pp. 150-2.
- -- 2004b, 'Vine mealybug, *Planococcus ficus* (Signoret) (Hemiptera: Pseudococcidae), a key pest in South African vineyards: a review', *South African Journal of Enology and Viticulture*, vol. 25, no. 2, pp. 54-62.
- Wan Chow Wah, YF & Symons, RH 1997, 'A high sensitivity RT-PCR assay for the diagnosis of viroids in grapevines in the field and in tissue culture', *Journal of Virological Methods*, vol. 63, pp. 57-69.
- -- 1999, 'Transmission of viroids via grape seeds', *Journal of Phytopathology*, vol. 147, pp. 285-91.
- Wangikar, J, Raut, G & Gopalkrishna, N 1969, 'Drying of grape vines caused by *Hendersonula toruloidea*', *Indian Phytopathology*, vol. 22, pp. 403-4.
- Ward, D 2001, *Control of European wasps in vineyards*, Under Control: pest plant and animal management news, Number 16, State of Victoria, Department of natural resources and environment.
- Washington State University 2008, *Noctuid pests in Washington orchards: spotted cutworm (Xestia c-nigrum)*, Tree Fruit Research and Extension Center, Washington State University, available at <http://entomology.tfrec.wsu.edu/noctuid/id.html>.
- Washington, WS & Nancarrow, RJ 1983, *List of diseases recorded on fruit and vegetable crops in Victoria before June 30, 1980*, Technical report series, Report No. 66, Victorian Department of Agriculture, Melbourne.
- Waterhouse, DF 1993, *The major arthropod pests and weeds of agriculture in southeast Asia: distribution, importance and origin*, ACIAR, Canberra.
- Watson, JR 1923, 'Insects of the grape', *Proceedings of the Florida State Horticultural Society*, vol. 36, pp. 104-8.

- Watson, RN 2004, 'Internal biosecurity: a realistic objective for plant nematodes?', *New Zealand Plant Protection*, vol. 57, pp. 151-5.
- Weinert, MP, Shivas, RG, Pitkethley, RN & Daly, AM 2003, 'First record of grapevine leaf rust in the Northern Territory, Australia', *Australasian Plant Pathology*, vol. 32, no. 1, pp. 117-8.
- Weintraub, PG & Beanland, L 2006, 'Insect vectors of phytoplasmas: supplemental material', *Annual Review of Entomology*, vol. 51, pp. 91-111.
- Weintraub, PG & Jones, P, (eds) 2010, *Phytoplasmas: genomes, plant hosts and vectors*, CAB International, Wallingford.
- White, IM & Elson-Harris, MM 1992, *Fruit flies of economic significance: their identification and bionomics*, CAB International, Wallingford, UK.
- Wilcox, WF 2003, *Grapes: black rot (Guignardia bidwellii (Ellis) Viala and Ravaz)*, available at http://www.nysipm.cornell.edu/factsheets/grapes/diseases/grape_br.pdf (pdf 413 kb).
- Wilkinson, KG, Sivasithamparam, K, Dixon, KW, Fahy, PC & Bradley, JK 1994, 'Identification and characterisation of bacteria associated with Western Australian orchids', *Soil Biology and Biochemistry*, vol. 26, no. 1, pp. 137-42.
- Williams, DJ 1989, 'The mealybug genus *Rastrococcus* Ferris (Hemiptera: Pseudococcidae)', *Systematic Entomology*, vol. 14, no. 4, pp. 433-86.
- -- 2004, *Mealybugs of Southern Asia*, Natural History Museum and Southdene, Kuala Lumpur.
- Wilson, MR & Turner, JA 2010, *Leafhopper, planthopper and psyllid vectors of plant disease*, Amgueddfa Cymru - National Museum of Wales, available at <http://naturalhistory.museumwales.ac.uk/Vectors>.
- Woodruff, RE, Gerberg, EJ & Spilman, TJ 2011, *A false powder-post beetle, *Xylopsocus capucinus* (Fabridius) (Insecta: Coleoptera: Bostrichidae)*, University of Florida, IFAS Extension, available at <http://edis.ifas.ufl.edu/in336>.
- WSUE 2010, *SWD Interest Group, 2010-10-01 09:46:00. WSU Extension*, Washington State University, available at <http://lyris.cahe.wsu.edu/read/?forum=swdinterestgroup&sb=1>.
- WTO 1995, 'Agreement on the application of sanitary and phytosanitary measures', World Trade Organization. Geneva.
- Wu, T & Lo, K 1989, 'Integrated control of citrus red spider mite, *Panonychus citri*, (Acari: Tetranychidae)' (in Chinese), *Chinese Journal of Entomology*, vol. 3, pp. 139-43.
- Wu, YF 2005, 'Geographical distribution model of Oriental fruit fly', *Journal of Fujian Agricultural University*, vol. 34, no. 2, pp. 168-71.
- Wysoki, M 1986, 'New records of Lepidopterous pests of macadamia in Israel', *Phytoparasitica*, vol. 14, no. 2, p. 147.
- Wysoki, M, Yehuda, SB & Rosen, D 1993, 'Reproductive behavior of the honeydew moth, *Cryptoblabes gnidiella*', *Invertebrate Reproduction and Development*, vol. 24, no. 3, pp. 217-24.
- Xi, D, Li, J, Han, C, Li, D, Yu, J & Zhou, X 2008, 'Complete nucleotide sequence of a new strain of *Tobacco necrosis virus A* infecting soybean in China and infectivity of its full-length cDNA clone', *Virus Genes*, vol. 36, pp. 259-66.

- Xu, L, Kusakari, S, Hosomi, A, Toyoda, H & Ouchi, S 1999, 'Postharvest disease of grape caused by *Pestalotiopsis* species' (in Japanese), *Annals of the Phytopathological Society of Japan*, vol. 65, pp. 305-11.
- Yaguchi, S & Takahashi, T 1984, 'Survival of *Hop stunt viroid* in the hop garden', *Phytopathologische Zeitschrift*, vol. 109, pp. 32-44.
- Yamato, H 1995, 'White rot of grape caused by *Coniella* spp.' (in Japanese), *Annals of the Phytopathological Society of Japan*, vol. 61, p. 635.
- Yano, K 1963, 'Taxonomic and biological studies of Pterophoridae of Japan (Lepidoptera)', *Pacific Insects*, vol. 5, no. 1, pp. 65-209.
- Ye, H & Liu, JH 2005, 'Population dynamics of the Oriental fruit fly, *Bactrocera dorsalis* (Diptera: Tephritidae) in the Kunming area, Southwestern China', *Insect Science*, vol. 12, pp. 387-92.
- Yehuda, SB, Wysoki, M & Rosen, D 1991, 'Phenology of the honeydew moth, *Cryptoblabes gnidiella* (Milliere) (Lepidoptera: Pyralidae), on avocado in Israel', *Israel Journal of Entomology*, vol. 25-26, pp. 149-60.
- Yokoyama, VY & Miller, GT 2000, 'Response of omnivorous leafroller (Lepidoptera: Tortricidae) and onion thrips (Thysanoptera: Thripidae) to low-temperature storage', *Journal of Economic Entomology*, vol. 93, no. 3, pp. 1031-4.
- Zalom, FG, Varela, LG & Cooper, M 2011, *European grapevine moth (Lobesia botrana): provisional guidelines*, UC IPM Online: Statewide Integrated Pest Management Program, available at <http://www.ipm.ucdavis.edu/EXOTIC/eurograpevinemoth.html#>.
- Zaspel, JM & Branham, MA 2008, 'World checklist of tribe Calpini (Lepidoptera: Noctuidae: Calpinae)', *Insecta Mundi*, vol. 47, pp. 1-15.
- Zhang, BC 1994, *Index of economically important Lepidoptera*, CAB International, Wallingford, UK.
- Zhang, L, French, R & Langenberg, WG 1993, 'Molecular cloning and sequencing of the coat protein gene of a Nebraskan isolate of *Tobacco necrosis virus*: the deduced coat protein sequence has only moderate homology with those of strain A and strain D', *Archives of Virology*, vol. 132, pp. 291-305.
- Zhang, YP 2005, *Identification and control of pests and diseases of table grape (with colour plates)*, Jindun, Beijing.
- Zhou, X 2002, *Broad bean wilt virus 2*, Descriptions of Plant Viruses, available at <http://www.dpvweb.net/dpv/showdpv.php?dpvno=392>.
- Zimmerman, EC 1958, *Insects of Hawaii Volume 8, Lepidoptera: Pyraloidea*, University of Hawaii Press, Honolulu.
- Zitikaite, I & Staniulis, J 2009, 'Isolation and characterization of *Tobacco necrosis virus* detected on some vegetable species', *Biologija*, vol. 55, pp. 35-9.
- Zohary, D 1996, 'The domestication of the grapevine *Vitis vinifera* L. in the Near East', in *The origins and ancient history of wine*, McGovern, PE, Fleming, SJ & Katz, SH (eds), Gordon and Breach Publishers, London.